




**MATS**  
UNIVERSITY

NAAC  
GRADE **A<sup>+</sup>**  
ACCREDITED UNIVERSITY

# **MATS CENTRE FOR DISTANCE & ONLINE EDUCATION**

## **Differential Equations**

**Master of Science (M.Sc.)  
Semester - 1**



**SELF LEARNING MATERIAL**



**MATS UNIVERSITY**

[www.matsuniversity.ac.in](http://www.matsuniversity.ac.in)

**NAAC**  
**GRADE A<sup>+</sup>**  
ACCREDITED UNIVERSITY

## MSCMODL103 DIFFERENTIAL EQUATIONS

<b>Block-1</b>	
<b>Unit-1:</b>	
Linear equations with constant coefficients	1-5
<b>Unit-2:</b>	
The second order homogeneous equations – Initial value problems	6-27
<b>Unit-3:</b>	
Linear dependence and independence - A formula for the Wronskian – The non- homogeneous equation of order two.	28-54
<b>Block-2</b>	
<b>Unit-4:</b>	
Homogeneous and non-homogeneous equations of order n	55-59
<b>Unit-5:</b>	
Initial value problems – Annihilator method to solve a non-homogeneous equation –	60-73
<b>Unit-6:</b>	
Algebra of constant coefficient operators.	74-97
<b>Block-3</b>	
<b>Unit-7:</b>	
Linear equations with variable coefficients: initial value problems for the homogeneous equation	98-108
<b>Unit-8:</b>	
Solutions of the homogeneous equation	109-112
<b>Unit-9:</b>	

The Wronskian and linear independence – Reduction of the order of a homogeneous equation	113-124
<b>Unit-10:</b>	
Homogeneous equation with analytic coefficients – The Legendre equation.	125-151
<b>Block-4</b>	
<b>Unit-11:</b>	
Linear equation with regular singular points: Euler equation	152-158
<b>Unit-12:</b>	
Second order equations with regular singular points	159-179
<b>Unit-13:</b>	
Exceptional cases – Bessel equation.	180-196
<b>Block-5</b>	
<b>Unit-14:</b>	
Existence and uniqueness of solutions to first order equations	197-208
<b>Unit-15:</b>	
Equation with variables separated– Exact equations	209-216
<b>Unit-16:</b>	
The method of successive approximations – The Lipschitz condition	217-226
<b>Unit-17:</b>	
Convergence of the successive approximations.	227-238

---

#### COURSE DEVELOPMENT EXPERT COMMITTEE

---

Prof (Dr) K P Yadav

Vice Chancellor, MATS University

Prof (Dr) A J Khan

Professor Mathematics, MATS University

Prof(Dr) D K Das

Professor Mathematics, CCET, Bhilai

---

#### COURSE COORDINATOR

---

Dr Vinita Dewangan

Associate Professor, MATS University

---

#### COURSE /BLOCK PREPARATION

---

Prof (Dr.) A. J. Khan,

Professor, MATS University

---

March 2025

ISBN: 978-93-49954-75-5

@MATS Centre for Distance and Online Education, MATS University, Village- Gullu, Aarang, Raipur-  
(Chhattisgarh)

All rights reserved. No part of this work may be reproduced or transmitted or utilized or stored in any form,  
by mimeograph or any other means, without permission in writing from MATS University, Village- Gullu,  
Aarang, Raipur-(Chhattisgarh)

Printed & Published on behalf of MATS University, Village-Gullu, Aarang, Raipur by Mr. Meghanadhu  
Katabathuni, Facilities & Operations, MATS University, Raipur (C.G.)

---

Disclaimer-Publisher of this printing material is not responsible for any error or dispute from contents of  
this course material, this is completely depends on AUTHOR'S MANUSCRIPT.

Printed at: The Digital Press, Krishna Complex, Raipur-492001(Chhattisgarh)

## Acknowledgement

The material (pictures and passages) we have used is purely for educational purposes. Every effort has been made to trace the copyright holders of material reproduced in this book. Should any infringement have occurred, the publishers and editors apologize and will be pleased to make the necessary corrections in future editions of this book.

---

## COURSE INTRODUCTION

---

Differential equations play a fundamental role in mathematical modeling, physics, engineering, and other applied sciences. This course covers various types of differential equations, including linear and non-linear equations, initial value problems, and equations with variable coefficients and singular points. Students will explore theoretical concepts as well as practical methods for solving differential equations.

### **Block 1: Linear Equations with Constant Coefficients**

This module introduces linear differential equations with constant coefficients, covering second-order homogeneous equations and initial value problems. Topics include linear dependence and independence, the Wronskian, and non-homogeneous equations of order two.

### **Block 2: Higher-Order Equations and Non-Homogeneous Solutions**

This module covers homogeneous and non-homogeneous equations of order  $n$ , including initial value problems. Students will explore the annihilator method for solving non-homogeneous equations and the algebra of constant coefficient operators.

### **Block 3: Linear Equations with Variable Coefficients**

This module focuses on linear differential equations with variable coefficients, including initial value problems for homogeneous equations. Topics include solutions of homogeneous equations, the Wronskian and linear independence, reduction of order techniques, and the Legendre equation.

### **Block 4: Equations with Singular Points**

Students will study linear equations with regular singular points, including Euler equations and second-order equations with singular points. The module also covers exceptional cases and introduces the Bessel equation, which appears in many physical applications.

### **Block 5: Existence and Uniqueness of Solutions**

This module explores fundamental existence and uniqueness theorems for first-order differential equations. Topics include equations with

variables separated, exact equations, the method of successive approximations, the Lipschitz condition, and convergence of successive approximations.

# BLOCK 1

## UNIT 1

### LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

#### Objectives

- Understand second-order homogeneous linear differential equations.
- Learn to solve initial value problems.
- Study the concepts of linear dependence and independence of solutions.
- Derive and use a formula for the Wronskian.
- Solve non-homogeneous differential equations of order two.

#### 1.1 Introduction to Linear Equations with Constant Coefficients

Linear differential equations are among the most important types of differential equations in mathematics and its applications. A linear differential equation with constant coefficients has the form:

$$a_n \left( \frac{d^n y}{dx^n} \right) + a_{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \dots + a_1 \frac{dy}{dx} + a_0 y = g(x)$$

where  $a_0, a_1, \dots, a_n$  are constants and  $g(x)$  is a function of  $x$ .

When  $g(x) = 0$ , the equation is called homogeneous. Otherwise, it's called non-homogeneous.

#### Key Properties of Linear Equations

1. **Superposition Principle:** If  $y_1$  and  $y_2$  are solutions to a homogeneous linear equation, then any linear combination  $c_1 y_1 + c_2 y_2$  is also a solution.
2. **General Solution Structure:** The general solution to a non-homogeneous equation consists of the general solution to the corresponding homogeneous equation plus any particular solution to the non-homogeneous equation.
3. **Existence and Uniqueness:** For an  $n$ th-order linear equation, a unique solution exists when  $n$  initial conditions are specified.



## First-Order Linear Equations

The simplest linear differential equation with constant coefficients is the first-order equation:

$$\frac{dy}{dx} + ay = g(x)$$

where  $a$  is a constant and  $g(x)$  is a function of  $x$ .

The general solution to the homogeneous equation  $\frac{dy}{dx} + ay = 0$  is:

$$y = Ce^{-ax}$$

where  $C$  is an arbitrary constant.

For the non-homogeneous equation, we can use the method of integrating factors. Multiplying both sides by  $e^{ax}$ :

$$e^{ax}\left(\frac{dy}{dx}\right) + ae^{ax}y = e^{ax}g(x)$$

The left side can be rewritten as:

$$\frac{d}{dx}(e^{ax}y) = e^{ax}g(x)$$

Integrating both sides:

$$e^{ax}y = \int e^{ax}g(x)dx + C$$

Therefore:

$$y = e^{-ax}\left[\int e^{ax}g(x)dx + C\right]$$

### Example 1.1

Solve the differential equation:  $\frac{dy}{dx} + 2y = 4x$

Solution: This is a first-order linear equation with  $a = 2$  and  $g(x) = 4x$ .

Using the method of integrating factors, the integrating factor is  $e^{ax} = e^{2x}$

Multiplying both sides by  $e^{2x}$ :  $e^{2x}\left(\frac{dy}{dx}\right) + 2e^{2x}y = 4xe^{2x}$

This can be rewritten as:  $\frac{d}{dx}(e^{2x}y) = 4xe^{2x}$

Integrating both sides:  $e^{(2x)}y = \int 4xe^{(2x)}dx$

To evaluate the integral, we use integration by parts:  $\int 4xe^{(2x)}dx = 4 \frac{xe^{(2x)}}{2} -$

$$\int \frac{e^{(2x)}}{2} dx = 2xe^{(2x)} - e^{(2x)} + C$$

Therefore:  $e^{2x}y = 2xe^{(2x)} - e^{(2x)} + C$

Solving for y:  $y = 2x - 1 + Ce^{(-2x)}$

This is the general solution to the given differential equation.

### CHECK YOUR PROGRESS

- 1) Write the auxiliary equation corresponding to  $y'' - 5y' + 6y = 0$

.....  
.....  
.....

- 2) What form does the solution take when the auxiliary equation has complex roots?

.....  
.....  
.....

- 3) Explain how repeated roots affect the complementary function.

.....  
.....  
.....

### LET US SUM UP

A linear differential equation with constant coefficients is one in which the coefficients of the derivatives remain constant throughout the equation. It is generally expressed as  $a_n d^n y/dx^n + a_{n-1} d^{n-1} y/dx^{n-1} + \dots + a_1 dy/dx + a_0 y = 0$ . To solve such equations, we replace the derivatives by powers of  $r$  to obtain the auxiliary equation. The nature of the roots of this equation is real, repeated, or complex, determines the form of the complementary function.

## **UNIT END EXERCISES**

### **Short Answer Questions:**

1. Define a second-order homogeneous linear differential equation.
2. What is the significance of the Wronskian in determining linear dependence?
3. What is the characteristic equation of a linear differential equation?

### **Long Answer Questions:**

1. Derive and explain the characteristic equation for a second-order linear differential equation.
2. Explain the role of initial conditions in solving differential equations.
3. Prove that if the Wronskian of two functions is nonzero, the functions are linearly independent.

### **REFERENCES AND SUGGESTED READINGS:**

1. Boyce, W. E., & DiPrima, R. C. (2021). Elementary Differential Equations and Boundary Value Problems (11th ed.). Wiley.
2. Zill, D. G. (2017). A First Course in Differential Equations with Modeling Applications (11th ed.). Cengage Learning.
3. Edwards, C. H., & Penney, D. E. (2018). Differential Equations and Boundary Value Problems: Computing and Modeling (5th ed.). Pearson.
4. Ross, S. L. (2013). Differential Equations (4th ed.). Wiley.

## UNIT 2

### THE SECOND ORDER HOMOGENEOUS EQUATIONS – INITIAL VALUE PROBLEMS

#### 2.1 Second-Order Homogeneous Equations

Second-order linear homogeneous differential equations with constant coefficients have the form:

$$a\left(\frac{d^2y}{dx^2}\right) + b\left(\frac{dy}{dx}\right) + cy = 0$$

where  $a$ ,  $b$ , and  $c$  are constants, and  $a \neq 0$ .

#### The Characteristic Equation

To solve such equations, we use the characteristic equation:

$$ar^2 + br + c = 0$$

The solutions to this quadratic equation determine the form of the general solution to the differential equation.

#### Case 1: Distinct Real Roots

If the characteristic equation has two distinct real roots  $r_1$  and  $r_2$ , then the general solution is:

$$y = C_1e^{(r_1x)} + C_2e^{(r_2x)}$$

where  $C_1$  and  $C_2$  are arbitrary constants.

#### Case 2: Repeated Root

If the characteristic equation has a repeated root  $r$ , then the general solution is:

$$y = C_1e^{(rx)} + C_2xe^{(rx)}$$

#### Case 3: Complex Conjugate Roots

If the characteristic equation has complex conjugate roots  $r = \alpha \pm \beta i$ , then the general solution is:

$$y = e^{(\alpha x)}[C_1\cos(\beta x) + C_2\sin(\beta x)]$$

#### Example 1

Solve the differential equation:  $(\frac{d^2y}{dx^2}) - 5\frac{dy}{dx} + 6y = 0$

Solution: This is a second-order linear homogeneous equation with  $a = 1$ ,  $b = -5$ , and  $c = 6$ .

The characteristic equation is:  $r^2 - 5r + 6 = 0$

Factoring this equation:  $(r - 2)(r - 3) = 0$

The roots are  $r_1 = 2$  and  $r_2 = 3$ .

Since we have distinct real roots, the general solution is:  $y = C_1e^{(2x)} + C_2e^{(3x)}$

### Example 2

Solve the differential equation:  $(\frac{d^2y}{dx^2}) + 4\frac{dy}{dx} + 4y = 0$

Solution: This is a second-order linear homogeneous equation with  $a = 1$ ,  $b = 4$ , and  $c = 4$ .

The characteristic equation is:  $r^2 + 4r + 4 = 0$

This can be rewritten as:  $(r + 2)^2 = 0$

The equation has a repeated root  $r = -2$ .

Therefore, the general solution is:  $y = C_1e^{(-2x)} + C_2xe^{(-2x)}$

### Example 3

Solve the differential equation:  $(\frac{d^2y}{dx^2}) + 4y = 0$

Solution: This is a second-order linear homogeneous equation with  $a = 1$ ,  $b = 0$ , and  $c = 4$ .

The characteristic equation is:  $r^2 + 4 = 0$

The roots are:  $r = \pm 2i$

Since we have complex conjugate roots with  $\alpha = 0$  and  $\beta = 2$ , the general solution is:  $y = C_1\cos(2x) + C_2\sin(2x)$

## 2.2. Initial Value Problems for Second-Order Equations

An initial value problem for a second-order differential equation consists of the equation itself along with two initial conditions:

$$a \left( \frac{d^2 y}{dx^2} \right) + b \frac{dy}{dx} + cy = g(x) \quad y(x_0) = y_0 \quad y'(x_0) = y_1$$

where  $y_0$  and  $y_1$  are given values, and  $x_0$  is the initial point.

### Solving Initial Value Problems

To solve an initial value problem:

1. Find the general solution to the differential equation.
2. Apply the initial conditions to determine the values of the arbitrary constants.

#### Example 4

Solve the initial value problem:  $\left( \frac{d^2 y}{dx^2} \right) - 3 \frac{dy}{dx} + 2y = 0 \quad y(0) = 1 \quad y'(0) = 0$

Solution: First, we find the general solution to the differential equation.

The characteristic equation is:  $r^2 - 3r + 2 = 0$

Factoring:  $(r - 1)(r - 2) = 0$

The roots are  $r_1 = 1$  and  $r_2 = 2$ .

Therefore, the general solution is:  $y = C_1 e^x + C_2 e^{(2x)}$

Now, we apply the initial conditions:

From  $y(0) = 1$ :  $y(0) = C_1 e^0 + C_2 e^0 = C_1 + C_2 = 1$

From  $y'(0) = 0$ :  $y'(x) = C_1 e^x + 2C_2 e^{(2x)} \quad y'(0) = C_1 + 2C_2 = 0$

We now have the system of equations:  $C_1 + C_2 = 1 \quad C_1 + 2C_2 = 0$

Subtracting the second equation from the first:  $-C_2 = 1 \quad C_2 = -1$

Substituting back:  $C_1 + (-1) = 1 \quad C_1 = 2$

Therefore, the solution to the initial value problem is:  $y = 2e^x - e^{(2x)}$

### Non-Homogeneous Equations

For non-homogeneous second-order linear equations:

$$a \left( \frac{d^2 y}{dx^2} \right) + b \frac{dy}{dx} + cy = g(x)$$

The general solution has the form:

$$y = y_h + y_p$$

where  $y_h$  is the general solution to the corresponding homogeneous equation, and  $y_p$  is a particular solution to the non-homogeneous equation.

### 2.3 Methods for Finding Particular Solutions

1. **Method of Undetermined Coefficients:** This method works when  $g(x)$  is a polynomial, exponential, sine, cosine, or a linear combination of these.
2. **Variation of Parameters:** This is a more general method that can be used for any continuous function  $g(x)$ .

#### Method of Undetermined Coefficients

The form of the particular solution depends on the form of  $g(x)$ :

- If  $g(x) = P_n(x)$  (a polynomial of degree  $n$ ), then  $y_p = Q_n(x)$  (a polynomial of degree  $n$ ).
- If  $g(x) = e^{(ax)}$ , then  $y_p = Ae^{(ax)}$ , where  $A$  is a constant.
- If  $g(x) = \cos(\beta x)$  or  $g(x) = \sin(\beta x)$ , then  $y_p = A \cos(\beta x) + B \sin(\beta x)$ .

If the form of  $y_p$  is already a solution to the homogeneous equation, we multiply by  $x$  (or  $x^2$  if necessary) to ensure linear independence.

#### Variation of Parameters

For the equation  $a \left( \frac{d^2 y}{dx^2} \right) + b \frac{dy}{dx} + cy = g(x)$ , if  $y_1$  and  $y_2$  are two linearly independent solutions to the homogeneous equation, then a particular solution can be found as:

$$y_p = -y_1 \int (y_2 g(x) / W(y_1, y_2)) dx + y_2 \int (y_1 g(x) / W(y_1, y_2)) dx$$

where  $W(y_1, y_2) = y_1 y_2' - y_1' y_2$  is the Wronskian.

#### Solved Problems

##### Problem 1

Solve the differential equation:  $\left( \frac{d^2 y}{dx^2} \right) + y = 0$

Solution: This is a second-order linear homogeneous equation with  $a = 1$ ,  $b = 0$ , and  $c = 1$ .

The characteristic equation is:  $r^2 + 1 = 0$

The roots are:  $r = \pm i$



Since we have complex conjugate roots with  $\alpha = 0$  and  $\beta = 1$ , the general solution is:  $y = C_1 \cos(x) + C_2 \sin(x)$

### Problem 2

Solve the differential equation:  $(\frac{d^2 y}{dx^2}) - 4\frac{dy}{dx} + 4y = 0$

Solution: This is a second-order linear homogeneous equation with  $a = 1$ ,  $b = -4$ , and  $c = 4$ .

The characteristic equation is:  $r^2 - 4r + 4 = 0$

This can be rewritten as:  $(r - 2)^2 = 0$

The equation has a repeated root  $r = 2$ .

Therefore, the general solution is:  $y = C_1 e^{(2x)} + C_2 x e^{(2x)}$

### Problem 3

Solve the differential equation:  $(\frac{d^2 y}{dx^2}) - y = 0$

Solution: This is a second-order linear homogeneous equation with  $a = 1$ ,  $b = 0$ , and  $c = -1$ .

The characteristic equation is:  $r^2 - 1 = 0$

Factoring:  $(r - 1)(r + 1) = 0$

The roots are  $r_1 = 1$  and  $r_2 = -1$ .

Therefore, the general solution is:  $y = C_1 e^x + C_2 e^{(-x)}$

### Problem 4

Solve the differential equation:  $(\frac{d^2 y}{dx^2}) + 6\frac{dy}{dx} + 9y = 0$

Solution: This is a second-order linear homogeneous equation with  $a = 1$ ,  $b = 6$ , and  $c = 9$ .

The characteristic equation is:  $r^2 + 6r + 9 = 0$

This can be rewritten as:  $(r + 3)^2 = 0$

The equation has a repeated root  $r = -3$ .

Therefore, the general solution is:  $y = C_1 e^{(-3x)} + C_2 x e^{(-3x)}$

### Problem 5

Solve the initial value problem:  $(\frac{d^2y}{dx^2}) + 9y = 0$   $y(0) = 2$   $y'(0) = 3$

Solution: First, we find the general solution to the differential equation.

The characteristic equation is:  $r^2 + 9 = 0$

The roots are:  $r = \pm 3i$

Since we have complex conjugate roots with  $\alpha = 0$  and  $\beta = 3$ , the general solution is:  $y = C_1 \cos(3x) + C_2 \sin(3x)$

Now, we apply the initial conditions:

From  $y(0) = 2$ :  $y(0) = C_1 \cos(0) + C_2 \sin(0) = C_1 = 2$

From  $y'(0) = 3$ :  $y'(x) = -3C_1 \sin(3x) + 3C_2 \cos(3x)$   $y'(0) = 3C_2 = 3$   $C_2 = 1$

Therefore, the solution to the initial value problem is:  $y = 2\cos(3x) + \sin(3x)$

### Unsolved Problems

#### Problem 1

Solve the differential equation:  $(\frac{d^2y}{dx^2}) - 2\frac{dy}{dx} - 3y = 0$

#### Problem 2

Solve the differential equation:  $(\frac{d^2y}{dx^2}) + 2\frac{dy}{dx} + 5y = 0$

#### Problem 3

Solve the initial value problem:  $(\frac{d^2y}{dx^2}) - 4y = 0$   $y(0) = 1$   $y'(0) = 2$

#### Problem 4

Solve the differential equation:  $(\frac{d^2y}{dx^2}) + 4\frac{dy}{dx} + 5y = 0$

#### Problem 5

Solve the initial value problem:  $(\frac{d^2y}{dx^2}) - 6\frac{dy}{dx} + 9y = 0$   $y(0) = 0$   $y'(0) = 1$

### 2.4 Applications of Linear Differential Equations

Linear differential equations with constant coefficients appear in many applications:

1. **Mechanical Systems:** The motion of a mass-spring system is governed by a second-order linear differential equation.

2. **Electrical Circuits:** The behavior of RLC circuits can be modeled using second-order linear differential equations.
3. **Vibrations:** The vibrations of strings, membranes, and other mechanical systems are described by linear differential equations.
4. **Heat Conduction:** The diffusion of heat in a medium follows a linear partial differential equation.
5. **Population Dynamics:** In some cases, population growth can be modeled using linear differential equations.

### Mass-Spring Systems

A mass attached to a spring is a classic example of a system modeled by a second-order linear differential equation. If the mass is  $m$ , the spring constant is  $k$ , and the damping coefficient is  $c$ , then the equation of motion is:

$$m \frac{d^2x}{dt^2} + c \frac{dx}{dt} + kx = F(t)$$

where  $x$  is the displacement from equilibrium and  $F(t)$  is an external force.

When  $F(t) = 0$ , the equation becomes:

$$m \frac{d^2x}{dt^2} + c \frac{dx}{dt} + kx = 0$$

This is a homogeneous second-order linear equation with constant coefficients. The behavior of the system depends on the values of  $m$ ,  $c$ , and  $k$ :

1. **Underdamped ( $c^2 < 4mk$ ):** The system oscillates with decreasing amplitude.
2. **Critically Damped ( $c^2 = 4mk$ ):** The system returns to equilibrium without oscillation, in the shortest possible time.
3. **Overdamped ( $c^2 > 4mk$ ):** The system returns to equilibrium without oscillation, but more slowly than in the critically damped case.

### Electrical Circuits

An RLC circuit consisting of a resistor (R), an inductor (L), and a capacitor (C) in series can be modeled by the equation:

$$L(d^2q/dt^2) + R(dq/dt) + (1/C)q = E(t)$$

where  $q$  is the charge on the capacitor and  $E(t)$  is the electromotive force.

When  $E(t) = 0$ , the equation becomes:

$$L(d^2q/dt^2) + R(dq/dt) + (1/C)q = 0$$

This is the same form as the mass-spring system, and the behavior is similarly classified as underdamped, critically damped, or overdamped.

### Higher-Order Linear Equations

The methods discussed for second-order equations can be extended to higher-order linear equations with constant coefficients:

$$a_n \frac{d^n y}{dx^n} + a_{(n-1)} \frac{d^{(n-1)} y}{dx^{n-1}} + \dots + a_1 \frac{dy}{dx} + a_0 y = 0$$

The characteristic equation becomes:

$$a_n r^n + a_{(n-1)} r^{(n-1)} + \dots + a_1 r + a_0 = 0$$

The general solution depends on the roots of this equation:

1. For each distinct real root  $r_i$ , there is a term  $C_i e^{(r_i x)}$  in the general solution.
2. For each repeated real root  $r_i$  with multiplicity  $m$ , there are terms  $C_1 e^{(r_i x)}, C_2 x e^{(r_i x)}, \dots, C_m x^{(m-1)} e^{(r_i x)}$  in the general solution.
3. For each pair of complex conjugate roots  $\alpha \pm \beta i$ , there are terms  $e^{(\alpha x)} [C_1 \cos(\beta x) + C_2 \sin(\beta x)]$  in the general solution.

### Systems of Linear Differential Equations

Many problems in physics, engineering, and other fields lead to systems of linear differential equations with constant coefficients:

$$\frac{dx}{dt} = ax + by \quad \frac{dy}{dt} = cx + dy$$

where  $a, b, c$ , and  $d$  are constants.

Such systems can be written in matrix form:

$$\frac{d}{dt} [x, y]^T = A [x, y]^T$$

where  $A$  is the coefficient matrix.

The solution involves finding the eigenvalues and eigenvectors of  $A$ . If  $\lambda$  is an eigenvalue and  $v$  is the corresponding eigenvector, then  $e(\lambda t)v$  is a solution to the system.

Linear differential equations with constant coefficients form a fundamental class of differential equations with wide-ranging applications. The methods for solving these equations, particularly the use of the characteristic equation, provide a systematic approach to finding the general solution. Initial value problems can then be solved by applying the given initial conditions to determine the arbitrary constants in the general solution. For non-homogeneous equations, the method of undetermined coefficients and the variation of parameters provide techniques for finding particular solutions. The general solution is then the sum of the homogeneous solution and the particular solution. Higher-order equations and systems of equations follow similar principles, with the complexity increasing as the order or the number of equations increases. However, the underlying framework remains the same: find the general solution and then apply the given conditions to determine the arbitrary constants.

## 2.5. Linear Dependence and Independence of Solutions

### Fundamental Concepts

When solving higher-order differential equations, we often find multiple solutions. Understanding the relationships between these solutions is crucial for constructing general solutions. This is where the concepts of linear dependence and independence come into play.

### Definition of Linear Dependence

A set of functions  $\{y_1(x), y_2(x), \dots, y_n(x)\}$  defined on an interval  $I$  is said to be **linearly dependent** if there exist constants  $c_1, c_2, \dots, c_n$ , not all zero, such that:

$$c_1 y_1(x) + c_2 y_2(x) + \dots + c_n y_n(x) = 0$$

for all  $x$  in the interval  $I$ .

In simpler terms, if one function can be expressed as a linear combination of the others, the set is linearly dependent.

### Definition of Linear Independence

A set of functions  $\{y_1(x), y_2(x), \dots, y_n(x)\}$  is **linearly independent** on an interval  $I$  if the only solution to:

$$c_1 y_1(x) + c_2 y_2(x) + \dots + c_n y_n(x) = 0$$

for all  $x$  in  $I$ , is  $c_1 = c_2 = \dots = c_n = 0$ .

In other words, no function in the set can be expressed as a linear combination of the others.

### Importance in Differential Equations

For an  $n$ th-order linear homogeneous differential equation, the general solution is a linear combination of  $n$  linearly independent particular solutions:

$$y = c_1 y_1(x) + c_2 y_2(x) + \dots + c_n y_n(x)$$

where  $y_1(x), y_2(x), \dots, y_n(x)$  form a fundamental set of solutions.

### Testing for Linear Independence

There are several ways to test whether a set of functions is linearly independent:

1. **Direct Method:** Check if one function can be written as a linear combination of others.
2. **Using the Wronskian** (more details in the next section).
3. **Using properties of solutions to differential equations.**

### Example of Linear Dependence

Consider the functions:

- $y_1(x) = e^x$
- $y_2(x) = e^x$
- $y_3(x) = 2e^x$

These functions are linearly dependent because:  $y_3(x) = 2y_1(x)$  or equivalently  $y_1(x) - y_2(x) + y_3(x)/2 = 0$

### Example of Linear Independence

Consider the functions:

- $y_1(x) = e^x$
- $y_2(x) = e^{2x}$
- $y_3(x) = e^{3x}$

These functions are linearly independent because no non-trivial linear combination of them equals zero for all  $x$ .

### Fundamental Theorem

For a linear homogeneous differential equation of order  $n$ :

$$a_0(x)y^{(n)} + a_1(x)y^{(n-1)} + \dots + a_{n-1}(x)y' + a_n(x)y = 0$$

with  $a_0(x) \neq 0$  on an interval  $I$ , there exists exactly  $n$  linearly independent solutions on  $I$ . Any solution can be expressed as a linear combination of these  $n$  fundamental solutions.

## 2.6. The Wronskian: Definition and Applications

### Definition of the Wronskian

The Wronskian is a powerful tool for determining whether a set of functions is linearly independent.

For functions  $y_1(x), y_2(x), \dots, y_n(x)$  that have derivatives up to order  $n-1$ , the Wronskian  $W(x)$  is defined as the determinant:

$$W(x) = \begin{vmatrix} y_1(x) & y_2(x) & \dots & y_n(x) & y_1'(x) & y_2'(x) & \dots & y_n'(x) & \dots & \dots & \dots & y_1^{(n-1)}(x) & y_2^{(n-1)}(x) \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \end{vmatrix}$$

For two functions, the Wronskian simplifies to:

$$W_{(y_1, y_2)}(x) = y_1(x)y_2'(x) - y_2(x)y_1'(x)$$

For three functions, it becomes:

$$W_{(y_1, y_2, y_3)}(x) = \begin{vmatrix} y_1(x) & y_2(x) & y_3(x) & y_1'(x) & y_2'(x) & y_3'(x) & y_1''(x) & y_2''(x) & y_3''(x) \end{vmatrix}$$

### Theorem on the Wronskian

The key theorem regarding the Wronskian states:

If  $y_1(x)$ ,  $y_2(x)$ , ...,  $y_n(x)$  are solutions to a linear homogeneous differential equation on an interval  $I$ , then:

1. Either  $W(x) = 0$  for all  $x$  in  $I$ , or
2.  $W(x) \neq 0$  for all  $x$  in  $I$ .

Moreover, if  $W(x) \neq 0$  at even a single point in  $I$ , then the functions are linearly independent on  $I$ .

### Abel's Identity

For an  $n$ th-order linear homogeneous differential equation in the form:

$$Y^{(n)} + p_1(x)y^{(n-1)} + \dots + p_{n-1}(x)y' + p_n(x)y = 0$$

If  $W(x)$  is the Wronskian of  $n$  solutions, then:

$$W(x) = W(x_0) \cdot \exp\left[-\int p_1(x)dx\right]$$

where  $x_0$  is any point in the interval  $I$ .

This formula, known as Abel's Identity, allows us to compute the Wronskian without evaluating the determinant directly.

### Applications of the Wronskian

The Wronskian has several important applications:

1. **Testing for Linear Independence:** If  $W(x) \neq 0$  at any point, the functions are linearly independent.
2. **Constructing General Solutions:** For linear homogeneous differential equations.
3. **Method of Variation of Parameters:** For solving non-homogeneous equations.
4. **Reduction of Order:** For finding additional solutions when one solution is known.

### Computing the Wronskian: Examples

#### Example 1: Second-Order Case

For  $y_1(x) = e^x$  and  $y_2(x) = e^{2x}$ :



$$W(x) = \begin{vmatrix} e^x & e^{2x} & e^x & 2e^{2x} \end{vmatrix}$$

$$W(x) = e^x \cdot 2e^{2x} - e^{2x} \cdot e^x = 2e^{3x} - e^{3x} = e^{3x}$$

Since  $W(x) \neq 0$  for all  $x$ , the functions are linearly independent.

### Example 2: Third-Order Case

For  $y_1(x) = 1$ ,  $y_2(x) = x$ ,  $y_3(x) = x^2$ :

$$W(x) = \begin{vmatrix} 1 & x & x^2 & 0 & 1 & 2x & 0 & 0 & 2 \end{vmatrix}$$

$$W(x) = 1 \cdot 1 \cdot 2 = 2$$

Since  $W(x) = 2 \neq 0$  for all  $x$ , these functions are linearly independent.

### Special Cases and Properties

1. **Zero Wronskian:** If  $W(x) = 0$  for all  $x$ , the functions may or may not be linearly dependent (a zero Wronskian is a necessary but not sufficient condition for linear dependence).
2. **Wronskian of a Fundamental Set:** If the functions form a fundamental set of solutions for an  $n$ th-order homogeneous linear differential equation, their Wronskian is never zero.
3. **Wronskian and Initial Conditions:** For an initial value problem, the Wronskian evaluated at the initial point helps determine whether a unique solution exists.

## 2.7. Non-Homogeneous Equations of Order Two

### Structure of Non-Homogeneous Equations

A second-order linear non-homogeneous differential equation has the form:

$$a(x)y'' + b(x)y' + c(x)y = f(x)$$

where  $f(x) \neq 0$  is the non-homogeneous term (also called the forcing function or input).

### General Solution Structure

The general solution to a non-homogeneous equation consists of two parts:

$$y(x) = y_h(x) + y_p(x)$$

where:

- $y_h(x)$  is the general solution to the corresponding homogeneous equation (called the complementary function)
- $y_p(x)$  is any particular solution to the non-homogeneous equation

### **Methods for Finding Particular Solutions**

There are several methods for finding particular solutions:

#### **1. Method of Undetermined Coefficients**

This method works when  $f(x)$  and its derivatives form a finite set of linearly independent functions. We assume a solution form based on  $f(x)$  and determine the coefficients.

#### **When to Use**

This method is effective when  $f(x)$  is:

- A polynomial
- An exponential function ( $e^{ax}$ )
- A sine or cosine function
- A product of the above types

#### **Procedure**

1. Identify the form of  $f(x)$
2. Propose a trial solution  $y_p(x)$  with undetermined coefficients
3. Substitute into the differential equation
4. Solve for the coefficients by equating like terms

#### **Important Note**

If any term in the trial solution is already a solution to the homogeneous equation, multiply the entire trial solution by  $x$  (or  $x^2$  if necessary) to make it linearly independent from the homogeneous solutions.

#### **2. Method of Variation of Parameters**

This is a general method that works for any continuous  $f(x)$ .

### Procedure

For a second-order equation, if  $y_1(x)$  and  $y_2(x)$  are linearly independent solutions to the homogeneous equation, then:

$$y_p(x) = u_1(x)y_1(x) + u_2(x)y_2(x)$$

where  $u_1(x)$  and  $u_2(x)$  are determined by solving:

$$u_1'(x)y_1(x) + u_2'(x)y_2(x) = 0 \quad u_1'(x)y_1'(x) + u_2'(x)y_2'(x) = f(x)/a(x)$$

The solutions are:

$$u_1'(x) = -y_2(x)f(x)/[a(x)W(x)] \quad u_2'(x) = y_1(x)f(x)/[a(x)W(x)]$$

where  $W(x)$  is the Wronskian of  $y_1$  and  $y_2$ .

Integrating to find  $u_1(x)$  and  $u_2(x)$  gives the particular solution:

$$y_p(x) = -y_1(x) \int [y_2(x)f(x)/(a(x)W(x))]dx + y_2(x) \int [y_1(x)f(x)/(a(x)W(x))]dx$$

### 3. Operator Method

This involves using differential operators to factor and solve the equation.

#### Behavior of Solutions

The behavior of solutions to non-homogeneous equations depends on:

1. **Transient Response:** Governed by the homogeneous solution  $y_h(x)$ , which typically decays over time in stable systems.
2. **Steady-State Response:** Governed by the particular solution  $y_p(x)$ , which persists and matches the pattern of the input  $f(x)$ .

#### Resonance

A special situation occurs when  $f(x)$  contains terms that are solutions to the homogeneous equation. This leads to resonance, where the response can grow without bound.

For example, if  $y'' + y = \sin(x)$ , the solution contains terms with  $x \cdot \sin(x)$ , showing amplitude growth over time.

### Solved Problems

#### Problem 1: Testing Linear Independence Using Definition

**Problem:** Determine whether the functions  $y_1(x) = x$ ,  $y_2(x) = x|x|$ , and  $y_3(x) = x^3$  are linearly independent on the interval  $(-\infty, \infty)$ .

**Solution:**

Let's assume there exist constants  $c_1$ ,  $c_2$ , and  $c_3$ , not all zero, such that:

$$c_1x + c_2x|x| + c_3x^3 = 0 \text{ for all } x \in (-\infty, \infty)$$

For  $x > 0$ , we have  $|x| = x$ , so the equation becomes:  $c_1x + c_2x^2 + c_3x^3 = 0$

For this to be true for all  $x > 0$ , each coefficient must be zero:  $c_1 = c_2 = c_3 = 0$

But for  $x < 0$ , we have  $|x| = -x$ , so the equation becomes:  $c_1x - c_2x^2 + c_3x^3 = 0$

Again, for this to be true for all  $x < 0$ , each coefficient must be zero:  $c_1 = -c_2 = c_3 = 0$

Combining these constraints:

- From the first case:  $c_1 = c_2 = c_3 = 0$
- From the second case:  $c_1 = -c_2 = c_3 = 0$

This implies  $c_1 = 0$  and  $c_2 = 0$ , which is consistent. Therefore, the only solution is  $c_1 = c_2 = c_3 = 0$ , meaning the functions are linearly independent on  $(-\infty, \infty)$ .

## Problem 2: Computing and Interpreting the Wronskian

**Problem:** Compute the Wronskian of  $y_1(x) = e^x$ ,  $y_2(x) = e^{-x}$ , and determine if they form a fundamental set of solutions for the differential equation  $y'' - y = 0$ .

**Solution:**

First, let's compute the Wronskian:

$$W(y_1, y_2)(x) = \begin{vmatrix} e^x & e^{-x} \\ e^x & -e^x \end{vmatrix}$$

$$W(x) = e^x \cdot (-e^{-x}) - e^{-x} \cdot e^x = -e^0 - e^0 = -2$$

Since  $W(x) = -2 \neq 0$  for all  $x$ , the functions are linearly independent.

Now, let's check if they satisfy the differential equation  $y'' - y = 0$ :

$$\text{For } y_1(x) = e^x: y_1'(x) = e^x, y_1''(x) = e^x, y_1''(x) - y_1(x) = e^x - e^x = 0 \quad \checkmark$$

For  $y_2(x) = e^{-x}$ :  $y_2'(x) = -e^{-x}$   $y_2''(x) = e^{-x}$   $y_2''(x) - y_2(x) = e^{-x} - e^{-x} = 0$  ✓

Both functions satisfy the differential equation. Since they are also linearly independent, they form a fundamental set of solutions for  $y'' - y = 0$ .

The general solution is:  $y(x) = c_1 e^x + c_2 e^{-x}$

where  $c_1$  and  $c_2$  are arbitrary constants.

### Problem 3: Using Abel's Identity to Find the Wronskian

**Problem:** Use Abel's Identity to find the Wronskian of solutions to the differential equation:  $y'' - 2y' + y = 0$

**Solution:**

First, we rewrite the equation in standard form:  $y'' - 2y' + y = 0$

Comparing with the standard form  $y'' + p_1(x)y' + p_2(x)y = 0$ :  $p_1(x) = -2$   $p_2(x) = 1$

By Abel's Identity, if  $W(x)$  is the Wronskian of two linearly independent solutions, then:  $W(x) = W(x_0) \cdot \exp[-\int p_1(x) dx] = W(x_0) \cdot \exp[-\int (-2) dx] = W(x_0) \cdot \exp[2x]$

To find  $W(x_0)$ , we need the actual solutions. The characteristic equation for  $y'' - 2y' + y = 0$  is:  $r^2 - 2r + 1 = 0$   $(r - 1)^2 = 0$   $r = 1$  (repeated root)

So the solutions are:  $y_1(x) = e^x$   $y_2(x) = xe^x$

Let's compute  $W(x_0)$  at  $x_0 = 0$ :  $W(0) = \begin{vmatrix} e^0 & 0 \cdot e^0 \\ 0 \cdot e^0 & 0 \cdot e^0 + 1 \cdot e^0 \end{vmatrix} = \begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} = 1 \cdot 1 - 0 \cdot 1 = 1$

Therefore, by Abel's Identity:  $W(x) = 1 \cdot e^{2x} = e^{2x}$

We can verify this by direct computation:  $W(x) = \begin{vmatrix} e^x & xe^x \\ e^x & e^x + xe^x \end{vmatrix} = e^x(e^x + xe^x) - xe^x \cdot e^x = e^{2x} + xe^{2x} - xe^{2x} = e^{2x}$

which confirms our result from Abel's Identity.

#### Problem 4: Solving a Non-Homogeneous Equation Using Undetermined Coefficients

**Problem:** Solve the non-homogeneous differential equation:  $y'' + 4y = 3\sin(2x)$

**Solution:**

Step 1: Find the complementary solution (homogeneous solution). The characteristic equation for  $y'' + 4y = 0$  is:  $r^2 + 4 = 0$   $r = \pm 2i$

Therefore, the complementary solution is:  $y_h(x) = c_1\cos(2x) + c_2\sin(2x)$

Step 2: Find the particular solution using the method of undetermined coefficients. Since  $3\sin(2x)$  is already included in the complementary solution, we need to use a modified form:  $y_p(x) = Axcos(2x) + Bxsin(2x)$

Step 3: Find the derivatives of  $y_p(x)$ .  $y_p'(x) = A[\cos(2x) - 2x\sin(2x)] + B[\sin(2x) + 2xcos(2x)] = A\cos(2x) - 2Axs\sin(2x) + B\sin(2x) + 2Bxcos(2x)$

$y_p''(x) = -2A\sin(2x) - 2A[\sin(2x) + 2xcos(2x)] - 2B\cos(2x) + 2B[\cos(2x) - 2xsin(2x)] = -2A\sin(2x) - 2A\sin(2x) - 4Axcos(2x) - 2B\cos(2x) + 2B\cos(2x) - 4Bxsin(2x) = -4A\sin(2x) - 4Axcos(2x) - 4Bxsin(2x)$

Step 4: Substitute into the original equation.  $y'' + 4y = 3\sin(2x)$   $[-4A\sin(2x) - 4Axcos(2x) - 4Bxsin(2x)] + 4[Axcos(2x) + Bxsin(2x)] = 3\sin(2x) - 4A\sin(2x) - 4Axcos(2x) - 4Bxsin(2x) + 4Axcos(2x) + 4Bxsin(2x) = 3\sin(2x) - 4A\sin(2x) = 3\sin(2x)$

Step 5: Equate coefficients.  $-4A = 3$   $A = -3/4$   $B$  does not appear in the equation, so we can set  $B = 0$ .

Step 6: Write the particular solution.  $y_p(x) = -3/4 \cdot xcos(2x)$

Step 7: Combine the complementary and particular solutions.  $y(x) = y_h(x) + y_p(x) = c_1\cos(2x) + c_2\sin(2x) - 3/4 \cdot xcos(2x)$

### Problem 5: Solving a Non-Homogeneous Equation Using Variation of Parameters

**Problem:** Solve the non-homogeneous differential equation:  $y'' - y = \sec^2(x)$

**Solution:**

Step 1: Find the complementary solution. The characteristic equation for  $y'' - y = 0$  is:  $r^2 - 1 = 0$   $r = \pm 1$

The complementary solution is:  $y_h(x) = c_1 e^x + c_2 e^{-x}$

Step 2: Apply the method of variation of parameters. Let  $y_1(x) = e^x$  and  $y_2(x) = e^{-x}$

Calculate the Wronskian:  $W(x) = \begin{vmatrix} e^x & e^{-x} \\ e^x & -e^x \end{vmatrix} = -e^x \cdot e^{-x} - e^{-x} \cdot e^x = -2$

Step 3: Compute the integrals for variation of parameters.  $u_1'(x) = -y_2(x)f(x)/W(x) = -e^{-x} \cdot \sec^2(x)/(-2) = e^{-x} \cdot \sec^2(x)/2$   $u_2'(x) = y_1(x)f(x)/W(x) = e^x \cdot \sec^2(x)/(-2) = -e^x \cdot \sec^2(x)/2$

Step 4: Integrate to find  $u_1(x)$  and  $u_2(x)$ . Using the identity  $\sec^2(x) = 1 + \tan^2(x)$ :

$$u_1(x) = \int e^{-x} \cdot \sec^2(x)/2 \, dx = 1/2 \int e^{-x} \cdot (1 + \tan^2(x)) \, dx = 1/2 \left[ \int e^{-x} \, dx + \int e^{-x} \cdot \tan^2(x) \, dx \right]$$

The first integral is  $-e^{-x}/2$ . The second integral is more complex. Using integration by parts and the substitution  $\tan(x) = u$ , we get:

$$u_1(x) = -e^{-x}/2 - e^{-x} \cdot \tan(x)/2 + C_1$$

$$\text{Similarly: } u_2(x) = -e^x \cdot \tan(x)/2 + C_2$$

Step 5: Form the particular solution.  $y_p(x) = u_1(x)y_1(x) + u_2(x)y_2(x) = [-e^{-x}/2 - e^{-x} \cdot \tan(x)/2] \cdot e^x + [-e^x \cdot \tan(x)/2] \cdot e^{-x} = -1/2 - \tan(x)/2 - \tan(x)/2 = -1/2 - \tan(x)$

Step 6: Write the general solution.  $y(x) = y_h(x) + y_p(x) = c_1 e^x + c_2 e^{-x} - 1/2 - \tan(x)$

## Unsolved Problems

### Problem 1

Determine whether the functions  $y_1(x) = x^2$ ,  $y_2(x) = |x|$ ,  $y_3(x) = x^4$  are linearly independent on the interval  $(-\infty, \infty)$ .

### Problem 2

Calculate the Wronskian of the functions  $y_1(x) = \sin(2x)$ ,  $y_2(x) = \cos(2x)$ ,  $y_3(x) = e^x$  and determine if they form a fundamental set of solutions for any third-order linear homogeneous differential equation.

### Problem 3

Use Abel's Identity to find the Wronskian of solutions to the differential equation:  $x^2y'' + xy' - y = 0$

### Problem 4

Solve the non-homogeneous differential equation:  $y'' + 9y = x \cdot \cos(3x)$  using the method of undetermined coefficients.

### Problem 5

Solve the non-homogeneous differential equation:  $y'' - 4y' + 4y = e^{2x} \cdot \ln(x)$  using the method of variation of parameters, given that  $y_1(x) = e^{2x}$  and  $y_2(x) = xe^{2x}$  are solutions to the homogeneous equation.



## CHECK YOUR PROGRESS

- 1) Define a homogeneous second-order differential equation.

.....  
.....  
.....

- 2) State the superposition principle for homogeneous equations.

.....  
.....  
.....

- 3) How do initial conditions determine the constants of integration?

.....  
.....  
.....

## LET US SUM UP

A second-order homogeneous differential equation is one in which every term contains either the dependent variable  $y$  or its derivatives. It is typically written as  $y'' + P(x)y' + Q(x)y = 0$ . The superposition principle applies, meaning that any linear combination of two independent solutions is also a solution. When initial conditions such as  $y(x_0)$  and  $y'(x_0)$  are given, the arbitrary constants can be uniquely determined.

## UNIT END EXERCISES

### Short Answer Questions:

1. How do you solve an initial value problem for a second-order linear equation?
2. Explain how complex roots affect the general solution of a second-order equation.
3. What is the particular solution of a non-homogeneous equation?

### Long Answer Questions:

1. Solve the equation  $y''+3y'+2y=0$  using the characteristic equation method.
2. Explain and prove the method of variation of parameters for solving non-homogeneous equations.
3. Solve the equation  $y''-y'-6y=0$  using the characteristic equation.

### REFERENCES AND SUGGESTED READINGS:

1. Boyce, W. E., & DiPrima, R. C. (2021). Elementary Differential Equations and Boundary Value Problems (11th ed.). Wiley.
2. Tenenbaum, M., & Pollard, H. (2012). Ordinary Differential Equations: An Elementary Textbook for Students of Mathematics, Engineering, and the Sciences. Dover Publications.
3. Ross, S. L. (2013). Differential Equations (4th ed.). Wiley.

## UNIT 3

### LINEAR DEPENDENCE AND INDEPENDENCE - A FORMULA FOR THE WRONSKIAN – THE NON-HOMOGENEOUS EQUATION OF ORDER TWO.

#### 3.1 Key Concepts

##### 1. Linear Dependence and Independence:

- Functions are linearly dependent if one can be expressed as a linear combination of others.
- The general solution to an  $n$ th-order homogeneous linear differential equation requires  $n$  linearly independent solutions.

##### 2. The Wronskian:

- A determinant that helps determine linear independence of functions.
- If the Wronskian is non-zero at any point, the functions are linearly independent.
- Abel's Identity provides a formula for the Wronskian without direct computation.

##### 3. Non-Homogeneous Equations:

- The general solution consists of the complementary function (homogeneous solution) plus a particular solution.
- Methods for finding particular solutions include undetermined coefficients and variation of parameters.
- Resonance occurs when the forcing function matches the natural frequency of the system.

These concepts are fundamental to understanding and solving differential equations, with applications in physics, engineering, economics, and many other fields.

#### 3.2 Applications of Second-Order Linear Equations

Second-order linear differential equations play a crucial role in modeling physical systems across numerous fields including physics, engineering, and applied mathematics. These equations help describe phenomena ranging from simple harmonic motion to more complex scenarios like damped oscillations and forced vibrations.

### **The General Form and Physical Significance**

A second-order linear differential equation typically takes the form:

$$a(x) \cdot y''(x) + b(x) \cdot y'(x) + c(x) \cdot y(x) = f(x)$$

Where:

- $y''(x)$  represents the second derivative of  $y$  with respect to  $x$
- $y'(x)$  represents the first derivative
- $a(x)$ ,  $b(x)$ , and  $c(x)$  are coefficients that may be constants or functions of  $x$
- $f(x)$  is the non-homogeneous term (when  $f(x) = 0$ , we have a homogeneous equation)

In physical systems, the terms often represent:

- The second derivative ( $y''$ ) typically corresponds to acceleration
- The first derivative ( $y'$ ) typically corresponds to velocity or a damping term
- The function itself ( $y$ ) typically corresponds to position or displacement
- The coefficients represent physical parameters like mass, damping coefficient, or spring constant

### **Common Physical Applications**

#### **1. Spring-Mass Systems**

One of the most fundamental applications is modeling a spring-mass system. The equation takes the form:

$$m \cdot y''(t) + c \cdot y'(t) + k \cdot y(t) = F(t)$$

Where:

- $m$  represents mass
- $c$  represents the damping coefficient
- $k$  represents the spring constant
- $F(t)$  represents an external force
- $y(t)$  represents displacement from equilibrium

Depending on the values of these parameters, we observe different behaviors:

- When  $c = 0$  and  $F(t) = 0$ : Simple harmonic motion
- When  $0 < c < 2\sqrt{(km)}$  and  $F(t) = 0$ : Underdamped oscillation
- When  $c = 2\sqrt{(km)}$  and  $F(t) = 0$ : Critically damped motion
- When  $c > 2\sqrt{(km)}$  and  $F(t) = 0$ : Overdamped motion
- When  $F(t) \neq 0$ : Forced oscillation

## 2. RLC Circuits

Electrical circuits with resistors, inductors, and capacitors are modeled using second-order equations:

$$L \cdot d^2q/dt^2 + R \cdot dq/dt + (1/C) \cdot q = E(t)$$

Where:

- $L$  is inductance
- $R$  is resistance
- $C$  is capacitance
- $q$  is electric charge
- $E(t)$  is the applied voltage

This is mathematically identical to the spring-mass system, highlighting the parallel between mechanical and electrical systems.

## 3. Beam Deflection

The equation for the deflection  $y(x)$  of a uniform beam is:

$$EI \cdot d^4y/dx^4 = w(x)$$

Where:

- E is Young's modulus
- I is the area moment of inertia
- $w(x)$  is the distributed load

This is a fourth-order equation but can be reduced to a system of second-order equations.

#### 4. Heat Transfer and Diffusion

The one-dimensional heat equation:

$$\partial^2 u / \partial x^2 = (1/\alpha) \cdot \partial u / \partial t$$

Where  $u(x,t)$  is temperature, can be solved using techniques for second-order equations.

#### Solving Second-Order Linear Equations

The general approach to solving second-order linear equations involves:

1. For homogeneous equations ( $f(x) = 0$ ):
  - Find the general solution  $y_h$  using characteristic equations or other methods
2. For non-homogeneous equations ( $f(x) \neq 0$ ):
  - Find a particular solution  $y_p$  using methods like undetermined coefficients or variation of parameters
  - The complete solution is  $y = y_h + y_p$

#### Solved Problems

##### Solved Problem 1: Simple Harmonic Motion

**Problem:** A mass of 2 kg is attached to a spring with spring constant  $k = 8$  N/m. If the mass is displaced 0.5 meters from equilibrium and released from rest, find the position of the mass as a function of time.

##### Solution:

The differential equation for this system is:  $m \cdot y''(t) + k \cdot y(t) = 0$

Substituting the given values:  $2 \cdot y''(t) + 8 \cdot y(t) = 0$   $y''(t) + 4 \cdot y(t) = 0$

This is a homogeneous second-order equation with constant coefficients.

The characteristic equation is:  $r^2 + 4 = 0$   $r = \pm 2i$

The general solution is:  $y(t) = C_1 \cdot \cos(2t) + C_2 \cdot \sin(2t)$

Given initial conditions:  $y(0) = 0.5$  (initial displacement)  $y'(0) = 0$  (released from rest)

Applying the first condition:  $y(0) = C_1 \cdot \cos(0) + C_2 \cdot \sin(0) = 0.5$   $C_1 = 0.5$

Applying the second condition:  $y'(t) = -2C_1 \cdot \sin(2t) + 2C_2 \cdot \cos(2t)$   $y'(0) = -2C_1 \cdot \sin(0) + 2C_2 \cdot \cos(0) = 0$   $2C_2 = 0$   $C_2 = 0$

Therefore, the position as a function of time is:  $y(t) = 0.5 \cdot \cos(2t)$

This represents simple harmonic motion with amplitude 0.5 meters and angular frequency 2 rad/s. The period of oscillation is  $\pi$  seconds.

### Solved Problem 2: Damped Oscillations

**Problem:** A mass-spring-damper system is governed by the equation  $y''(t) + 4y'(t) + 4y(t) = 0$ . If  $y(0) = 2$  and  $y'(0) = -4$ , find the position function  $y(t)$ .

**Solution:**

The differential equation is:  $y''(t) + 4y'(t) + 4y(t) = 0$

This is a homogeneous second-order equation with constant coefficients.

The characteristic equation is:  $r^2 + 4r + 4 = 0$   $(r + 2)^2 = 0$   $r = -2$  (repeated root)

For a repeated root, the general solution is:  $y(t) = (C_1 + C_2t) \cdot e^{(-2t)}$

Given initial conditions:  $y(0) = 2$   $y'(0) = -4$

Applying the first condition:  $y(0) = C_1 = 2$

To find  $C_2$ , we compute the derivative:  $y'(t) = -2(C_1 + C_2t)e^{(-2t)} + C_2e^{(-2t)} = (-2C_1 + C_2 - 2C_2t)e^{(-2t)}$

Applying the second condition:  $y'(0) = -2C_1 + C_2 = -4$   $-2(2) + C_2 = -4$   $-4 + C_2 = -4$   $C_2 = 0$

Therefore, the position function is:  $y(t) = 2e^{(-2t)}$

This represents a critically damped system where the mass approaches equilibrium without oscillating. The system returns to equilibrium asymptotically as  $t$  increases.

### Solved Problem 3: Forced Vibrations

**Problem:** A spring-mass system is described by the equation  $y''(t) + 9y(t) = 3\cos(3t)$ . If  $y(0) = 0$  and  $y'(0) = 2$ , find the solution  $y(t)$ .

#### Solution:

The differential equation is:  $y''(t) + 9y(t) = 3\cos(3t)$

This is a non-homogeneous equation. We first find the complementary solution (solution to the homogeneous equation):  $y''(t) + 9y(t) = 0$

The characteristic equation is:  $r^2 + 9 = 0$   $r = \pm 3i$

So the complementary solution is:  $y_h(t) = C_1\cos(3t) + C_2\sin(3t)$

Next, we find a particular solution. Since the right side involves  $\cos(3t)$  and this term also appears in the complementary solution, we use:  $y_p(t) = t(A\cos(3t) + B\sin(3t))$

Taking derivatives:  $y_p'(t) = A\cos(3t) + B\sin(3t) + t(-3A\sin(3t) + 3B\cos(3t))$   
 $y_p''(t) = -3A\sin(3t) + 3B\cos(3t) + t(-3A\cdot 3\cos(3t) - 3B\cdot 3\sin(3t))$   
 $+ (-3A\sin(3t) + 3B\cos(3t)) = -6A\sin(3t) + 6B\cos(3t) - 9At\cos(3t) - 9Bt\sin(3t)$

Substituting into the original equation:  $y_p''(t) + 9y_p(t) = 3\cos(3t)$   
 $[-6A\sin(3t) + 6B\cos(3t) - 9At\cos(3t) - 9Bt\sin(3t)] + 9[t(A\cos(3t) + B\sin(3t))] = 3\cos(3t)$   
 $-6A\sin(3t) + 6B\cos(3t) - 9At\cos(3t) - 9Bt\sin(3t) + 9At\cos(3t) + 9Bt\sin(3t) = 3\cos(3t)$   
 $-6A\sin(3t) + 6B\cos(3t) = 3\cos(3t)$

Comparing coefficients:  $-6A = 0$ , so  $A = 0$   $6B = 3$ , so  $B = 1/2$

Therefore,  $y_p(t) = (t/2)\sin(3t)$

The complete solution is:  $y(t) = y_h(t) + y_p(t)$   $y(t) = C_1\cos(3t) + C_2\sin(3t) + (t/2)\sin(3t)$

Applying the initial condition  $y(0) = 0$ :  $y(0) = C_1\cos(0) + C_2\sin(0) + (0/2)\sin(0) = 0$   $C_1 = 0$



For the second condition,  $y'(0) = 2$ , we need to compute  $y'(t)$ :  $y'(t) = -3C_1\sin(3t) + 3C_2\cos(3t) + (1/2)\cdot\sin(3t) + (t/2)\cdot 3\cos(3t) = -3C_1\sin(3t) + 3C_2\cos(3t) + (1/2)\cdot\sin(3t) + (3t/2)\cdot\cos(3t)$

At  $t = 0$ :  $y'(0) = -3C_1\sin(0) + 3C_2\cos(0) + (1/2)\cdot\sin(0) + (3\cdot 0/2)\cdot\cos(0) = 3C_2 = 2 \Rightarrow C_2 = 2/3$

Therefore, the complete solution is:  $y(t) = (2/3)\cdot\sin(3t) + (t/2)\cdot\sin(3t)$   $y(t) = \sin(3t)\cdot(2/3 + t/2)$

This solution represents forced vibrations, where the system exhibits resonance because the forcing frequency matches the natural frequency of the system.

#### Solved Problem 4: RLC Circuit

**Problem:** An RLC circuit has an inductance  $L = 1$  H, resistance  $R = 6 \Omega$ , and capacitance  $C = 1/16$  F. If the initial current is zero and the initial charge on the capacitor is 2 coulombs, find the charge  $q(t)$  on the capacitor as a function of time.

#### Solution:

The differential equation for the charge  $q(t)$  in an RLC circuit is:  $L \cdot d^2q/dt^2 + R \cdot dq/dt + (1/C) \cdot q = 0$

Substituting the given values:  $1 \cdot d^2q/dt^2 + 6 \cdot dq/dt + 16 \cdot q = 0$   $d^2q/dt^2 + 6 \cdot dq/dt + 16 \cdot q = 0$

This is a homogeneous second-order equation with constant coefficients.

The characteristic equation is:  $r^2 + 6r + 16 = 0$

Using the quadratic formula:  $r = (-6 \pm \sqrt{(36 - 64)})/2 = (-6 \pm \sqrt{(-28)})/2 = (-6 \pm 2\sqrt{7}i)/2 = -3 \pm \sqrt{7}i$

The general solution is:  $q(t) = e^{(-3t)} \cdot [C_1 \cdot \cos(\sqrt{7}t) + C_2 \cdot \sin(\sqrt{7}t)]$

Given initial conditions:  $q(0) = 2$  (initial charge)  $dq/dt(0) = 0$  (initial current is zero)

Applying the first condition:  $q(0) = e^{(0)} \cdot [C_1 \cdot \cos(0) + C_2 \cdot \sin(0)] = C_1 = 2$

To find  $C_2$ , we compute the derivative:  $dq/dt = -3e^{(-3t)} \cdot [C_1 \cdot \cos(\sqrt{7}t) + C_2 \cdot \sin(\sqrt{7}t)] + e^{(-3t)} \cdot [-C_1 \cdot \sqrt{7} \cdot \sin(\sqrt{7}t) + C_2 \cdot \sqrt{7} \cdot \cos(\sqrt{7}t)] = e^{(-3t)} \cdot [-3C_1 \cdot \cos(\sqrt{7}t) - 3C_2 \cdot \sin(\sqrt{7}t) - C_1 \cdot \sqrt{7} \cdot \sin(\sqrt{7}t) + C_2 \cdot \sqrt{7} \cdot \cos(\sqrt{7}t)]$

Applying the second condition:  $\frac{dq}{dt}(0) = e(0) \cdot [-3C_1 \cdot \cos(0) - 3C_2 \cdot \sin(0) - C_1 \cdot \sqrt{7} \cdot \sin(0) + C_2 \cdot \sqrt{7} \cdot \cos(0)] = -3C_1 + C_2 \cdot \sqrt{7} = 0 = -3(2) + C_2 \cdot \sqrt{7} = 0 = -6 + C_2 \cdot \sqrt{7} = 0 = C_2 = 6/\sqrt{7} = 6\sqrt{7}/7$

Therefore, the charge as a function of time is:  $q(t) = e^{(-3t)} \cdot [2 \cdot \cos(\sqrt{7}t) + (6\sqrt{7}/7) \cdot \sin(\sqrt{7}t)]$

This represents an underdamped RLC circuit where the charge oscillates with decreasing amplitude due to the resistance.

### Solved Problem 5: Beam Deflection

**Problem:** A uniform beam of length  $L$  is simply supported at both ends and carries a uniform load  $w$  per unit length. Find the equation for the deflection curve.

#### Solution:

The differential equation for the deflection  $y(x)$  of a uniform beam under a distributed load  $w$  is:  $EI \cdot \frac{d^4y}{dx^4} = w$

Where  $E$  is Young's modulus,  $I$  is the moment of inertia, and  $w$  is the load per unit length.

For a constant load  $w$ , we can integrate this equation directly:  $EI \cdot \frac{d^3y}{dx^3} = wx + C_1$   
 $EI \cdot \frac{d^2y}{dx^2} = (w/2)x^2 + C_1x + C_2$   
 $EI \cdot \frac{dy}{dx} = (w/6)x^3 + (C_1/2)x^2 + C_2x + C_3$   
 $EI \cdot y = (w/24)x^4 + (C_1/6)x^3 + (C_2/2)x^2 + C_3x + C_4$

For a simply supported beam, the boundary conditions are:  $y(0) = 0$  (deflection at left end is zero)  $y(L) = 0$  (deflection at right end is zero)  
 $\frac{d^2y}{dx^2}(0) = 0$  (bending moment at left end is zero)  $\frac{d^2y}{dx^2}(L) = 0$  (bending moment at right end is zero)

Applying  $y(0) = 0$ :  $EI \cdot y(0) = C_4 = 0$

Applying  $\frac{d^2y}{dx^2}(0) = 0$ :  $EI \cdot \frac{d^2y}{dx^2}(0) = C_2 = 0$

From the remaining two conditions:  $y(L) = (w/24)L^4 + (C_1/6)L^3 + C_3L = 0$   
 $\frac{d^2y}{dx^2}(L) = wL^2 + C_1L = 0$

From the last equation:  $C_1 = -wL$

Substituting into  $y(L) = 0$ :  $(w/24)L^4 - (wL/6)L^3 + C_3L = 0$   $(w/24)L^4 - (wL^4/6) + C_3L = 0$   $(wL^4/24) - (wL^4/6) + C_3L = 0$   $(wL^4/24) - (4wL^4/24) + C_3L = 0$   $(-3wL^4/24) + C_3L = 0$   $C_3 = (3wL^3/24) = (wL^3/8)$

Therefore, the deflection equation is:  $EI \cdot y = (w/24)x^4 - (wL/6)x^3 + (wL^3/8)x$

Simplifying:  $y = (w/24EI)[x^4 - 4Lx^3 + 3L^3x]$

This equation describes the deflection of the beam at any point  $x$  along its length under the uniform load  $w$ .

### **Unsolved Problems**

#### **Unsolved Problem 1: Damped Spring-Mass System**

A mass of 0.5 kg is attached to a spring with spring constant  $k = 12$  N/m and a damper with damping coefficient  $c = 3$  N·s/m. The mass is pulled down 10 cm from equilibrium and released with an initial velocity of 0.2 m/s upward. Find the position function  $y(t)$  and determine whether the system is underdamped, critically damped, or overdamped.

#### **Unsolved Problem 2: Forced Vibrations with Damping**

Consider a spring-mass-damper system described by the equation:  $y''(t) + 4y'(t) + 13y(t) = 10\sin(2t)$

If  $y(0) = 0$  and  $y'(0) = 0$ , find the complete solution and determine the steady-state response.

#### **Unsolved Problem 3: RLC Circuit with Applied Voltage**

An RLC circuit with inductance  $L = 2$  H, resistance  $R = 8$   $\Omega$ , and capacitance  $C = 0.02$  F is connected to a voltage source  $E(t) = 12\cos(5t)$  V. If the initial charge on the capacitor is zero and the initial current is zero, find the charge  $q(t)$  on the capacitor as a function of time.

#### **Unsolved Problem 4: Heat Transfer in a Rod**

A rod of length  $L$  has its ends maintained at temperature 0. The initial temperature distribution in the rod is given by  $f(x) = \sin(\pi x/L)$ . Find the temperature  $u(x,t)$  at any point  $x$  and time  $t$ , given that the heat equation is:  $\partial^2 u / \partial x^2 = (1/\alpha) \cdot \partial u / \partial t$

With boundary conditions  $u(0,t) = u(L,t) = 0$  and initial condition  $u(x,0) = f(x)$ .

#### **Unsolved Problem 5: Cantilever Beam**

A cantilever beam of length  $L$  is fixed at one end ( $x = 0$ ) and free at the other end ( $x = L$ ). The beam carries a point load  $P$  at the free end. Find the equation for the deflection curve  $y(x)$ .

## **Applications in Various Fields**

### **Mechanical Engineering**

Second-order linear equations are essential in analyzing:

- Vibration analysis of structures
- Stress and strain in materials
- Control systems for mechanical devices
- Automotive suspension systems
- Structural dynamics of buildings

### **Electrical Engineering**

Key applications include:

- Circuit analysis (RLC circuits)
- Signal processing and filter design
- Control systems for electrical devices
- Power systems stability
- Electromagnetic wave propagation

### **Civil Engineering**

Applications encompass:

- Structural analysis of buildings and bridges
- Beam and column deflection
- Dynamic response of structures to earthquakes
- Fluid flow in pipes and channels
- Soil mechanics and foundation design

### **Aerospace Engineering**

Critical uses include:

- Aircraft and spacecraft dynamics
- Aeroelasticity (flutter analysis)
- Launch vehicle trajectory optimization
- Control system design
- Structural vibration of airframes

## **Advanced Topics**

### **Variable Coefficient Equations**

Many real-world problems lead to second-order equations with variable coefficients:  $a(x) \cdot y''(x) + b(x) \cdot y'(x) + c(x) \cdot y(x) = f(x)$

These are often more challenging to solve and may require numerical methods or series solutions like:

- Frobenius method
- Variation of parameters
- WKB approximation
- Numerical techniques (Runge-Kutta, finite differences)

### **Systems of Second-Order Equations**

Complex mechanical systems with multiple degrees of freedom lead to systems of coupled second-order equations that can be written in matrix form:  $[M]\{\ddot{x}\} + [C]\{\dot{x}\} + [K]\{x\} = \{F(t)\}$

Where:

- $[M]$  is the mass matrix
- $[C]$  is the damping matrix
- $[K]$  is the stiffness matrix
- $\{x\}$  is the displacement vector
- $\{F(t)\}$  is the forcing vector

These systems are typically solved using:

- Modal analysis
- Numerical integration
- State-space methods

### **Nonlinear Second-Order Equations**

Many physical systems exhibit nonlinear behavior, leading to nonlinear second-order equations such as:

- Duffing equation (nonlinear spring):  $\ddot{x} + \delta\dot{x} + \alpha x + \beta x^3 = F_0 \cos(\omega t)$
- Van der Pol equation (nonlinear damping):  $\ddot{x} - \mu(1-x^2)\dot{x} + x = 0$
- Pendulum equation (large displacements):  $\ddot{\theta} + (g/L)\sin(\theta) = 0$

These equations often exhibit complex behaviors like:

- Multiple equilibria
- Limit cycles
- Chaos
- Bifurcations

### **Computational Methods**

Modern approaches to solving second-order differential equations often involve computational methods:

#### **Finite Difference Methods**

Approximate derivatives using differences between discrete points:

- Forward difference:  $f'(x) \approx [f(x+h) - f(x)]/h$
- Central difference:  $f'(x) \approx [f(x+h) - f(x-h)]/(2h)$
- Second derivative:  $f''(x) \approx [f(x+h) - 2f(x) + f(x-h)]/h^2$

#### **Runge-Kutta Methods**

Higher-order methods that propagate a solution by combining information from several steps:

- RK4 (fourth-order Runge-Kutta) is widely used for its balance of accuracy and efficiency

## **Finite Element Methods**

Particularly useful for complex geometries and boundary conditions:

- Divide the domain into small elements
- Approximate the solution within each element
- Assemble a global system of equations
- Solve the resulting system

Second-order linear differential equations provide a powerful framework for modeling and analyzing a wide range of physical phenomena. From simple harmonic oscillators to complex structural dynamics, these equations form the mathematical foundation for understanding how systems respond to various inputs and disturbances. The applications span across multiple engineering disciplines, including mechanical, electrical, civil, and aerospace engineering. Understanding these equations and their solutions is essential for engineers and scientists working on problems involving motion, vibration, wave propagation, and structural analysis. As computational capabilities continue to advance, more complex systems can be modeled and analyzed using these fundamental equations, leading to improved designs and better understanding of physical phenomena.

## **Second-Order Differential Equations: Practical Applications in Contemporary Engineering and Science**

In the contemporary technologically advanced world, second-order differential equations constitute the mathematical basis for various engineering and scientific fields. These equations represent systems where the rate of change of a rate of change is essential, encompassing the oscillations of mechanical systems and the flow of electric current in circuits. The importance of knowing these equations is paramount, since they offer the analytical framework for comprehending and forecasting intricate dynamic behaviors in real-world situations.

Second-order differential equations are expressed as  $a(x)y'' + b(x)y' + c(x)y = f(x)$ , with the homogeneous case arising when  $f(x) = 0$ . Engineers, physicists, and applied mathematicians routinely confront these equations whether examining structure vibrations, devising control systems, modeling population dynamics, or creating electronic filters. The capacity to resolve

these equations effectively converts abstract mathematical principles into practical instruments for creativity and problem-solving.

### **Homogeneous Linear Differential Equations: Applications in Contemporary Structural Analysis**

Contemporary structural engineers predominantly utilize homogeneous second-order differential equations to assess building responses to environmental pressures. Examine a contemporary skyscraper exposed to wind forces or seismic events. The displacement  $y$  of the building as a function of time  $t$  typically adheres to the equation  $my'' + cy' + ky = 0$ , where  $m$  denotes the mass of the building,  $c$  signifies the damping coefficient from structural components, and  $k$  represents the stiffness of construction materials.

In the design of the Burj Khalifa or comparable supertall edifices, engineers must resolve these equations to forecast maximum displacements and guarantee safety margins. The characteristic equation  $mr^2 + cr + k = 0$  produces roots that indicate whether the structure will undergo critical damping (equal roots), underdamping (complex conjugate roots), or overdamping (distinct real roots). Each scenario necessitates distinct structural considerations—underdamped systems may require supplementary dampers to avert resonance, whereas overdamped systems may compromise responsiveness for stability. Contemporary computer techniques have transformed the practical use of these equations. Engineers utilize finite element analysis software that integrates these differential equations into millions of concurrent calculations, facilitating the optimization of structural parameters through numerous design iterations prior to actual construction.

### **Methods for Solving Homogeneous Equations**

Homogeneous second-order linear differential equations with constant coefficients ( $ay'' + by' + cy = 0$ ) are resolved by determining the roots of the characteristic equation  $ar^2 + br + c = 0$ . The structure of the general solution is contingent upon these roots:

One. For unique real roots  $r_1$  and  $r_2$ :  $y(x) = C_1e^{(r_1x)} + C_2e^{(r_2x)}$

Two. For repeated roots  $r_1 = r_2$ :  $y(x) = C_1e^{(r_1x)} + C_2xe^{(r_1x)}$

Three. For complex conjugate roots  $r_{1,2} = \alpha \pm \beta i$ :  $y(x) = e^{(\alpha x)}[C_1\cos(\beta x) + C_2\sin(\beta x)]$



These solutions represent physical phenomena such as damped oscillations in suspension systems, where the type of damping—over, under, or critical—correlates directly with the nature of the roots.

### **Initial Value Problems: Control Systems in Real-Time and Robotics**

Contemporary automated manufacturing facilities and autonomous cars utilize control systems that depend on resolving initial value problems (IVPs) linked to second-order differential equations. In designing a robotic arm for precise movement between positions, engineers must consider the initial location ( $y(0) = y_0$ ) and initial velocity ( $y'(0) = v_0$ ). A robotic surgical system, for example, may express arm movement as  $my'' + cy' + ky = F(t)$ , with  $F(t)$  being the input force. The surgical robot must operate with exceptional precision, frequently within microns, while ensuring smooth motion trajectories. Control engineers develop precise motion profiles that guarantee patient safety by resolving the corresponding initial value problem with defined initial conditions. The solutions are expressed as  $y(t) = C_1y_1(t) + C_2y_2(t)$ , where  $y_1$  and  $y_2$  are fundamental solutions to the homogeneous equation, and the constants  $C_1$  and  $C_2$  are ascertained from initial conditions. In fact, these constants directly correspond to control parameters in the system's software, determining the exact voltage or current applied to motors at certain millisecond intervals.

Contemporary machine learning methodologies have started to augment conventional IVP solutions, utilizing neural networks trained to forecast ideal constants derived from system identification data. This hybrid methodology facilitates adaptive regulation in dynamic contexts while preserving the mathematical precision of differential equation solutions.

### **Linear Independence and Dependence: Theoretical Basis and Practical Importance**

For a second-order differential equation, two solutions  $y_1(x)$  and  $y_2(x)$  are linearly independent on an interval  $I$  if the sole solution to  $c_1y_1(x) + c_2y_2(x) = 0$  for any  $x$  in  $I$  is  $c_1 = c_2 = 0$ . This abstract notion has significant practical ramifications across various domains. In contemporary vibration analysis, linear independence guarantees that engineers have identified all potential modes of vibration within a structure. Each linearly independent solution signifies a fundamental mode of

oscillation for the system. Omission of a mode may result in unforeseen resonance and structural failure.

The principle applies to signal processing, where linearly independent basis functions enable comprehensive representation of intricate signals. Contemporary compression methods such as JPEG and MP3 utilize transformations derived from linearly independent functions, facilitating efficient digital communication and storage. The Wronskian determinant serves as a practical test for linear independence, offering engineers a computational method to confirm the completeness of their solution sets.

### **Financial Modeling and Risk Evaluation via Differential Equations**

Financial analysts at contemporary investment firms employ the principles of linear independence and dependency when developing differential equation models for asset pricing and risk management. The value of a portfolio,  $V$ , may adhere to a second-order equation  $V'' + a(t)V' + b(t)V = f(t)$ , with  $f(t)$  denoting external market influences. Two solutions  $V_1$  and  $V_2$  are linearly independent if there are no constants  $c_1$  and  $c_2$  (not both zero) such that  $c_1V_1 + c_2V_2 = 0$  for all  $t$ . This independence signifies that the portfolio comprises genuinely diverse assets that react differently to market fluctuations—a vital factor in the current unstable financial environment. Quantitative analysts at companies such as Renaissance Technologies or Two Sigma utilize these mathematical principles in the creation of trading algorithms. By finding linearly independent variables influencing asset prices, they create more robust portfolios. This application encompasses advanced derivative pricing models, utilizing second-order differential equations to assess option prices under stochastic volatility conditions, surpassing mere stock diversification. The notion has acquired renewed importance due to the emergence of high-frequency trading, wherein algorithms must swiftly resolve these equations to detect arbitrage possibilities within microsecond intervals. The mathematical assurances of linear independence directly inform risk management techniques that have been essential during recent market volatility occurrences.

### **The Wronskian in Engineering Applications: Aerospace and Mechanical Systems**

Aerospace engineers developing contemporary commercial aircraft such as

the Boeing 787 or Airbus A350 frequently utilize the Wronskian determinant in their analysis of flight dynamics. The Wronskian  $W(y_1, y_2)(t) = y_1(t)y_2'(t) - y_1'(t)y_2(t)$  serves as an effective instrument to verify the linear independence of two candidate solutions  $y_1$  and  $y_2$  to a homogeneous second-order differential equation. In flutter analysis—a vital safety issue in aircraft design—engineers investigate aeroelastic processes via coupled differential equations. The Wronskian assists in determining when suggested solution sets are insufficient by exposing dependencies that could result in detrimental resonance circumstances. If  $W(y_1, y_2)(t) = 0$  for a certain  $t$ , the solutions are dependant, indicating possible structural weaknesses. Flight test engineers gather vibration data during aircraft certification and analyze the observed frequency responses in relation to projected outcomes. The Wronskian computation functions as a mathematical verification of the completeness of their analytical models. Contemporary airplane certification necessitates the demonstration that all critical vibration modes have been considered—a stipulation intrinsically connected to guaranteeing linearly independent solutions to the governing differential equations. The analytical expression for the Wronskian of a second-order linear homogeneous differential equation  $y'' + p(t)y' + q(t)y = 0$  is  $W(t) = W(0)e(-\int p(t)dt)$ . Engineers utilize this relationship to predict system behavior in untested operating conditions, hence ensuring safety margins within the aircraft's fly envelope.

### **Derivation and Application of the Wronskian Formula**

For a second-order linear homogeneous differential equation of the type  $y'' + p(x)y' + q(x)y = 0$ , the Wronskian  $W(x) = W(y_1, y_2)$  satisfies the differential equation:

$$W'(x) = -p(x)W(x)$$

This first-order equation possesses the solution:

$$W(x) = W(x_0) e^{-\int_{x_0}^x p(t)dt}$$

This formula offers numerous pragmatic insights:

1. The Wronskian is either identically zero or consistently non-zero over the specified interval.
2. If  $p(x) = 0$  (as in  $y'' + q(x)y = 0$ ), the Wronskian remains constant.

3. In standard form equations (where the coefficient of  $y''$  is 1), the behavior of the Wronskian is solely determined by the coefficient of  $y'$ .

Engineers employ this method to validate solution sets without the explicit computation of determinants at various places, hence enhancing efficiency in complex system analysis.

#### Non-Homogeneous Differential Equations: Communication and Signal Processing

The current telecommunications infrastructure heavily depends on the resolution of non-homogeneous differential equations. In the analysis of signal transmission via fiber optic networks or wireless channels, engineers utilize equations of the type  $y'' + a(t)y' + b(t)y = s(t)$ , with  $s(t)$  being the input signal.

Designers of 5G networks utilize these mathematical instruments to optimize antenna arrays and signal processing techniques. The comprehensive solution entails determining both the complementary function (solution to the homogeneous equation) and the particular integral (addressing the individual input). This mathematical paradigm immediately applies to actual filter design, modulation techniques, and error correction codes in contemporary communication systems.

Digital signal processing experts execute these solutions utilizing diverse strategies, such as change of parameters and the method of indeterminate coefficients. For example, when  $s(t)$  represents a sinusoidal carrier wave in radio communications, engineers want to find a specific solution of analogous form while circumventing resonance conditions where frequencies align with the system's intrinsic frequency—a phenomena that results in signal distortion. The variation of parameters method is particularly advantageous in contemporary adaptive filtering applications, where the system must adjust to fluctuating signal environments. Engineers design algorithms that maximize signal detection in noisy settings by creating solutions of the type  $y(t) = u_1(t)y_1(t) + u_2(t)y_2(t)$ , where  $u_1$  and  $u_2$  are functions defined by the non-homogeneity rather than constants.

#### Techniques for Resolving Non-Homogeneous Equations

A variety of techniques are available for determining specific solutions to non-homogeneous equations:

### Technique of Indeterminate Coefficients

When the non-homogeneous term  $f(x)$  is a polynomial, exponential, sine, cosine, or a product of these functions, engineers postulate a particular solution of analogous form with unspecified coefficients. This technique is extensively employed in electrical filter design, where input signals assume conventional formats.

For instance, if  $f(x) = 3x^2 + 2\sin(x)$ , we could propose:  $y_p(x) = Ax^2 + Bx + C + D\sin(x) + E\cos(x)$

Substituting this into the original equation and equating coefficients identifies the constants.

### Technique of Parameter Variation

In cases of intricate forcing functions or where the method of indeterminate coefficients proves cumbersome, the variation of parameters method offers a systematic solution. Having two linearly independent solutions  $y_1$  and  $y_2$  to the homogeneous equation, we proceed to construct:

$$y_p(x) = u_1(x)y_1(x) + u_2(x)y_2(x)$$

where:  $u_1'(x)y_1(x) + u_2'(x)y_2(x) = 0$  and  $u_1'(x)y_1'(x) + u_2'(x)y_2'(x) = f(x)$

This technique is very beneficial in contemporary control systems that need to react to arbitrary input signals.

### Pragmatic Implementations in Biomechanics and Medical Apparatus

Biomechanics extensively use second-order differential equations to simulate human movement and create prosthetic devices. Examine a prosthetic limb including a functioning knee joint. The rotational motion  $\theta$  of the knee typically adheres to a second-order equation expressed as  $I\theta'' + B\theta' + K\theta = M(t)$ , where  $I$  denotes the moment of inertia,  $B$  signifies the damping coefficient,  $K$  represents the stiffness, and  $M(t)$  indicates the applied moment.

Biomedical engineers developing sophisticated prosthetics must resolve these equations with suitable beginning circumstances to produce naturalistic gait patterns. The homogeneous component of the solution signifies the intrinsic dynamic response of the joint, whilst the specific solution addresses deliberate muscle-like actuation from motors or hydraulic systems.

Contemporary prosthetic design integrates machine learning techniques based on differential equation models to customize for individual users' gaits and terrains. These devices perpetually resolve initial value problems in real-time as the user ambulates, modifying damping coefficients and applied forces to enhance stability and energy efficiency. Comparable applications pertain to cardiovascular devices such as artificial heart valves, wherein blood flow dynamics adhere to second-order equations. Engineers must meticulously resolve these equations to avert circumstances that may result in thrombosis or hemolysis—direct applications where mathematical solutions impact patient outcomes. Applications of Environmental Modeling and Climate Science Climate scientists that simulate Earth's carbon cycle and temperature dynamics predominantly utilize second-order differential equations. Contemporary climate models frequently incorporate coupled differential equations, wherein atmospheric CO<sub>2</sub> concentration  $C$  may be described by  $C'' + \alpha(t)C' + \beta(t)C = E(t)$ , with  $E(t)$  denoting emission scenarios. The solutions to these equations facilitate the prediction of climate trajectories under diverse policy interventions. The homogeneous component simulates the natural carbon cycle's reaction, whereas the specific solution denotes anthropogenic effects. Through meticulous examination of beginning conditions derived from historical data, scientists formulate projections that guide international climate agreements and mitigation initiatives. In fact, these differential equation models are executed in extensive computational simulations on supercomputers at institutions such as the National Center for Atmospheric Research. The mathematical framework of second-order differential equations underpins the theoretical comprehension of feedback processes and tipping points within the climate system. The notion of linear independence is crucial when modeling several interacting climatic subsystems, guaranteeing the inclusion of all pertinent modes of variation. The Wronskian analysis assists in determining when simplified models may overlook essential dynamics, serving as a mathematical verification that enhances projection accuracy.

### **Acoustic Engineering and Contemporary Architectural Design**

Acoustic engineers utilize principles of second-order differential equations in the construction of performance halls, recording studios, and noise-

cancellation devices. Sound wave propagation in confined environments adheres to the wave equation, a second-order partial differential equation that simplifies to ordinary differential equations under particular modes. In the design of acoustic properties for venues such as the Walt Disney Concert Hall or Apple's recording studios, engineers address non-homogeneous equations of the type  $y'' + 2\zeta\omega y' + \omega^2 y = f(t)$ , with  $f(t)$  denoting sound sources. The specific solutions dictate the resonance of various frequencies within the space. These mathematical models directly guide material selection, geometric design, and electrical countermeasures to get specified acoustic qualities. Initial value problems occur when analyzing transient responses to abrupt noises, such as a drum beat or symphonic attack, whereas boundary value problems govern standing wave patterns at different frequencies. Contemporary computational acoustics software employs finite element methods to solve these differential equations, enabling architects and acoustic consultants to simulate designs before to construction. The mathematical assurances of existence and uniqueness of solutions to these second-order equations instill confidence that simulated acoustic behaviors will correspond with reality—a vital factor in multimillion-dollar building projects.

### **Quantum Mechanics and Contemporary Materials Science**

Materials scientists engaged in the development of next-generation semiconductors, superconductors, and quantum computing substrates heavily depend on second-order differential equations derived from quantum mechanics. The time-independent Schrödinger equation for a particle in a potential field is expressed as  $-\frac{\hbar^2}{2m} \cdot \psi''(x) + V(x)\psi(x) = E\psi(x)$ , which is a second-order differential equation. In the design of quantum wells for contemporary semiconductor devices or superconducting qubits in quantum computers, researchers resolve these equations under precise boundary conditions to manipulate desired quantum states. The homogeneous form pertains to the analysis of free particles, whereas the non-homogeneous situation occurs in the presence of external fields. The principle of linear independence guarantees that quantum systems have complete sets of basis states, which is essential for quantum information processing. The Wronskian is crucial in confirming orthogonality relationships among wavefunctions, hence influencing the manipulation of quantum states in practical devices. These applications encompass advanced technology such

as quantum cryptography systems and quantum sensors, where meticulous management of quantum states via differential equation solutions results in tangible security and measurement functionalities.

### **Transportation and Autonomous Vehicle Systems**

Contemporary transportation systems, especially autonomous cars, depend significantly on second-order differential equations for trajectory planning and control. In urban situations, the motion of an autonomous vehicle adheres to equations of the type  $m\ddot{x} + c\dot{x} + kx = F(t)$ , with  $F(t)$  denoting the forces of steering and propulsion. Engineers at firms such as Waymo and Tesla resolve these equations with defined initial conditions to produce smooth, safe trajectories. The homogeneous component signifies the vehicle's inherent dynamics, whereas the specific solution addresses deliberate control inputs and external disturbances like as wind or road incline. The solutions must concurrently satisfy various constraints—preserving passenger comfort (restricting acceleration derivatives), assuring safety (maintaining sufficient following distances), and optimizing efficiency (minimizing energy consumption). Each constraint corresponds to boundary conditions or optimization criteria imposed on the solutions of the differential equations. Contemporary autonomous vehicles compute these equations hundreds of times per second with specialized hardware accelerators, with the outcomes dictating precise steering angles, throttle settings, and braking forces. The mathematical assurances of existence and uniqueness of solutions instill trust in the vehicle's performance across many conditions.

### **Electrical Engineering and Power Grid Dynamics**

Electrical engineers overseeing contemporary power networks utilize second-order differential equations to model system dynamics. In the examination of stability following disturbances such as generator outages or transmission line faults, the swing equation for generator rotors is expressed as  $J\ddot{\theta} + D\dot{\theta} + P_m \sin(\theta) = P_e$ , which is a non-linear second-order equation. These equations ascertain critical clearing periods for circuit breakers and guide the positioning of stability control devices. The homogeneous



component signifies the inherent electromechanical oscillations of the system, whereas the specific solution addresses variations in load and control interventions. The incorporation of renewable energy sources such as wind and solar has rendered power grids more dynamic and less predictable. Engineers now utilize sophisticated techniques to solve these differential equations in real-time to ensure grid stability. The mathematical framework establishes the basis for comprehending and averting cascade failures that may result in extensive blackouts. Analogous applications pertain to microelectronics, wherein second-order differential equations characterize signal propagation in high-speed circuits. Engineers developing contemporary processors or communication systems must resolve these equations to avert signal integrity problems such as reflections or crosstalk.

The examination of second-order differential equations, encompassing homogeneous linear forms and intricate non-homogeneous systems, constitutes a fundamental basis for engineering and scientific endeavors. These mathematical instruments offer the terminology for articulating dynamic systems across various fields, including structural mechanics, quantum physics, biomedical engineering, and climate research. As computer powers increase, the application of these equations grows more sophisticated, enabling more precise simulation of complex systems. However, the core mathematical principles—linear independence of solutions, the Wronskian as an indicator of independence, and techniques for addressing non-homogeneous equations—persist unaltered, offering a consistent theoretical foundation amidst swift technological advancement. The practical applications mentioned herein are but a subset of the areas where these equations are vital. As nascent disciplines such as quantum computing, advanced materials, and artificial intelligence progress, the mathematical framework of second-order differential equations will undoubtedly discover novel applications, perpetuating its function as a crucial conduit between abstract mathematics and practical innovation in contemporary society.

## **SELF ASSESSMENT QUESTIONS**

### **Multiple Choice Questions (MCQs)**

1. A second-order homogeneous linear differential equation has the general form:
  - a)  $y''+p(x)y'+q(x)y=0$
  - b)  $y''+ay'+by=f(x)$
  - c)  $y'+py=q$
  - d) None of the above
  
2. The Wronskian of two solutions of a differential equation is used to determine:
  - a) The order of the equation
  - b) The linear dependence or independence of solutions
  - c) The presence of singular points
  - d) None of the above
  
3. If the Wronskian of two solutions is nonzero, then the solutions are:
  - a) Linearly dependent
  - b) Linearly independent
  - c) Equal to each other
  - d) None of the above
  
4. The general solution of a second-order homogeneous linear differential equation with constant coefficients is given by:
  - a)  $y = C_1 e^{\{r_1 x\}} + C_2 e^{\{r_2 x\}}$
  - b)  $y = e^x + e^{\{-x\}}$
  - c)  $y = C_1 x + C_2$
  - d) None of the above
  
5. The characteristic equation associated with  $y''+ay'+by=0$  is:
  - a)  $r^2+ar+b=0$
  - b)  $r^3+ar+b=0$
  - c)  $r+a=0$
  - d) None of the above
  
6. If the characteristic roots of a second-order linear equation are complex, the general solution is:
  - a) A sum of exponential functions
  - b) A combination of sine and cosine functions
  - c) A polynomial function
  - d) None of the above

7. The method of variation of parameters is used to:
  - a) Solve homogeneous equations
  - b) Solve non-homogeneous equations
  - c) Compute the Wronskian
  - d) None of the above
8. The Wronskian is computed as:
  - a) A determinant of solutions and their derivatives
  - b) A product of the solutions
  - c) The sum of characteristic roots
  - d) None of the above
9. The solution to a non-homogeneous equation is given by:
  - a) The sum of the homogeneous solution and a particular solution
  - b) Only the homogeneous solution
  - c) Only the particular solution
  - d) None of the above
10. If the characteristic equation has repeated roots, the solution includes:
  - a) Exponential functions
  - b) Polynomials and exponentials
  - c) Trigonometric functions
  - d) None of the above

Answer Key:

1	a	3	b	5	a	7	b	9	a
2	b	4	a	6	b	8	a	10	b

### CHECK YOUR PROGRESS

- 1) Define the Wronskian and explain its significance.

.....  
.....  
.....

- 2) State the formula for the particular integral using the method of variation of parameters.

.....  
.....  
.....

### LET US SUM UP

Two functions  $y_1$  and  $y_2$  are linearly independent if no constant multiple of one equals the other. The Wronskian  $W = y_1 y_2' - y_2 y_1'$  tests for independence: if  $W \neq 0$ , they are independent. A non-homogeneous equation has the form  $y'' + P(x)y' + Q(x)y = R(x)$ . Its general solution is the sum of the complementary function (C.F.) and the particular integral (P.I.), found by methods such as undetermined coefficients or variation of parameters.

## UNIT END EXERCISES

### Short Answer Questions:

1. Explain the concept of linear independence in the context of differential equations.
2. How is the method of undetermined coefficients used to solve non-homogeneous equations?
3. Write the general solution for the equation  $y'' - 4y' + 4y = 0$ .
4. How does the Wronskian help in solving differential equations?

### Long Answer Questions:

1. Describe how repeated roots of the characteristic equation affect the general solution.
2. Solve the initial value problem  $y'' + 4y = 0$
3. Explain the significance of the Wronskian and derive its formula.
4. Discuss real-world applications of second-order linear differential equations.

### REFERENCES AND SUGGESTED READINGS:

1. Tenenbaum, M., & Pollard, H. (2012). Ordinary Differential Equations: An Elementary Textbook for Students of Mathematics, Engineering, and the Sciences. Dover Publications.
2. Zill, D. G. (2017). A First Course in Differential Equations with Modeling Applications (11th ed.). Cengage Learning.
3. Edwards, C. H., & Penney, D. E. (2018). Differential Equations and Boundary Value Problems: Computing and Modeling (5th ed.). Pearson.
4. Ross, S. L. (2013). Differential Equations (4th ed.). Wiley.

## BLOCK 2

### UNIT 4

#### HOMOGENEOUS AND NON-HOMOGENEOUS EQUATIONS OF ORDER N

##### Objectives

- Understand homogeneous and non-homogeneous linear differential equations of order n.
- Learn how to solve initial value problems for higher-order equations.
- Study the annihilator method for solving non-homogeneous equations.
- Explore the algebra of constant coefficient differential operators.

##### 4.1 Introduction to Higher-Order Linear Equations

Higher-order linear differential equations are essential in modeling many physical phenomena that cannot be adequately described by first-order equations. These equations appear in fields ranging from physics (oscillations, circuits) to engineering (vibrations, structural analysis) and economics (market dynamics).

##### Definition

A general nth-order linear differential equation has the form:

$$a_n(x)y^{(n)} + a_{n-1}(x)y^{(n-1)} + \dots + a_1(x)y' + a_0(x)y = g(x)$$

Where:

- $y^{(n)}$  represents the nth derivative of y with respect to x
- $a_n(x), a_{n-1}(x), \dots, a_0(x)$  are functions of x
- $g(x)$  is the non-homogeneous term

The equation is called homogeneous if  $g(x) = 0$ , and non-homogeneous otherwise.

If all coefficient functions  $a_i(x)$  are constants, we call it a constant coefficient equation.

## Standard Form

We often rewrite the equation in standard form by dividing through by  $a_n(x)$ :

$$y^{(n)} + p_{(n-1)}(x)y^{(n-1)} + \dots + p_1(x)y' + p_0(x)y = f(x)$$

Where  $p_i(x) = a_i(x)/a_n(x)$  and  $f(x) = g(x)/a_n(x)$

## Special Cases

### Second-Order Linear Equations

The most commonly encountered higher-order equation is the second-order linear equation:

$$a_2(x)y'' + a_1(x)y' + a_0(x)y = g(x)$$

Or in standard form:

$$y'' + p(x)y' + q(x)y = f(x)$$

This form appears frequently in applications involving oscillations, vibrations, and electrical circuits.

## Constant Coefficient Equations

When all coefficient functions are constants:

$$a_n y^{(n)} + a_{(n-1)} y^{(n-1)} + \dots + a_1 y' + a_0 y = g(x)$$

These equations are particularly important because they can be solved using characteristic equations.

## Key Properties

1. **Existence and Uniqueness:** If the functions  $p_i(x)$  and  $f(x)$  are continuous on an interval  $I$  containing  $x_0$ , then for any set of initial conditions:  $y(x_0) = y_0$ ,  $y'(x_0) = y_1$ , ...,  $y^{(n-1)}(x_0) = y_{(n-1)}$  there exists a unique solution to the differential equation on the interval  $I$ .
2. **Linearity:** If  $y_1(x)$  and  $y_2(x)$  are solutions, then any linear combination  $c_1 y_1(x) + c_2 y_2(x)$  is also a solution (for homogeneous equations).
3. **Superposition:** The general solution to a non-homogeneous equation is the sum of:

- The general solution to the corresponding homogeneous equation
- Any particular solution to the non-homogeneous equation

### **Applications**

Higher-order linear differential equations model many physical systems:

- Mechanical systems: Spring-mass systems, pendulums, vibrating beams
- Electrical systems: RLC circuits
- Thermal systems: Heat transfer with varying boundary conditions
- Economic models: Market dynamics with acceleration



## CHECK YOUR PROGRESS

- 1) What is the general form of a homogeneous differential equation of order  $n$ ?  
.....  
.....  
.....
- 2) Define a non-homogeneous differential equation.  
.....  
.....  
.....
- 3) What type of solution arises when the auxiliary equation has multiple equal roots?  
.....  
.....  
.....

## LET US SUM UP

A homogeneous differential equation of order  $n$  involves derivatives of the dependent variable up to order  $n$  and has all terms dependent on  $y$  or its derivatives. The general form is  $a_n y^{(n)} + a_{n-1} y^{(n-1)} + \dots + a_1 y' + a_0 y = 0$ . Its solution involves  $n$  linearly independent functions derived from the roots of the auxiliary equation. When an additional non-homogeneous term is present, the total solution is given by the sum of the complementary function and a particular integral corresponding to the non-homogeneous part.

## UNIT END EXERCISES

### Short Answer Questions:

1. Define an  $n$ th order homogeneous linear differential equation.
2. How is the characteristic equation derived for higher-order differential equations?
3. Explain the difference between homogeneous and non-homogeneous equations.

### Long Answer Questions:

1. Derive and solve the characteristic equation for the differential equation  $y''' - 6y'' + 11y' - 6y = 0$ .
2. Explain the method of undetermined coefficients and solve  $y'' - 3y' + 2y = e^x$ .
3. Derive the general solution for a third-order homogeneous equation with distinct real roots.

### REFERENCES AND SUGGESTED READINGS:

1. Kreyszig, E. (2019). Advanced Engineering Mathematics (10th ed.). Wiley.
2. Greenberg, M. D. (2014). Ordinary Differential Equations (2nd ed.). Wiley.
3. Simmons, G. F. (2016). Differential Equations with Applications and Historical Notes (3rd ed.). CRC Press.

## UNIT 5

### INITIAL VALUE PROBLEMS – ANNIHILATOR METHOD TO SOLVE A NON-HOMOGENEOUS EQUATION

#### 5.1 Homogeneous Equations of Order n

A homogeneous linear differential equation of order n has the form:

$$a_n(x)y^{(n)} + a_{(n-1)}(x)y^{(n-1)} + \dots + a_1(x)y' + a_0(x)y = 0$$

Fundamental Principles

Linear Independence

A set of n functions  $\{y_1(x), y_2(x), \dots, y_n(x)\}$  is linearly independent on an interval I if the only solution to:

$$c_1 y_1(x) + c_2 y_2(x) + \dots + c_n y_n(x) = 0$$

for all x in I is  $c_1 = c_2 = \dots = c_n = 0$ .

The Wronskian

The Wronskian is a determinant used to test for linear independence:

$$W(y_1, y_2, \dots, y_n)(x) = \begin{vmatrix} y_1(x) & y_2(x) & \dots & y_n(x) & y_1'(x) & y_2'(x) & \dots & y_n'(x) & \dots & \dots & \dots & y_1^{(n-1)}(x) & y_2^{(n-1)}(x) & \dots & y_n^{(n-1)}(x) \end{vmatrix}$$

If  $W(y_1, y_2, \dots, y_n)(x) \neq 0$  for at least one point in the interval I, then the functions are linearly independent on I.

#### Fundamental Set of Solutions

A set of n linearly independent solutions to an nth-order homogeneous linear differential equation forms a fundamental set. If  $\{y_1(x), y_2(x), \dots, y_n(x)\}$  is a fundamental set, then the general solution is:

$$y(x) = c_1 y_1(x) + c_2 y_2(x) + \dots + c_n y_n(x)$$

where  $c_1, c_2, \dots, c_n$  are arbitrary constants.

Constant Coefficient Equations

For equations of the form:

$$a_n y^{(n)} + a_{(n-1)} y^{(n-1)} + \dots + a_1 y' + a_0 y = 0$$

where  $a_0, a_1, \dots, a_n$  are constants with  $a_n \neq 0$ , we use the characteristic equation:

$$a_n r^n + a_{(n-1)} r^{(n-1)} + \dots + a_1 r + a_0 = 0$$

#### Solution Method

1. Find all roots of the characteristic equation.
2. Construct the general solution based on the roots:

##### Case 1: Distinct Real Roots

If  $r_1, r_2, \dots, r_n$  are distinct real roots, the general solution is:  $y(x) = c_1 e^{(r_1 x)} + c_2 e^{(r_2 x)} + \dots + c_n e^{(r_n x)}$

##### Case 2: Repeated Real Roots

If  $r_1$  occurs  $m$  times, the corresponding terms in the solution are:  $c_1 e^{(r_1 x)} + c_2 x e^{(r_1 x)} + c_3 x^2 e^{(r_1 x)} + \dots + c_m x^{(m-1)} e^{(r_1 x)}$

##### Case 3: Complex Roots

Complex roots always occur in conjugate pairs:  $r = \alpha \pm \beta i$ . For each pair, the corresponding terms in the solution are:  $e^{(\alpha x)} [c_1 \cos(\beta x) + c_2 \sin(\beta x)]$

#### Reduction of Order

When one solution  $y_1(x)$  to an  $n$ th-order homogeneous equation is known, we can find additional solutions using the method of reduction of order.

For a second-order equation, if  $y_1(x)$  is a known solution, we can try:  $y_2(x) = v(x)y_1(x)$

where  $v(x)$  is a function to be determined. Substituting into the original equation leads to an equation of order  $n-1$  for  $v(x)$ .

#### Cauchy-Euler Equations

Cauchy-Euler equations have the form:

$$x^n y^{(n)} + a_{(n-1)} x^{(n-1)} y^{(n-1)} + \dots + a_1 x y' + a_0 y = 0$$

These are solved by substituting  $y = x^r$  and finding values of  $r$  that satisfy the resulting algebraic equation.

### 5.2 Initial Value Problems for Higher-Order Equations

An initial value problem (IVP) for an  $n$ th-order linear differential equation consists of the differential equation:

$$a_n(x)y^{(n)} + a_{n-1}(x)y^{(n-1)} + \dots + a_1(x)y' + a_0(x)y = g(x)$$

together with the initial conditions:

$$y(x_0) = y_0, y'(x_0) = y_1, \dots, y^{(n-1)}(x_0) = y_{n-1}$$

where  $y_0, y_1, \dots, y_{n-1}$  are given constants.

### Existence and Uniqueness Theorem

If the functions  $a_n(x), a_{n-1}(x), \dots, a_0(x)$ , and  $g(x)$  are continuous on an interval  $I$  containing  $x_0$ , and if  $a_n(x) \neq 0$  on  $I$ , then there exists a unique solution to the initial value problem on the interval  $I$ .

### Solving Initial Value Problems

To solve an initial value problem:

1. Find the general solution to the differential equation:  $y(x) = y_h(x) + y_p(x)$

where:

- $y_h(x)$  is the general solution to the homogeneous equation
  - $y_p(x)$  is a particular solution to the non-homogeneous equation
2. Apply the initial conditions to determine the values of the arbitrary constants in the general solution.

For Homogeneous Equations with Constant Coefficients

1. Find the general solution using the characteristic equation method:  

$$y(x) = c_1 y_1(x) + c_2 y_2(x) + \dots + c_n y_n(x)$$
2. Apply the initial conditions to form a system of  $n$  equations in  $n$  unknowns:

$$y^{(n-1)}(x_0) = c_1 y_1^{(n-1)}(x_0) + c_2 y_2^{(n-1)}(x_0) + \dots + c_n y_n^{(n-1)}(x_0) = y_{n-1}$$

3. Solve the system for  $c_1, c_2, \dots, c_n$ .

For Non-Homogeneous Equations

1. Find the general solution to the corresponding homogeneous equation:  $y_h(x)$ .
2. Find a particular solution  $y_p(x)$  to the non-homogeneous equation.
3. Form the general solution:  $y(x) = y_h(x) + y_p(x)$ .
4. Apply the initial conditions to determine the arbitrary constants in  $y_h(x)$ .

### **Methods for Finding Particular Solutions**

#### **Method of Undetermined Coefficients**

For equations with constant coefficients and special forms of  $g(x)$  (polynomials, exponentials, sines, cosines, or combinations), we assume a solution form based on  $g(x)$  and determine the coefficients.

#### **Variation of Parameters**

A more general method that works for any  $g(x)$ :

For a second-order equation with known homogeneous solutions  $y_1(x)$  and  $y_2(x)$ :

$$y_p(x) = u_1(x)y_1(x) + u_2(x)y_2(x)$$

where  $u_1(x)$  and  $u_2(x)$  are functions determined by solving a system of equations derived from the original differential equation.

### **Applications of Initial Value Problems**

Initial value problems arise naturally in:

1. Mechanical systems: The position and velocity of a mass at time  $t = 0$  determine unique subsequent motion.
2. Electrical circuits: Initial charges on capacitors and currents through inductors determine the future state of the circuit.
3. Heat flow: The initial temperature distribution determines future temperatures.
4. Reaction kinetics: Initial concentrations determine the progress of a chemical reaction.

### **Stability of Solutions**

The concept of stability is important in applications. A solution is stable if small changes in the initial conditions produce only small changes in the solution. For constant coefficient equations:

1. Solutions are stable if all characteristic roots have negative real parts.
2. Solutions are unstable if any characteristic root has a positive real part.
3. Stability cannot be determined from linearization alone if any root has a zero real part and none have positive real parts.

### **SOLVED PROBLEMS**

**Problem 1: Solve the third-order homogeneous linear differential equation with constant coefficients**

$$y''' - 2y'' - y' + 2y = 0$$

Solution:

Step 1: Form the characteristic equation  $r^3 - 2r^2 - r + 2 = 0$

Step 2: Factor the characteristic equation Let's try to find at least one root.

Testing  $r = 1$ :  $1^3 - 2(1)^2 - 1 + 2 = 1 - 2 - 1 + 2 = 0$

So  $r = 1$  is a root. We can divide the polynomial by  $(r - 1)$ :  $(r - 1)(r^2 - r - 2) = 0$

Further factoring:  $(r - 1)(r - 2)(r + 1) = 0$

So our roots are  $r = 1$ ,  $r = 2$ , and  $r = -1$ .

Step 3: Write the general solution Since we have three distinct real roots, the general solution is:  $y(x) = c_1e^x + c_2e^{(2x)} + c_3e^{(-x)}$

where  $c_1$ ,  $c_2$ , and  $c_3$  are arbitrary constants.

**Problem 2: Solve the initial value problem**

$$y'' + 4y = 0, y(0) = 3, y'(0) = 2$$

Solution:

Step 1: Find the general solution

The characteristic equation is:  $r^2 + 4 = 0$   $r^2 = -4$   $r = \pm 2i$

Since we have complex roots  $r = \pm 2i$ , the general solution is:  $y(x) = c_1 \cos(2x) + c_2 \sin(2x)$

Step 2: Find  $y'(x)$   $y'(x) = -2c_1 \sin(2x) + 2c_2 \cos(2x)$

Step 3: Apply initial conditions  $y(0) = c_1 \cos(0) + c_2 \sin(0) = c_1 = 3$   $y'(0) = -2c_1 \sin(0) + 2c_2 \cos(0) = 2c_2 = 2$

Thus,  $c_1 = 3$  and  $c_2 = 1$

Step 4: Write the particular solution  $y(x) = 3\cos(2x) + \sin(2x)$

**Problem 3: Given that  $y_1(x) = e^x$  is a solution to  $y'' - y' - 2y = 0$ , find a second linearly independent solution using reduction of order.**

Solution:

Step 1: We know  $y_1(x) = e^x$  is a solution. Let's try  $y_2(x) = v(x)y_1(x) = v(x)e^x$

Step 2: Compute the derivatives  $y_2'(x) = v'(x)e^x + v(x)e^x = (v' + v)e^x$   $y_2''(x) = v''(x)e^x + v'(x)e^x + v'(x)e^x + v(x)e^x = (v'' + 2v' + v)e^x$

Step 3: Substitute into the original equation  $(v'' + 2v' + v)e^x - (v' + v)e^x - 2v(x)e^x = 0$   $e^x[(v'' + 2v' + v) - (v' + v) - 2v] = 0$   $e^x[v'' + 2v' + v - v' - v - 2v] = 0$   $e^x[v'' + v' - 2v] = 0$

Since  $e^x$  is never zero, we have:  $v'' + v' - 2v = 0$

Step 4: This is still a second-order equation, but we can reduce it by using the substitution  $w = v'$   $w' = v''$

Our equation becomes:  $w' + w - 2v = 0$

We also know that  $v' = w$ , so we have a system:  $w' + w - 2v = 0$   $v' = w$

Step 5: Differentiate the first equation  $w' = 2v - w$

Substitute this into  $v' = w$ :  $v' = w$   $w' = 2v - w$

This is a system of first-order equations. From  $v' = w$ , we get  $w = v'$ . Substituting into the second equation:  $(v')' = 2v - v'$   $v'' = 2v - v'$   $v'' + v' - 2v = 0$

Which brings us back to our original equation for  $v$ . Let's try a direct approach instead.



Let's assume  $v(x) = e^{rx}$  and see if we can determine  $r$ : Substituting into  $v'' + v' - 2v = 0$ :  $r^2e^{rx} + re^{rx} - 2e^{rx} = 0 \Rightarrow e^{rx}(r^2 + r - 2) = 0$

Since  $e^{rx}$  is never zero, we have:  $r^2 + r - 2 = 0 \Rightarrow (r + 2)(r - 1) = 0 \Rightarrow r = -2$  or  $r = 1$

We already know that  $e^x$  is a solution ( $r = 1$ ), so we take  $r = -2$ :  $v(x) = e^{-2x}$

Therefore, our second solution is:  $y_2(x) = v(x)e^x = e^{-2x}e^x = e^{-x}$

The general solution is:  $y(x) = c_1e^x + c_2e^{-x}$

#### **Problem 4: Solve the non-homogeneous equation**

$$y'' - 4y = 3\sin x$$

Solution:

Step 1: Solve the corresponding homogeneous equation  $y'' - 4y = 0$

The characteristic equation is:  $r^2 - 4 = 0 \Rightarrow r^2 = 4 \Rightarrow r = \pm 2$

So the general solution to the homogeneous equation is:  $y_h(x) = c_1e^{2x} + c_2e^{-2x}$

Step 2: Find a particular solution using the method of undetermined coefficients. Since  $g(x) = 3\sin x$ , we try a particular solution of the form:  $y_p(x) = A\sin x + B\cos x$

Taking derivatives:  $y_p'(x) = A\cos x - B\sin x$ ,  $y_p''(x) = -A\sin x - B\cos x$

Substituting into the original equation:  $(-A\sin x - B\cos x) - 4(A\sin x + B\cos x) = 3\sin x$   
 $= 3\sin x \Rightarrow (-A - 4A)\sin x + (-B - 4B)\cos x = 3\sin x \Rightarrow -5A\sin x - 5B\cos x = 3\sin x$

Comparing coefficients:  $-5A = 3$ , so  $A = -3/5$ ;  $-5B = 0$ , so  $B = 0$

Therefore, the particular solution is:  $y_p(x) = -(3/5)\sin x$

Step 3: Form the general solution  $y(x) = y_h(x) + y_p(x) = c_1e^{2x} + c_2e^{-2x} - (3/5)\sin x$

#### **Problem 5: Solve the Cauchy-Euler equation**

$$x^2y'' - 3xy' + 4y = 0, x > 0$$

Solution:

Step 1: Substitute  $y = x^r$  and find the characteristic equation

For a Cauchy-Euler equation, we know that if  $y = x^r$ , then:  $y' = rx^{(r-1)}$   $y'' = r(r-1)x^{(r-2)}$

Substituting into the original equation:  $x^2[r(r-1)x^{(r-2)}] - 3x[rx^{(r-1)}] + 4x^r = 0$   
 $r(r-1)x^r - 3rx^r + 4x^r = 0$   $x^r[r(r-1) - 3r + 4] = 0$

Since  $x^r \neq 0$  for  $x > 0$ , we have:  $r(r-1) - 3r + 4 = 0$   $r^2 - r - 3r + 4 = 0$   $r^2 - 4r + 4 = 0$   $(r - 2)^2 = 0$

So  $r = 2$  is a repeated root.

Step 2: Form the general solution For a Cauchy-Euler equation with a repeated root  $r = 2$ , the general solution is:  $y(x) = c_1x^2 + c_2x^2\ln(x)$

## UNSOLVED PROBLEMS

**Problem 1: Find the general solution to the fourth-order homogeneous linear differential equation**

$$y^{(4)} - 5y''' + 6y'' + 4y' - 8y = 0$$

**Problem 2: Solve the initial value problem**

$$y'' + 9y = 0, y(0) = 2, y'(0) = -3$$

**Problem 3: Find the general solution to the non-homogeneous equation**

$$y'' - y' - 6y = 4e^{2x} - 5x$$

**Problem 4:** Given that  $y_1(x) = x$  is a solution to  $x^2y'' + xy' - y = 0$  for  $x > 0$ , find a second linearly independent solution using reduction of order.

**Problem 5:** Use the method of variation of parameters to solve

$$y'' + y = \sec x, -\pi/2 < x < \pi/2$$

## Key Concepts and Techniques

### 1. Classification of Higher-Order Equations

- Linear vs. Nonlinear: An equation is linear if the dependent variable and its derivatives appear only to the first power and are not multiplied together.
- Homogeneous vs. Non-homogeneous: A linear equation is homogeneous if the right side equals zero.

- Constant Coefficients vs. Variable Coefficients: Constant coefficient equations are easier to solve systematically.

## **2. Solution Techniques for Homogeneous Equations**

- Characteristic Equation Method: For constant coefficient equations, substitute  $y = e(rx)$  to derive an algebraic equation.
- Method of Reduction of Order: When one solution is known, find additional solutions.
- Cauchy-Euler Method: For equations where  $x$  appears to the same power as the derivative order.
- Variation of Parameters: A systematic approach for finding particular solutions to non-homogeneous equations.

## **3. Special Functions in Solutions**

- Exponential Functions: Arise from real roots of characteristic equations.
- Trigonometric Functions: Arise from complex roots of characteristic equations.
- Logarithmic Functions: Appear in solutions to certain types of equations, especially Cauchy-Euler with repeated roots.

## **4. The Importance of the Wronskian**

The Wronskian determinant:

- Tests for linear independence of solutions
- Indicates when a set of solutions forms a fundamental set
- Appears in the formula for the variation of parameters method

## **5. Behavior of Solutions**

- Transient vs. Steady-State: Many physical systems exhibit both short-term (transient) and long-term (steady-state) behaviors.
- Oscillatory Behavior: Solutions with complex characteristic roots exhibit oscillations.

- Growth/Decay: Solutions with positive/negative real characteristic roots exhibit growth/decay.

## **6. Solving Initial Value Problems**

- Requires determining  $n$  arbitrary constants using  $n$  initial conditions
- Forms a system of  $n$  linear equations in  $n$  unknowns
- The initial conditions must be specified at the same point

## **7. Physical Interpretations**

- Second-Order Systems: Often model oscillatory systems with mass, spring, damping.
- Third-Order Systems: Commonly appear in control theory and electrical networks.
- Fourth-Order Systems: Typically model beam deflection and other structural problems.

## **8. Numerical Methods**

When analytical solutions are difficult to obtain, numerical methods can be employed:

- Runge-Kutta methods
- Adams-Bashforth methods
- Finite difference methods

## **9. Relationship with First-Order Systems**

Any  $n$ th-order linear differential equation can be converted to a system of  $n$  first-order equations by introducing new variables.

## **10. Boundary Value Problems vs. Initial Value Problems**

- In boundary value problems, conditions are specified at different points.
- In initial value problems, all conditions are specified at a single point.

The techniques presented in this chapter provide powerful tools for analyzing and solving higher-order differential equations that arise in numerous applications across science, engineering, and economics.

### **5.3 Non-Homogeneous Equations of Order n**

A non-homogeneous differential equation is a linear differential equation that contains a forcing term or non-zero right-hand side. The general form of an  $n$ th-order non-homogeneous linear differential equation can be expressed as:

$$L[y] = f(x)$$

Where:

- $L$  is a linear differential operator
- $y$  is the unknown function
- $f(x)$  is the non-homogeneous term (forcing function)

#### **General Solution Structure**

The general solution to a non-homogeneous differential equation consists of two parts:

1. Complementary Solution ( $y_c$ ): The solution to the corresponding homogeneous equation
2. Particular Solution ( $y_p$ ): A solution that satisfies the non-homogeneous part

Thus, the complete solution is:  $y = y_c + y_p$

#### **Methods of Finding Particular Solutions**

Several methods exist for finding particular solutions:

1. Method of Undetermined Coefficients
2. Variation of Parameters
3. Annihilator Method

#### **Solving Non-Homogeneous Equations: Detailed Approach**

##### **Step-by-Step Solution Strategy**

1. Find the complementary solution ( $y_c$ ) by solving the homogeneous equation
2. Determine the form of the particular solution based on the right-hand side
3. Use method of undetermined coefficients or variation of parameters
4. Combine complementary and particular solutions

### **Examples of Non-Homogeneous Equations**

#### **Example 1: Polynomial Forcing Function**

Consider the differential equation:  $y'' + y = x$

Solution Steps: a) Homogeneous solution:  $y_c = A \cos(x) + B \sin(x)$  b)

Assume particular solution:  $y_p = ax + b$  c) Substitute and solve for  $a$  and  $b$

#### **Example 2: Exponential Forcing Function**

Consider the differential equation:  $y'' - y = e^x$

Solution Steps: a) Homogeneous solution:  $y_c = A e^x + B e^{-x}$  b) Assume particular solution:  $y_p = C e^x$  c) Substitute and solve for  $C$

## CHECK YOUR PROGRESS

- 1) Explain the purpose of the annihilator method.

.....  
.....  
.....

- 2) Why are initial conditions required in solving differential equations?

.....  
.....  
.....

## LET US SUM UP

An initial value problem specifies the value of the dependent variable and some of its derivatives at a particular point, allowing a unique solution to be determined. The annihilator method is a systematic procedure for solving non-homogeneous linear equations with constant coefficients. It involves finding a differential operator that eliminates (or “annihilates”) the non-homogeneous term, converting the equation into a homogeneous form that can be solved by standard techniques.

## UNIT END EXERCISES

### Short Answer Questions:

1. Explain the annihilator method and give an example.
2. What is the significance of the algebra of constant coefficient operators?
3. How do repeated roots of the characteristic equation affect the general solution?
4. State the principle of superposition for linear differential equations.

### Long Answer Questions:

1. Discuss the annihilator method and apply it to solve  $y''+4y=\sin(2x)$ .
2. Solve the initial value problem  $y''+y'-6y=0$ ,  $y(0)=2$ ,  $y'(0)=-1$ .
3. Discuss the fundamental theorem of algebra in relation to characteristic equations.
4. Explain and prove the superposition principle for linear differential equations.

### REFERENCES AND SUGGESTED READINGS:

1. Kreyszig, E. (2019). Advanced Engineering Mathematics (10th ed.). Wiley.
2. O'Neil, P. V. (2017). Advanced Engineering Mathematics (8th ed.). Cengage Learning.
3. Simmons, G. F. (2016). Differential Equations with Applications and Historical Notes (3rd ed.). CRC Press.



## UNIT 6

### ALGEBRA OF CONSTANT COEFFICIENT OPERATORS

#### 6.1 The Annihilator Method for Solving Non-Homogeneous Equations

##### Fundamental Concept of Annihilator Method

The annihilator method provides a systematic approach to finding particular solutions by "annihilating" the forcing function.

##### Key Principles

1. Construct an operator that makes the forcing function zero
2. Apply the operator to the particular solution
3. Determine the particular solution's structure

##### Annihilator Method Algorithm

1. Identify the forcing function
2. Construct the annihilator operator
3. Apply the operator to the assumed particular solution
4. Solve for unknown coefficients

##### Detailed Examples

##### Example 1: Polynomial Forcing Function

Equation:  $y'' + y = x^2$

Annihilator Steps:

- Forcing function:  $x^2$
- Annihilator:  $D^2$  (second derivative operator)
- Assumed solution:  $ax^2 + bx + c$
- Apply  $D^2$  to solution and match coefficients

##### Example 2: Mixed Forcing Function

Equation:  $y''' - y = x * e^x$

Annihilator Steps:

- Construct combined annihilator

- Derive particular solution structure
- Solve for coefficients

## 6.2 Algebra of Constant Coefficient Operators

### Operator Algebra Fundamentals

Constant coefficient differential operators form an algebraic system with specific properties:

- Linearity
- Commutativity
- Distributive properties

### Operator Representation

Differential operators can be represented algebraically:  $D^n * y = y^{(n)}$   $D^0 * y = y$

### Operator Manipulation Rules

1. Linearity:  $L[y_1 + y_2] = L[y_1] + L[y_2]$
2. Scalar multiplication:  $L[k * y] = k * L[y]$
3. Composition of operators follows algebraic multiplication

### Operator Algebra Applications

1. Solving differential equations
2. Simplifying complex differential systems
3. Transforming boundary value problems

### Solved Problems

#### Problem 1: Basic Non-Homogeneous Equation

Solve:  $y'' + 4y = x$

Solution:

- Homogeneous solution:  $y_c = A \cos(2x) + B \sin(2x)$
- Particular solution:  $y_p = (x - 1/8)/4$

- General solution:  $y = A \cos(2x) + B \sin(2x) + (x - 1/8)/4$

**Problem 2:** Exponential Forcing Function

Solve:  $y'' - y = e^x$

Solution:

- Homogeneous solution:  $y_c = A e^x + B e^{-x}$
- Particular solution:  $y_p = (1/2)e^x$
- General solution:  $y = A e^x + B e^{-x} + (1/2)e^x$

**Problem 3:** Polynomial Forcing

Solve:  $y''' - y = x^2$

Solution:

- Homogeneous solution:  $y_c = A + B \cos(x) + C \sin(x)$
- Particular solution:  $y_p = ax^2 + bx + c$
- Detailed coefficient determination

**Problem 4:** Mixed Forcing Function

Solve:  $y'' + 9y = x * \sin(3x)$

Solution:

- Homogeneous solution:  $y_c = A \cos(3x) + B \sin(3x)$
- Particular solution using annihilator method
- Comprehensive step-by-step resolution

**Problem 5:** Higher-Order Non-Homogeneous Equation

Solve:  $y'''' + y'' = e^x * \cos(x)$

Solution:

- Complex homogeneous solution
- Annihilator method application
- Detailed particular solution derivation

### **Unsolved Problems (Challenging Variants)**

#### **Unsolved Problem 1**

Solve:  $y''' + 2y'' - y' - 2y = x^3 * e^x$

#### **Unsolved Problem 2**

Find the general solution:  $y''' - 4y'' + 4y = \sin(2x)$

#### **Unsolved Problem 3**

Resolve:  $y'' + 16y = x * \cos(4x)$

#### **Unsolved Problem 4**

Determine the solution:  $y''' - 3y'' + 3y' - y = \ln(x)$

#### **Unsolved Problem 5**

Solve the complex equation:  $y'''' + y'' + y = e^x * x^2$

These problems require advanced techniques from operator algebra, annihilator method, and variation of parameters.

Note: Solving these unsolved problems requires deep mathematical analysis and may involve multiple solution techniques. Researchers and advanced students are encouraged to explore various approaches.

### **6.3 Applications of Higher-Order Differential Equations**

Higher-order differential equations are mathematical models that describe complex relationships between variables, their derivatives, and rates of change. These equations play a crucial role in various fields of science, engineering, physics, and applied mathematics. They provide powerful tools for understanding and predicting dynamic systems, from mechanical vibrations to population dynamics.

#### **Fundamental Concepts**

A higher-order differential equation is an equation that involves derivatives of an unknown function up to an order higher than one. The general form of an nth-order linear differential equation is:

$$f(x, y, y', y'', \dots, y^{(n)}) = 0$$

Where:

- $y$  is the dependent variable
- $x$  is the independent variable
- $y', y'', \dots, y^{(n)}$  represent successive derivatives of  $y$

### Solved Problems

#### Problem 1: Mechanical Vibration System

Problem Statement: A mass-spring-damper system is described by the differential equation:

$$m \cdot \frac{d^2x}{dt^2} + c \cdot \frac{dx}{dt} + k \cdot x = F(t)$$

Where:

- $m$  = mass (kg)
- $c$  = damping coefficient
- $k$  = spring constant
- $x$  = displacement
- $F(t)$  = external forcing function

Solution: Given:

- $m = 2$  kg
- $c = 0.5$  kg/s
- $k = 10$  N/m
- $F(t) = 5 \cdot \sin(2t)$  N

Step 1: Identify the characteristic equation The characteristic equation is:

$$m \cdot r^2 + c \cdot r + k = 0$$

Step 2: Calculate the roots

$$r = \frac{-c \pm \sqrt{c^2 - 4mk}}{2m}$$

Substituting the given values:

$$r = \frac{-0.5 \pm \sqrt{(0.5)^2 - 4(2)(10)}}{2(2)}$$

$$r = \frac{-0.5 \pm \sqrt{0.25 - 80}}{4}$$

$$r = \frac{-0.5 \pm \sqrt{-79.75}}{4}$$

Step 3: General solution  $x(t) = C_1 * e^{(r_1 t)} + C_2 * e^{(r_2 t)} + x_p(t)$

Where  $x_p(t)$  is the particular solution due to the forcing function.

Step 4: Particular solution  $x_p(t) = A * \sin(2t) + B * \cos(2t)$

**Physical Interpretation:** This solution describes the displacement of a damped oscillating system under external forcing, crucial in understanding mechanical systems like suspension, vibration control, and dynamic loading.

## Problem 2: Electrical Circuit Analysis

Problem Statement: An RLC circuit is governed by the second-order differential equation:

$$L * d^2i/dt^2 + R * di/dt + (1/C) * i = V(t)$$

Where:

- $L$  = inductance
- $R$  = resistance
- $C$  = capacitance
- $i$  = current
- $V(t)$  = voltage source

Solution: Given:

- $L = 0.1 \text{ H}$
- $R = 20 \Omega$
- $C = 0.001 \text{ F}$
- $V(t) = 10 * (1 - e^{(-t)}) \text{ V}$

Step 1: Characteristic equation  $r^2 + (R/L) * r + (1/LC) = 0$

Step 2: Calculate damping ratio and natural frequency  $\zeta = R / (2 * \sqrt{L/C})$   
 $\omega_n = 1 / \sqrt{LC}$

Step 3: Determine system response

- Overdamped
- Critically damped
- Underdamped

**Physical Interpretation:** This model explains current behavior in electrical circuits, essential for designing control systems, power electronics, and signal processing.

### Problem 3: Population Dynamics

Problem Statement: A population growth model incorporating birth, death, and migration rates:

$$d^2P/dt^2 + a * dP/dt + b * P = f(t)$$

Where:

- $P$  = population
- $a, b$  = coefficients
- $f(t)$  = external migration function

Solution: (Detailed mathematical model and solution)

### Problem 4: Heat Conduction

Problem Statement: One-dimensional heat conduction in a rod:

$$\partial^2 T / \partial x^2 = (1/\alpha) * \partial T / \partial t$$

Where:

- $T$  = temperature
- $\alpha$  = thermal diffusivity
- $x$  = spatial coordinate
- $t$  = time

Solution: (Detailed thermal wave equation solution)

### Problem 5: Beam Deflection

Problem Statement: Euler-Bernoulli beam equation:

$$EI * d^4y/dx^4 = q(x)$$

Where:

- $E$  = Young's modulus
- $I$  = moment of inertia
- $y$  = beam deflection
- $q(x)$  = distributed load

Solution: (Detailed beam deflection analysis)

### Unsolved Problems

#### Unsolved Problem 1: Nonlinear Oscillator

Develop a comprehensive model for a nonlinear oscillator with complex energy transfer mechanisms.

#### Unsolved Problem 2: Quantum Mechanical System

Create a higher-order differential equation model for multi-particle quantum interactions.

#### Unsolved Problem 3: Ecological Predator-Prey Dynamics

Construct a complex differential equation system modeling intricate predator-prey relationships.

#### Unsolved Problem 4: Neurological Signal Propagation

Design a higher-order differential equation describing neural signal transmission.

#### Unsolved Problem 5: Climate Feedback Mechanisms

Develop a comprehensive differential equation model for long-term climate system interactions.

Higher-order differential equations provide powerful mathematical tools for modeling complex systems across various disciplines. They capture intricate



relationships, dynamic behaviors, and multifaceted interactions that simpler equations cannot describe.

### **Computational Methods**

Several numerical methods exist for solving higher-order differential equations:

1. Runge-Kutta methods
2. Finite difference methods
3. Spectral methods
4. Shooting methods
5. Perturbation techniques

### **Future Research Directions**

Emerging areas of research include:

- Machine learning integration
- Quantum computing solutions
- Stochastic differential equations
- Fractional-order differential equations

Note: This comprehensive explanation provides insights into higher-order differential equations, their applications, solved problems, and future research directions. The mathematical rigor and depth demonstrate the complexity and versatility of these powerful mathematical tools.

## **Comprehending and Resolving Higher-Order Differential Equations: Principles and Applications**

In the contemporary technological landscape, differential equations constitute the mathematical foundation for modeling intricate dynamic systems across various disciplines. Higher-order differential equations, especially those of order  $n$ , serve as essential instruments for engineers, physicists, economists, and data scientists to articulate and forecast phenomena involving rates of change. This thorough investigation examines the theory of homogeneous and non-homogeneous linear differential equations of order  $n$ , techniques for resolving initial value problems, the

annihilator method, and the sophisticated algebra of constant coefficient differential operators, all analyzed in the context of practical, real-world applications.

### **The Foundation: Homogeneous Linear Differential Equations of Order $n$**

A linear differential equation of order  $n$  can be articulated in the general form:

$$a_0(x)y^n + a_1(x)y^{n-1} + \dots + a_{n-1}(x)y' + a_n(x)y = g(x)$$

A homogeneous equation occurs when  $g(x) = 0$ . The equation is non-homogeneous when  $g(x) \neq 0$ . Comprehending the differentiation between these two types is essential, as they necessitate separate solution methodologies and produce varying solution frameworks. In mechanical engineering, homogeneous differential equations characterize undamped and damped oscillations in mechanical systems devoid of external influences. Examine a multi-mass spring system employed in the design of automobile suspension. The vertical displacement of each component can be represented by higher-order homogeneous differential equations, with the order contingent upon the quantity of masses in the system. Engineers evaluate these equations to enhance ride comfort, handling stability, and traction performance. The fundamental theorem for homogeneous linear differential equations asserts that if the coefficient functions  $a_i(x)$  are continuous over an interval  $I$  and  $a_0(x) \neq 0$  for every  $x$  in  $I$ , then there exist  $n$  linearly independent solutions  $y_1(x)$ ,  $y_2(x)$ , ...,  $y_n(x)$  inside that interval. The general solution is a linear amalgamation of these fundamental solutions:

$$y(x) = c_1y_1(x) + c_2y_2(x) + \dots + c_ny_n(x)$$

where  $c_1$ ,  $c_2$ , ...,  $c_n$  are arbitrary constants established by initial conditions.

The Wronskian determinant is employed to verify the linear independence of solutions. For  $n$  functions  $y_1(x)$ ,  $y_2(x)$ , ...,  $y_n(x)$ , the Wronskian is defined as follows:

$$W(y_1, y_2, \dots, y_n)(x) = \det([y_1(x), y_2(x), \dots, y_n(x); y_1'(x), y_2'(x), \dots, y_n'(x); \dots; y_1^{(n-1)}(x), y_2^{(n-1)}(x), \dots, y_n^{(n-1)}(x)])$$

The solutions constitute a fundamental set if and only if their Wronskian is non-zero at some point within the interval I.

In acoustical engineering, the Wronskian facilitates the analysis of sound wave propagation in intricate situations. For example, in the design of concert halls, engineers utilize differential equations to predict sound wave dynamics. By guaranteeing linearly independent answers via Wronskian analysis, they may precisely forecast sound quality at various sites and execute architectural modifications to enhance acoustic performance.

### **Homogeneous Equations with Constant Coefficients: The Characteristic Equation Method**

For linear homogeneous differential equations characterized by constant coefficients:

$$a_0y^n + a_1y^{(n-1)} + \dots + a_{n-1}y' + a_ny = 0$$

The solution method use the characteristic equation:

$$a_0r^n + a_1r^{(n-1)} + \dots + a_{n-1}r + a_n = 0$$

The roots of this polynomial problem dictate the structure of the solution. Three scenarios must be examined:

1. Distinct real roots: If  $r_1, r_2, \dots, r_n$  are distinct real roots, the general solution is expressed as:  $y(x) = c_1e^{(r_1x)} + c_2e^{(r_2x)} + \dots + c_ne^{(r_nx)}$
2. Repeated real roots: If  $r_1$  has a multiplicity of  $k$ , the associated terms in the solution are:  $c_1e^{(r_1x)} + c_2xe^{(r_1x)} + c_3x^2e^{(r_1x)} + \dots + c_kx^{(k-1)}e^{(r_1x)}$
3. Complex conjugate roots: If  $a+bi$  and  $a-bi$  are roots, the associated terms in the solution are:  $e^{(ax)}(c_1\cos(bx) + c_2\sin(bx))$ .

This characteristic equation method is essential in electronic circuit design. Examine a series RLC circuit comprising a resistor, inductor, and capacitor. The present flow is regulated by a second-order differential equation. The circuit may demonstrate overdamped (distinct real roots), critically damped (repeated real roots), or underdamped (complex conjugate roots) behavior, contingent upon the component values. Engineers evaluate these scenarios to build circuits with specified transient responses for applications including power supplies and communication systems.

The aerospace sector utilizes higher-order differential equations with constant coefficients to simulate aircraft stability. The dynamics of longitudinal and lateral motion are generally expressed by fourth-order equations. The roots of the characteristic equation are directly related to flight stability characteristics. Real negative roots signify stable damping modes, but complex roots with positive real components suggest perilous instabilities that may result in catastrophic failures. Flight control systems are engineered to manipulate these roots to guarantee steady flight under diverse operational situations.

### **Initial Value Problems for Higher-Order Differential Equations**

Differential equations never exist independently in practical applications; they are typically accompanied with initial conditions that define the system's state at a specific moment. An  $n$ th-order equation necessitates  $n$  initial conditions to uniquely ascertain the solution. These generally assume the following format:

$$y(x_0) = y_0, y'(x_0) = y_1, \dots, y^{(n-1)}(x_0) = y_{n-1}$$

Upon deriving the general solution, the initial conditions are employed to ascertain the exact values of the arbitrary constants  $c_1, c_2, \dots, c_n$ .

In biomedical engineering, starting value problems are crucial for estimating drug concentration in multi-compartment pharmacokinetic models. The distribution of a medicine throughout different body tissues upon administration can be described using higher-order differential equations. Initial circumstances denote the initial concentration within each compartment. Healthcare practitioners utilize these models to construct appropriate dosing schedules, guaranteeing therapeutic drug concentrations while reducing adverse effects. Robotics engineers have analogous difficulties while programming the movements of robotic arms. The behavior of a multi-jointed robotic arm can be characterized by a set of higher-order differential equations. The initial circumstances delineate the initial location, velocity, and acceleration of each joint. Engineers create control algorithms by resolving these initial value difficulties, allowing robots to execute precise motions in production, surgery, and exploratory contexts.

The existence and uniqueness theorem for initial value problems guarantees that, under specific circumstances (continuous coefficients and right-hand side), a unique solution is present in a vicinity of the beginning point. This theorem supports the dependability of computational techniques employed in simulation software for engineering purposes.

### **Non-Homogeneous Linear Differential Equations**

When  $g(x) \neq 0$  in the original equation, it constitutes a non-homogeneous equation. The comprehensive solution to such an equation comprises two components:

$$y(x) = y_h(x) + y_p(x)$$

where  $y_h(x)$  represents the general solution to the associated homogeneous equation (complementary solution), and  $y_p(x)$  denotes any particular solution to the non-homogeneous equation.

In environmental engineering, non-homogeneous differential equations represent pollutant dispersal in watersheds. The homogeneous component delineates the natural dispersion and degradation of the pollutant, whereas the specific solution illustrates the impact of ongoing pollution sources. Through the analysis of both components, environmental experts formulate remediation techniques and determine safe discharge limits for industrial facilities.

A variety of techniques are available for identifying specific solutions, including:

1. Method of Undetermined Coefficients
2. Variation of parameters
3. The annihilation technique

The method of unknown coefficients is suitable when  $g(x)$  is a well-behaved function, often a polynomial, exponential, sine, cosine, or a combination thereof. The method entails formulating an informed hypothesis regarding the structure of the particular solution derived from  $g(x)$ , substituting this into the original equation, and resolving for the unknown coefficients. This method assists engineers in evaluating building reactions to harmonic loads from machinery in structural dynamics. The forcing function  $g(x)$  denotes the periodic force, whereas the particular solution illustrates the

steady-state vibrational response. Engineers utilize this information to devise vibration isolation devices that avert machinery-induced resonance in structural edifices.

### **The Annihilator Method: A Refined Technique for Non-Homogeneous Equations**

The annihilator method offers an alternate technique for determining individual solutions to non-homogeneous equations. The essential idea is to convert the non-homogeneous equation into a higher-order homogeneous equation by employing a suitable differential operator that eliminates the non-homogeneous term  $g(x)$ .

For instance, if  $g(x) = e^{(\alpha x)}$ , then the operator  $(D-\alpha)$ , where  $D = d/dx$ , annihilates  $g(x)$  since  $(D-\alpha)e^{(\alpha x)} = 0$ . Applying this operator to both sides of the original equation yields a homogeneous equation of superior order. Upon resolution, we derive the specific solution by isolating elements absent in the complimentary solution.

The annihilator approach in quantum mechanics is effective for solving time-dependent Schrödinger equations with certain potential functions. Quantum scientists employ this technique to examine particle behavior in dynamic fields, facilitating the advancement of quantum computing components and precision measurement instruments. The annihilator method is especially refined when addressing combinations of functions. If  $g(x) = g_1(x) + g_2(x)$ , and  $L_1$  and  $L_2$  are operators that annihilate  $g_1(x)$  and  $g_2(x)$  respectively, then the operator  $L_1L_2$  annihilates the entire function  $g(x)$ , provided that  $L_1$  and  $L_2$  commute, which is the case for constant coefficient operators. Financial analysts utilize the annihilator method to describe intricate economic systems with various driving functions. A nation's inflation rate may be affected by several cyclical causes (seasonal expenditure patterns) and exponential trends (monetary policy impacts). Through the application of suitable annihilator operators, economists construct intricate models that assist central banks in devising effective monetary policies.

## The Algebra of Differential Operators with Constant Coefficients

The examination of differential equations with constant coefficients inherently results in an algebraic framework for the collection of differential operators. Let  $D$  represent the operator  $d/dx$ . Any linear differential operator with constant coefficients can be expressed as a polynomial in  $D$ .

$$L = a_0 D^n + a_1 D^{(n-1)} + \dots + a_{n-1} D + a_n$$

These operators constitute an algebra characterized by the following properties:

1. Summation:  $(L_1 + L_2)y = L_1 y + L_2 y$
2. Scalar multiplication:  $(cL)y = c(Ly)$
3. Multiplication (composition):  $(L_1 L_2)y = L_1(L_2 y)$

The multiplication of these operators is commutative, a property not typically applicable to differential operators with variable coefficients. This commutativity enables the factoring of operators akin to polynomials, significantly streamlining solution techniques. This algebraic method aids in the design of intricate feedback controllers in control systems engineering. Engineers can algebraically alter formulas expressing both plant dynamics and the controller as differential operators to attain the required closed-loop behavior. This technique is essential for creating control systems in applications from driverless vehicles to industrial process control.

The factorization of differential operators is closely connected to the characteristic equation. If  $L = a_0 D^n + a_1 D^{(n-1)} + \dots$  and  $r_1, r_2, \dots, r_n$  denote the roots of the characteristic equation, then:

$$L = a_0 (D - r_1)(D - r_2) \dots (D - r_n)$$

This factored form elucidates the structure of solutions and facilitates the implementation of the annihilator approach.

In telecommunications, engineers employ operator factorization to create filters with defined frequency response attributes. The factored form illustrates the filter's impact on various frequency components, facilitating the development of accurate bandpass, notch, and equalizing filters vital for contemporary communication systems.

## **Pragmatic Implementations across Disciplines**

The aforementioned theoretical approach is applicable across various domains, tackling intricate real-world issues:

### **Mechanical and Structural Engineering**

In contemporary skyscraper architecture, wind-induced oscillation is a significant issue. The building's reaction to wind forces can be represented using non-homogeneous differential equations, with the wind force denoted by the  $g(x)$  term. The complementary solution delineates the building's inherent vibration modes, whereas the particular solution encapsulates the induced response to wind loads. Engineers evaluate these equations to deploy dampening systems—such as tuned mass dampers—that alleviate excessive oscillation during strong winds. Automotive engineers utilize higher-order differential equations in active suspension systems. In contrast to passive suspensions that solely utilize springs and dampers, active systems incorporate sensors, actuators, and controls to dynamically modify damping properties. The system's behavior is represented by non-homogeneous equations, with road irregularities acting as the forcing function. The vehicle's onboard computer can alter suspension characteristics in real-time by solving these equations, thereby optimizing comfort and handling for diverse road conditions.

### **Electrical Engineering and Signal Processing**

Contemporary digital filters apply methods to solve constant coefficient differential equations. In constructing filters for applications such as noise reduction in audio recordings or feature extraction in medical data, engineers initially determine the required frequency response. This is converted into a differential equation, thereafter solved and discretized for digital implementation. The annihilator method is very effective in the design of notch filters aimed at removing certain frequency components, such as 60Hz power line interference in biomedical signals.

In power grid management, the stability of interconnected generators is assessed by higher-order differential equations. The dynamics of each generator contribute to the overall system's order, leading to high-dimensional models. Engineers utilize the principles of linear differential equations to evaluate grid stability amongst many disturbance scenarios and



to devise protection measures that avert cascading failures resulting in extensive blackouts.

### **Biomedical Engineering and Physiological Simulation**

The glucose-insulin regulation systems in diabetes individuals are represented by higher-order differential equations. These models consider glucose absorption from diet, insulin secretion or administration, and glucose use by tissues. Medical researchers resolve these equations to create artificial pancreas devices that autonomously regulate insulin supply based on continuous glucose monitoring, thereby enhancing the quality of life for diabetic patients. Electroencephalography (EEG) records of brain activity can be evaluated employing differential equations via the annihilator method. Neurologists discern distinctive patterns linked to epileptic seizures by representing these signals as solutions to particular differential equations. This mathematical methodology facilitates the creation of early warning systems for seizure prediction and intervention.

### **Environmental Science and Climate Modeling**

Climate scientists utilize higher-order differential equations to model the dynamics of the carbon cycle. These equations delineate carbon exchange among the atmosphere, oceans, and terrestrial ecosystems. The non-homogeneous terms signify anthropogenic carbon emissions. Through the resolution of these equations across diverse emission scenarios, scientists forecast future atmospheric CO<sub>2</sub> levels and corresponding temperature variations, thereby guiding worldwide climate policy decisions. Water quality in river systems is represented by differential equations that incorporate pollution movement, dilution, and degradation mechanisms. Environmental engineers utilize the annihilator approach to assess the cumulative impacts of various pollution sources along a river. This mathematical methodology informs the formulation of watershed management policies that uphold water quality criteria while reconciling economic development requirements.

### **Economics and Finance**

In macroeconomic modeling, business cycles are depicted by higher-order differential equations. The interplay among variables such as GDP, inflation, unemployment, and interest rates generates intricate dynamics that can be

examined through the previously outlined mathematical framework. Policymakers resolve these equations to predict economic outcomes under various fiscal and monetary interventions, maximizing policy responses to economic recessions. Option pricing in financial markets entails resolving differential equations originating from stochastic processes. The Black-Scholes equation, essential to contemporary finance, is a second-order partial differential equation. Financial analysts ascertain fair pricing for intricate derivative products by implementing suitable transformations and boundary conditions, akin to starting conditions, hence facilitating effective risk management techniques for institutional investors.

### **Computational Techniques and Numerical Resolutions**

Although analytical methods yield significant insights, numerous practical applications necessitate numerical solutions owing to system complexity or non-linearities. Contemporary computational methodologies encompass:

1. Runge-Kutta techniques
2. Finite difference methodologies
3. Spectral techniques
4. Shooting methodologies for boundary value issues

These numerical methods apply the previously described theoretical ideas, broadening their use to scenarios where closed-form solutions are unavailable.

In aerospace engineering, flight simulators resolve intricate differential equations in real-time to precisely simulate aircraft dynamics. The equations encompass aerodynamic forces, engine performance, and control surface influences. Numerical integration techniques derived from initial value problem theory allow pilots to practice in virtual settings that accurately simulate aircraft reactions to control inputs across various flight conditions. Weather forecasting depends on extensive numerical simulations of differential equations that characterize atmospheric physics. These equations represent the dynamics of air movement, heat transport, moisture, and radiation processes. Notwithstanding its intricacy, the fundamental mathematical framework adheres to the ideas established for linear differential equations. Meteorologists utilize advanced numerical techniques

on these equations to produce forecasts that assist communities in preparing for extreme weather occurrences.

### **Novel Applications in Data Science and Machine Learning**

Recent advancements in machine learning have generated novel applications for the theory of differential equations. Neural ordinary differential equations (Neural ODEs) characterize the dynamics of neural networks as continuous-time models regulated by differential equations. The network parameters delineate the vector field of the ODE, and training entails improving these parameters to align with observed data paths. This method provides benefits in modeling time-series data characterized by unpredictable sample intervals, a prevalent issue in health monitoring and financial markets. Data scientists utilize the comprehensive theory of differential equations to create more interpretable machine learning models with enhanced generalization capabilities. In reinforcement learning, optimal control policies for robotics and autonomous systems are obtained from solutions to differential equations referred to as Hamilton-Jacobi-Bellman equations. These higher-order equations delineate the gradient of the value function throughout the state space. Engineers utilize numerical methods derived from the theory of initial value problems to resolve these equations, facilitating optimal decision-making by robots in intricate, dynamic settings.

### **Obstacles and Prospective Pathways**

Notwithstanding considerable progress, some obstacles persist in the theory and implementation of higher-order differential equations:

1. **Stiffness:** Systems exhibiting significant disparities in time scales result in numerical instability when employing conventional methodologies. Specialized implicit schemes are necessary but elevate computational expenses.
2. **High dimensionality:** Real-world systems frequently encompass multiple interrelated equations, rendering analytical methods impractical and numerical solutions computationally demanding.
3. **Parameter uncertainty:** In actual applications, coefficient values may be imprecise, requiring sensitivity analysis and rigorous solution methodologies.

4. Non-linearity: Numerous practical systems demonstrate non-linear behavior, necessitating linearization techniques or specific non-linear solution approaches.

Prospective avenues for research encompass:

1. Enhancing the efficacy of numerical techniques for high-dimensional systems
2. Incorporating uncertainty quantification into solution methodologies
3. Utilizing machine learning methodologies to estimate solutions for intricate differential equations
4. Investigating the convergence of differential equations and data-driven modeling

The theory of homogeneous and non-homogeneous linear differential equations of order  $n$  offers a robust framework for modeling and evaluating dynamic systems in several domains. Mathematical tools facilitate engineers, scientists, and analysts in describing, predicting, and controlling complicated events, spanning from classical mechanics to advanced artificial intelligence. The sophisticated interaction between differential operators and their algebraic characteristics, especially via the annihilator approach, provides both theoretical understanding and practical solution strategies. Initial value problems link abstract mathematical constructs to tangible physical conditions, facilitating accurate modeling of real-world systems. With the ongoing advancement of computational powers, the range and accuracy of differential equation models will broaden, extending the limits of what is achievable in science and engineering. The core notions delineated in this examination will persist as pivotal to these advancements, underscoring the lasting significance of mathematical theory in confronting humanity's most urgent issues. By learning these principles, contemporary practitioners acquire a mathematical toolkit adept at addressing challenges of unparalleled complexity and significance—ranging from climate forecasting to autonomous systems, from pandemic modeling to space exploration. The theory of differential equations is one of humanity's most important intellectual accomplishments, consistently broadening its influence across various fields of human activity.

## SELF ASSESSMENT QUESTIONS

### Multiple Choice Questions (MCQs)

1. The characteristic equation of an  $n^{\text{th}}$  order linear differential equation with constant coefficients is obtained by:
  - a) Substituting  $y=erx$  into the differential equation
  - b) Integrating the equation
  - c) Differentiating the equation
  - d) None of the above
2. If the characteristic equation has distinct real roots, the general solution is given by:
  - a) A sum of exponential functions
  - b) A sum of polynomial terms
  - c) A sum of sine and cosine functions
  - d) None of the above
3. The annihilator method is used to:
  - a) Solve homogeneous equations
  - b) Solve non-homogeneous equations
  - c) Find the Wronskian
  - d) None of the above
4. The method of undetermined coefficients is applicable when the non-homogeneous term is:
  - a) A polynomial, exponential, or trigonometric function
  - b) An arbitrary function
  - c) A discontinuous function
  - d) None of the above
5. The fundamental set of solutions of an  $n$ th order differential equation must consist of:
  - a)  $n$  linearly independent solutions
  - b)  $n-1$  solutions
  - c) Only one solution
  - d) None of the above
6. The operator equation  $(D-2)(D+3)y=0$  has a general solution of the form:
  - a)  $y=C_1e^{2x}+C_2e^{-3x}$

b)  $y=C_1e^{-2x}+C_2e^{3x}$

c)  $y=C_1e^x+C_2e^{-x}$

d) None of the above

7. The roots of the characteristic equation determine:

a) The form of the solution

b) The initial conditions

c) The uniqueness of the solution

d) None of the above

8. If a root of the characteristic equation is complex, the corresponding solution involves:

a) Exponential and trigonometric terms

b) Polynomials only

c) Logarithmic functions

d) None of the above

Answer Key:

1	a	3	b	5	a	7	a
2	a	4	a	6	a	8	a

## CHECK YOUR PROGRESS

- 1) Write two algebraic properties of the D-operator.

.....  
.....  
.....

- 2) Explain how inverse operators are used in solving differential equations.

.....  
.....  
.....

## LET US SUM UP

The differential operator  $D$ , defined as  $D = d/dx$ , obeys the rules of algebra. Operators can be added, multiplied, or factored like algebraic quantities. For example,  $D^m D^n = D^{m+n}$  and  $(D-a)(D-b) = D^2 - (a+b)D + ab$ . Using inverse operators such as  $1/(D-a)$ , we can express and compute particular integrals of non-homogeneous equations. This operator approach simplifies the process of solving linear equations with constant coefficients.

## UNIT END EXERCISES

### Short Answer Questions:

1. What type of functions can be handled using the method of undetermined coefficients?
2. Solve the characteristic equation  $r^3 - 3r^2 + 2r = 0$ .
3. What is the role of initial conditions in solving higher-order differential equations?

### Long Answer Questions:

1. Solve the equation  $y''' - y' = x^2$  using the method of undetermined coefficients.
2. How do we solve an equation with complex characteristic roots? Provide an example.
3. Discuss real-world applications of higher-order linear differential equations.

### REFERENCES AND SUGGESTED READINGS:

1. Kreyszig, E. (2019). Advanced Engineering Mathematics (10th ed.). Wiley.
2. O'Neil, P. V. (2017). Advanced Engineering Mathematics (8th ed.). Cengage Learning.
3. Greenberg, M. D. (2014). Ordinary Differential Equations (2nd ed.). Wiley.
4. Rainville, E. D., & Bedient, P. E. (2013). Elementary Differential Equations (8th ed.). Pearson.



## BLOCK 3

### UNIT 7

#### LINEAR EQUATIONS WITH VARIABLE COEFFICIENTS: INITIAL VALUE PROBLEMS FOR THE HOMOGENEOUS EQUATION

##### Objectives

- Understand and solve initial value problems for linear equations with variable coefficients.
- Study the solutions of homogeneous linear equations with variable coefficients.
- Explore the Wronskian and its role in determining linear independence.
- Learn the reduction of order method for solving second-order equations.
- Examine homogeneous equations with analytic coefficients.
- Understand and solve the Legendre equation.

##### 7.1 Introduction to Linear Equations with Variable Coefficients

Linear equations with variable coefficients represent a fascinating and fundamental area of mathematical study that bridges algebraic manipulation, mathematical reasoning, and practical problem-solving. These equations are characterized by their linear structure, where variables are raised to the first power and can have coefficients that themselves change or depend on other variables.

##### Fundamental Concepts and Definitions

A linear equation with variable coefficients can be generally expressed in the form:

$$a(x)y + b(x)y' + c(x)y = f(x)$$

Where:

- $y$  is the dependent variable
- $x$  is the independent variable

- $a(x)$ ,  $b(x)$ , and  $c(x)$  are functions of  $x$  that serve as coefficients
- $y'$  represents the first derivative of  $y$  with respect to  $x$
- $f(x)$  is a known function representing the right-hand side of the equation

### Key Characteristics

1. Linearity: The equation remains linear in the dependent variable ( $y$ ) and its derivatives.
2. Variable Coefficients: The coefficients are functions of the independent variable, not constant values.
3. Complexity: These equations are more sophisticated than standard linear equations with constant coefficients.

### Mathematical Framework

#### Classification of Linear Equations with Variable Coefficients

1. First-Order Linear Differential Equations
2. Second-Order Linear Differential Equations
3. Higher-Order Linear Differential Equations

### Solved Problems

#### Problem 1: Basic Variable Coefficient Linear Equation

Problem Statement: Solve the differential equation:  $y' + p(x)y = q(x)$ , where  $p(x)$  and  $q(x)$  are continuous functions.

Solution Steps:

1. Multiply both sides by the integrating factor  $e^{\int p(x)dx}$
2. Rearrange to obtain the general solution
3. Apply initial conditions if provided

Detailed Solution: Consider  $p(x) = 1/x$  and  $q(x) = x$  for  $x > 0$

Integrating factor:  $\exp(\int (1/x)dx) = \exp(\ln(x)) = x$

Multiply the original equation by  $x$ :  $x(y' + (1/x)y) = xy$

Rearranging:  $xy' + y = xy$

Integrate both sides:  $\int(xy')dx + \int y dx = \int(xy)dx$

Result:  $y = C/x + x$

Where C is an arbitrary constant determined by initial conditions.

### **Problem 2: Second-Order Variable Coefficient Equation**

Problem Statement: Solve the equation:  $x^2y'' + xy' - y = 0$ , valid for  $x > 0$

#### **Solution Methodology:**

1. Recognize this as a Cauchy-Euler equation
2. Assume solution of the form  $y = x^r$
3. Substitute and solve the characteristic equation
4. Determine general solution

Detailed Solution: Substituting  $y = x^r$ :  $x^2(r(r-1)x^{(r-2)}) + x(rx^{(r-1)}) - x^r = 0$

Simplifying:  $r(r-1) + r - 1 = 0$   $r^2 = 1$

Roots:  $r_1 = 1, r_2 = -1$

General solution:  $y = C_1x + C_2 1/x$

### **Problem 3: First-Order Nonhomogeneous Equation**

Problem Statement: Solve  $y' + (2/x)y = x^2$ , for  $x > 0$

#### **Solution Steps:**

1. Identify integrating factor
2. Multiply equation
3. Integrate to find general solution

Detailed Solution: Integrating factor:  $\exp(\int(2/x)dx) = x^2$

Multiplying equation by  $x^2$ :  $x^2y' + 2xy = x^4$

Integrating:  $x^2y = (x^4/2) + C$

Final solution:  $y = (x^2/2) + (C/x^2)$

#### **Problem 4: Legendre's Equation**

Problem Statement: Solve  $(1-x^2)y'' - 2xy' + n(n+1)y = 0$

Solution Approach:

1. Power series method
2. Frobenius method
3. Determine series solution

Detailed Solution: Assume solution:  $y = \sum_{k=0}^{\infty} a_k x^k$

Substitution leads to recurrence relations for coefficients, resulting in Legendre polynomials.

#### **Problem 5: Bessel's Equation**

Problem Statement: Solve  $x^2 y'' + xy' + (x^2 - n^2)y = 0$

Solution Methodology:

1. Power series solution
2. Frobenius method
3. Derive Bessel functions

Detailed Solution: Series solution converges to Bessel functions of the first and second kind.

#### **Unsolved Problems**

##### **Problem 1: Advanced Variable Coefficient Equation**

Prove existence and uniqueness of solutions for the equation:  $y'' + p(x)y' + q(x)y = f(x)$  Where  $p(x)$  and  $q(x)$  have specific continuity constraints.

##### **Problem 2: Singular Point Analysis**

Characterize singular points for the differential equation:  $x^2 y'' + axy' + by = 0$   
Determine conditions for regular and irregular singularities.

##### **Problem 3: Asymptotic Behavior**

Investigate asymptotic properties of solutions to:  $y'' + (1/x)y' + (\sin(x)/x^2)y = 0$   
As  $x$  approaches infinity.

#### **Problem4: Transformation Methods**

Develop a general transformation method to convert variable coefficient equations to constant coefficient forms.

#### **Problem 5: Numerical Stability**

Design a numerical method with guaranteed stability for solving high-order linear equations with rapidly changing coefficients.

#### **Theoretical Foundations**

##### **Existence and Uniqueness Theorems**

1. Picard-Lindelöf Theorem: Guarantees existence and uniqueness of solutions under certain continuity conditions.
2. Cauchy-Peano Theorem: Provides conditions for local existence of solutions.

##### **Computational Approaches**

1. Numerical Methods
  - Runge-Kutta methods
  - Predictor-corrector algorithms
  - Shooting methods
2. Symbolic Computation
  - Computer algebra systems
  - Symbolic manipulation techniques

Linear equations with variable coefficients represent a rich and complex domain of mathematical investigation. They bridge theoretical mathematics with practical applications in physics, engineering, and applied sciences. The exploration of these equations reveals intricate relationships between mathematical structures, computational methods, and fundamental principles of dynamic systems. Continued research in this area promises deeper insights into mathematical modeling, numerical analysis, and theoretical foundations of differential equations.

## 7.2 Initial Value Problems for Homogeneous Equations

### Theoretical Foundation

Initial value problems (IVPs) are fundamental in differential equations, representing mathematical models where we seek a solution to a differential equation that satisfies specific initial conditions. For homogeneous linear differential equations, these problems involve finding a solution that passes through predetermined points or satisfies specific constraints at the initial time.

### Basic Concept of Initial Value Problems

An initial value problem for a first-order linear homogeneous differential equation can be generally expressed as:

$$dy/dx + P(x)y = Q(x)$$

Where:

- $y$  is the dependent variable
- $x$  is the independent variable
- $P(x)$  and  $Q(x)$  are continuous functions in a given interval

### Key Components

1. Differential Equation: The mathematical relationship describing the rate of change
2. Initial Condition: Specific value of the solution at a starting point
3. Solution Domain: The interval where the solution is defined and continuous

### Solving Initial Value Problems: Methodological Approach

#### Step-by-Step Solution Strategy

1. Identify the type of differential equation
2. Determine the appropriate solution method
3. Apply initial conditions
4. Verify the solution's validity

## Solved Problems

### Problem 1: Standard Linear Homogeneous IVP

Problem Statement: Solve the differential equation  $dy/dx + 2y = 0$ , with the initial condition  $y(0) = 5$

Solution Process:

1. Recognize this as a first-order linear homogeneous equation
2. Separate variables:  $dy/y = -2dx$
3. Integrate both sides:  $\ln|y| = -2x + C$
4. Exponentiate:  $y = e^{(-2x + C)}$
5. Apply initial condition:  $5 = e^C$
6. Final solution:  $y = 5e^{(-2x)}$

Verification:

- Substituting back into original equation:  $dy/dx + 2y = -10e^{(-2x)} + 2(5e^{(-2x)}) = 0$  ✓
- Initial condition:  $y(0) = 5e^{(0)} = 5$  ✓

### Problem 2: Variable Coefficient Homogeneous IVP

Problem Statement: Solve  $dy/dx + xy = x$ , with  $y(0) = 2$

Solution Process:

1. Identify as a first-order linear non-homogeneous equation
2. Use integrating factor method
3. Integrating factor:  $\mu(x) = \exp(\int x \, dx) = \exp(x^2/2)$
4. Multiply equation by integrating factor
5. Integrate and solve
6. Final solution:  $y = 2e^{(-x^2/2)} + 1 - x^2/2$

Verification Steps:

- Check derivative conditions

- Validate initial condition
- Substitute back into original equation

### **Problem 3: Second-Order Homogeneous Linear IVP**

Problem Statement: Solve  $d^2y/dx^2 + 4y = 0$ , with  $y(0) = 3$  and  $dy/dx(0) = 1$

Solution Process:

1. Characteristic equation:  $r^2 + 4 = 0$
2. Roots:  $r = \pm 2i$
3. General solution:  $y = C_1 \cos(2x) + C_2 \sin(2x)$
4. Apply initial conditions:
  - $y(0) = 3$  implies  $C_1 = 3$
  - $dy/dx(0) = 1$  implies  $C_2 = 1/2$
5. Final solution:  $y = 3 \cos(2x) + (1/2)\sin(2x)$

### **Problem 4: Exponential Coefficient IVP**

Problem Statement: Solve  $dy/dx + e^x y = x$ , with  $y(0) = 1$

Solution Process:

1. Use variation of parameters
2. Construct fundamental solution
3. Apply integration techniques
4. Final solution:  $y = e^{(-ex)}(1 + \int x e^{(ex)} dx)$

### **Problem 5: Coupled Initial Value Problem**

Problem Statement: Solve the system:  $dy/dx = y + 2z$   $dz/dx = 3y - z$  Initial conditions:  $y(0) = 1, z(0) = 2$

Solution Process:

1. Use matrix exponential method
2. Construct state transition matrix
3. Apply initial condition vector



4. Derive complete solution

### **Unsolved Problems**

#### **Unsolved Problem 1: Advanced Nonlinear IVP**

Develop a solution method for:  $dy/dx = y^2 + \sin(x)$ ,  $y(0) = 1$

#### **Unsolved Problem 2: Fractional Order Differential Equation**

Investigate the solution of:  $D^{(0.5)}y + y = x$ , where  $D^{(0.5)}$  represents fractional derivative

#### **Unsolved Problem 3: Singular Point Analysis**

Analyze the behavior of solutions near singular points in the equation:  $x^2 ((dy)/[dx]) + x(dy/dx) - y = 0$

#### **Unsolved Problem 4: Stochastic Initial Value Problem**

Develop a probabilistic approach to solving:  $dy = (y + \text{noise})dx$ , with  $y(0) = a$

#### **Unsolved Problem 5: Multi-Point Boundary Conditions**

Explore solution techniques for:  $y''(x) + p(x)y'(x) + q(x)y(x) = f(x)$ , with mixed boundary conditions

## CHECK YOUR PROGRESS

- 1) Write the general form of a second-order homogeneous equation with variable coefficients.

.....  
.....  
.....

- 2) Explain why variable coefficients make solving differential equations more complex.

.....  
.....  
.....

- 3) What is the role of initial conditions in these problems?

.....  
.....  
.....

## LET US SUM UP

A second-order differential equation with variable coefficients has the general form  $y'' + P(x)y' + Q(x)y = 0$ , where  $P(x)$  and  $Q(x)$  vary with  $x$ . Such equations cannot generally be solved using simple exponential solutions. Instead, special techniques like the method of reduction of order or power series solutions are used. When initial conditions are provided, they help to determine the particular solution that fits those specific values.

## UNIT END EXERCISES

### Short Answer Questions:

1. Define a second-order linear equation with variable coefficients.
2. What is the Wronskian, and how is it used to determine linear independence?

### Long Answer Questions:

1. Derive and solve the Legendre equation for  $P_n(x)$ .
2. Explain the reduction of order method and solve  $y'' - 2y' + y = 0$  given that  $y_1 = e^x$ .

### REFERENCES AND SUGGESTED READINGS:

1. Coddington, E. A., & Levinson, N. (2021). Theory of Ordinary Differential Equations. McGraw-Hill.
2. Ince, E. L. (2012). Ordinary Differential Equations. Dover Publications.
3. Birkhoff, G., & Rota, G. C. (2010). Ordinary Differential Equations (4th ed.). Wiley.

## UNIT 8

### SOLUTIONS OF THE HOMOGENEOUS EQUATION

#### 8.1 Solutions of Homogeneous Equations with Variable Coefficients

##### Theoretical Overview

Homogeneous linear differential equations with variable coefficients represent a complex class of mathematical models encountered in various scientific disciplines, including physics, engineering, and applied mathematics.

##### Key Characteristics

1. Coefficients are functions of the independent variable
2. Solution methods are more intricate compared to constant coefficient equations
3. Require advanced mathematical techniques

##### Solution Techniques

###### 1. Power Series Method

- Assumes solution in the form of a power series
- Determines coefficients through recursive relationships
- Particularly useful near ordinary points

###### 2. Frobenius Method

- Extends power series approach
- Handles regular singular points
- Provides more robust solution techniques

###### 3. Asymptotic Expansion

- Approximates solutions for large or small independent variable values
- Useful in limit behavior analysis

##### Mathematical Framework

For a general linear homogeneous differential equation:

$$a_n(x)y^{(n)} + a_{(n-1)}(x)y^{(n-1)} + \dots + a_1(x)y' + a_0(x)y = 0$$

Where:

- $a_i(x)$  are continuous functions
- $y^{(k)}$  represents k-th derivative of y

Understanding initial value problems and solutions for homogeneous equations with variable coefficients requires advanced mathematical techniques, combining algebraic manipulation, series expansions, and deep analytical insights. The exploration of these mathematical models continues to be a rich area of research, offering profound insights into complex dynamic systems across scientific disciplines.

## CHECK YOUR PROGRESS

- 1) Define the reduction of order method.

.....  
.....  
.....

- 2) Given  $y_1 = e^x$ , find  $y_2$  for  $y'' - y' = 0$ .

.....  
.....  
.....

- 3) Write the formula for finding the second solution when one solution is known.

.....  
.....  
.....

- 4) Explain why two independent solutions are necessary.

.....  
.....  
.....

## LET US SUM UP

The solution of a homogeneous equation with variable coefficients can often be obtained using the method of reduction of order, particularly when one solution is already known. If  $y_1(x)$  is a known solution, then the second solution can be found using the formula  $y_2 = y_1 \int \frac{e^{-\int P(x)dx}}{y_1^2} dx$ . This ensures that  $y_1$  and  $y_2$  are linearly independent, forming the general solution of the differential equation.

## UNIT END EXERCISES

### Short Answer Questions:

1. Explain the reduction of order method with an example.
2. What are analytic coefficients, and why are they important?
3. Describe the Legendre equation and its significance.

### Long Answer Questions:

1. Discuss the role of the Wronskian in differential equations with variable coefficients.
2. Derive the variation of parameters formula and use it to solve  $y'' + p(x)y' + q(x)y = f(x)$ .

### REFERENCES AND SUGGESTED READINGS:

1. Coddington, E. A., & Levinson, N. (2021). Theory of Ordinary Differential Equations. McGraw-Hill.
2. Nagle, R. K., Saff, E. B., & Snider, A. D. (2018). Fundamentals of Differential Equations (9th ed.). Pearson.
3. Birkhoff, G., & Rota, G. C. (2010). Ordinary Differential Equations (4th ed.). Wiley.

## UNIT 9

### THE WRONSKIAN AND LINEAR INDEPENDENCE – REDUCTION OF THE ORDER OF A HOMOGENEOUS EQUATION

#### 9.1 The Wronskian and Linear Independence

The Wronskian is a powerful mathematical tool used in linear algebra and differential equations to determine the linear independence of a set of functions. Named after Józef Hoene-Wroński, a Polish mathematician and philosopher, this determinant-based method provides crucial insights into the relationship between different functions.

##### Fundamental Definition

For a set of  $n$  differentiable functions  $f_1(x)$ ,  $f_2(x)$ , ...,  $f_n(x)$ , the Wronskian  $W(x)$  is defined as the determinant of a matrix constructed from these functions and their successive derivatives:

$$W(x) = \det \begin{vmatrix} f_1(x) & f_2(x) & \dots & f_n(x) \\ f_1'(x) & f_2'(x) & \dots & f_n'(x) \\ f_1''(x) & f_2''(x) & \dots & f_n''(x) \\ \vdots & \vdots & \ddots & \vdots \\ f_1^{(n-1)}(x) & f_2^{(n-1)}(x) & \dots & f_n^{(n-1)}(x) \end{vmatrix}$$

##### Key Theoretical Insights

##### 1. Linear Independence Criterion

- If the Wronskian is non-zero at any point in an interval, the functions are linearly independent on that interval.
- If the Wronskian is zero at every point in an interval, the functions are linearly dependent.

##### 2. Differential Equation Connection

The Wronskian plays a critical role in solving linear differential equations, particularly in determining the general solution and understanding the relationship between solution functions.

##### Theoretical Foundation

##### Mathematical Formulation



Consider a system of  $n$  differential functions  $f_1(x)$ ,  $f_2(x)$ , ...,  $f_n(x)$ . The Wronskian provides a systematic method to assess their linear relationships through derivative analysis.

### **Computational Approach**

The Wronskian can be calculated through several methods:

1. Direct determinant computation
2. Recursive derivation
3. Symbolic manipulation

### **Properties of the Wronskian**

1. Symmetry and Antisymmetry
  - The Wronskian has specific symmetry properties based on function characteristics
  - Changes in function order can modify determinant sign
2. Derivative Relationship: The Wronskian satisfies a remarkable differential equation relationship, revealing deep connections between function derivatives.

### **Computational Methodology**

#### **Calculation Techniques**

1. Direct Matrix Determinant
  - Construct the matrix of functions and derivatives
  - Compute the determinant using standard linear algebra techniques
2. Recursive Computation
  - Develop algorithms for systematic Wronskian evaluation
  - Implement computational strategies for complex function sets

### **Algorithmic Representation**

Function `ComputeWronskian(functions[], interval):`

Initialize matrix M

For each function in functions:

Compute derivatives

Populate matrix rows

Compute determinant of matrix M

Return determinant value

## Solved Problems

### Problem 1: Basic Wronskian Calculation

Problem: Determine the Wronskian for functions  $f_1(x) = x$ ,  $f_2(x) = x^2$

Solution:

1. First function:  $f_1(x) = x$
2. First derivative:  $f_1'(x) = 1$
3. Second function:  $f_2(x) = x^2$
4. First derivative:  $f_2'(x) = 2x$

$$\text{Wronskian} = \det \begin{vmatrix} x & x^2 \\ 1 & 2x \end{vmatrix}$$

$$W(x) = x(2x) - x^2(1) = 2x^2 - x^2 = x^2$$

The Wronskian is non-zero for  $x \neq 0$ , indicating linear independence.

### Problem 2: Trigonometric Function Wronskian

Problem: Calculate the Wronskian for  $\sin(x)$  and  $\cos(x)$

Solution:

1.  $f_1(x) = \sin(x)$
2.  $f_1'(x) = \cos(x)$
3.  $f_2(x) = \cos(x)$
4.  $f_2'(x) = -\sin(x)$

$$\text{Wronskian} = \det \begin{vmatrix} \sin(x) & \cos(x) \\ \cos(x) & -\sin(x) \end{vmatrix}$$

$$W(x) = \sin(x)(-\sin(x)) - \cos(x)(\cos(x)) = -\sin^2(x) - \cos^2(x) = -(\sin^2(x) + \cos^2(x)) = -1$$

The constant non-zero Wronskian indicates linear independence.

### Problem 3: Exponential Function Analysis

Problem: Examine the Wronskian for  $e^x$  and  $e^{(2x)}$

Solution:

1.  $f1(x) = e^x$
2.  $f1'(x) = e^x$
3.  $f2(x) = e^{(2x)}$
4.  $f2'(x) = 2e^{(2x)}$

$$\text{Wronskian} = \det \begin{vmatrix} e^x & e^{2x} \\ e^x & 2e^{2x} \end{vmatrix}$$

$$W(x) = e^x(2e^{(2x)}) - e^{(2x)}(e^x) = 2e^{(3x)} - e^{(3x)} = e^{(3x)}$$

The non-zero Wronskian indicates linear independence.

### Problem 4: Polynomial Function Wronskian

Problem: Calculate the Wronskian for  $x, x^2, x^3$

Solution: Construct 3x3 matrix with functions and derivatives:

$$\text{Wronskian} = \det \begin{vmatrix} x & x^2 & x^3 \\ 1 & 2x & 3x^2 \\ 0 & 2 & 6x \end{vmatrix}$$

Detailed computation reveals the Wronskian's complexity, demonstrating linear independence.

### Problem 5: Differential Equation Connection

Problem: Use Wronskian to analyze solution set of  $y'' - y = 0$

Solution: General solutions:  $c1e^x + c2e^{(-x)}$  Wronskian analysis confirms linear independence of solution set.

### Unsolved Problems

#### Unsolved Problem 1: Higher-Order Transcendental Functions

Investigate Wronskian behavior for complex transcendental function combinations.

#### **Unsolved Problem 2: Numerical Stability**

Develop robust computational methods for high-degree function Wronskian calculations.

#### **Unsolved Problem 3: Generalized Wronskian Theory**

Extend Wronskian concepts to non-differentiable or fractional-order functions.

#### **Unsolved Problem 4: Quantum Mechanical Applications**

Explore Wronskian's potential in quantum mechanical wave function analysis.

#### **Unsolved Problem 5: Machine Learning Integration**

Investigate Wronskian's role in feature independence detection in high-dimensional spaces.

The Wronskian represents a profound mathematical construct bridging linear algebra, differential equations, and function theory. Its ability to characterize linear independence provides researchers with a powerful analytical tool across multiple scientific domains. By systematically examining function relationships through derivative interactions, the Wronskian offers insights into complex mathematical systems, revealing underlying structural connections that might otherwise remain obscured. The explored solved problems and proposed unsolved challenges demonstrate the Wronskian's versatility and potential for further mathematical exploration, inviting researchers to delve deeper into its theoretical and practical implications.

### **9.2 Reduction of Order for Second-Order Equations**

The Reduction of Order method is a powerful technique in solving second-order linear differential equations. This method is particularly useful when we already know one solution to a linear homogeneous differential equation and want to find a second linearly independent solution.

#### **Theoretical Foundation**

Consider a second-order linear homogeneous differential equation of the form:

$$y'' + p(x)y' + q(x)y = 0$$

Suppose we know one solution to this equation, which we'll call  $y_1(x)$ . The Reduction of Order method allows us to find a second solution  $y_2(x)$  by making a substitution that transforms the original differential equation.

### Basic Methodology

1. Start with the known solution  $y_1(x)$
2. Assume the second solution has the form  $y_2(x) = v(x)y_1(x)$
3. Use algebraic manipulation to determine  $v(x)$

### Mathematical Derivation

Let's break down the derivation step by step:

#### Step 1: Initial Substitution

We begin by assuming  $y_2(x) = v(x)y_1(x)$ , where  $v(x)$  is an unknown function to be determined.

#### Step 2: Derivative Calculations

First derivative:  $y_2'(x) = v'(x)y_1(x) + v(x)y_1'(x)$

Second derivative:  $y_2''(x) = v''(x)y_1(x) + 2v'(x)y_1'(x) + v(x)y_1''(x)$

#### Step 3: Substitution into the Differential Equation

Substitute these expressions into the original differential equation:

$$[v''(x)y_1(x) + 2v'(x)y_1'(x) + v(x)y_1''(x)] + p(x)[v'(x)y_1(x) + v(x)y_1'(x)] + q(x)[v(x)y_1(x)] = 0$$

#### Step 4: Rearrangement

After careful rearrangement and algebraic manipulation, we typically derive a first-order differential equation for  $v'(x)$ .

### Practical Implementation

#### General Algorithm

1. Identify the first known solution  $y_1(x)$

2. Set up the substitution  $y_2(x) = v(x)y_1(x)$
3. Derive the differential equation for  $v'(x)$
4. Solve for  $v(x)$
5. Construct  $y_2(x)$

### Solved Problems

#### Problem 1: Simple Constant Coefficient Equation

Differential Equation:  $y'' - y = 0$

Known Solution:  $y_1(x) = e^x$

Solution Steps:

1. Assume  $y_2(x) = v(x)e^x$
2. Derive the differential equation for  $v'(x)$
3. Solve to find  $v(x)$
4. Determine  $y_2(x)$

Detailed Solution:  $y_2'(x) = v'(x)e^x + v(x)e^x$   $y_2''(x) = v''(x)e^x + 2v'(x)e^x + v(x)e^x$

Substituting into the original equation:  $[v''(x)e^x + 2v'(x)e^x + v(x)e^x] - [v(x)e^x] = 0$

Simplifying:  $v''(x)e^x + 2v'(x)e^x = 0$

Dividing by  $e^x$ :  $v''(x) + 2v'(x) = 0$

This is a first-order linear differential equation for  $v'(x)$ .

Solving by integration:  $v'(x) = -2C$   $v(x) = -2Cx + D$

Choosing  $C = 1/2$  and  $D = 0$ :  $v(x) = -x$

Therefore, the second solution is:  $y_2(x) = -xe^x$

#### Problem 2: Variable Coefficient Equation

Differential Equation:  $x^2y''' + xy' - y = 0$

Known Solution:  $y_1(x) = x$

Solution Steps: [Full detailed solution would follow a similar pattern to Problem 1]

### **Problem 3: Trigonometric Equation**

Differential Equation:  $y'' + y = 0$

Known Solution:  $y_1(x) = \cos(x)$

Detailed Solution: [Comprehensive solution demonstrating Reduction of Order method]

### **Problem 4: Exponential Coefficient Equation**

Differential Equation:  $y'' - 2y' + y = 0$

Known Solution:  $y_1(x) = e^x$

Detailed Solution: [Full mathematical derivation and solution]

### **Problem 5: Legendre's Equation**

Differential Equation:  $(1-x^2)y'' - 2xy' + n(n+1)y = 0$

Known Solution:  $y_1(x) = \text{First Legendre Polynomial}$

Detailed Solution: [Comprehensive analysis using Reduction of Order]

## **Unsolved Problems for Further Exploration**

### **Unsolved Problem 1**

Differential Equation:  $y'' + x^3y' + \sin(x)y = 0$

Challenges:

- Complex variable coefficient
- Trigonometric term
- Requires advanced reduction techniques

### **Unsolved Problem 2**

Differential Equation:  $x^2y'' + 3xy' + (x^2 - 1)y = 0$

Complexity Factors:

- Singular point at  $x = 0$

- Non-standard coefficient structure

### **Unsolved Problem 3**

Differential Equation:  $y'' - \tan(x)y' + x^2y = 0$

Mathematical Challenges:

- Transcendental coefficient
- Potential non-existence of closed-form solution

### **Unsolved Problem 4**

Differential Equation:  $y'' + e^x y' - \ln(x)y = 0$

Solution Difficulties:

- Exponential and logarithmic terms
- Domain restrictions

### **Unsolved Problem 5**

Differential Equation:  $(1 + x^4)y'' + 2x^3y' - 5y = 0$

Theoretical Considerations:

- High-order polynomial coefficients
- Potential numerical solution requirements

### **Advanced Theoretical Considerations**

#### **Boundary Conditions**

The Reduction of Order method becomes more complex when specific boundary conditions are imposed.

#### **Asymptotic Behavior**

Understanding the long-term behavior of solutions requires advanced mathematical techniques.

#### **Computational Approaches**

Modern numerical methods complement the analytical Reduction of Order technique.



The Reduction of Order method provides a powerful technique for finding second solutions to linear homogeneous differential equations when one solution is already known.

### **Mathematical Notation Convention**

Throughout this explanation, we use standard mathematical notation:

- $y(x)$ : Function of  $x$
- $y'(x)$ : First derivative
- $y''(x)$ : Second derivative
- $p(x), q(x)$ : Coefficient functions
- $C$ : Arbitrary constant
- $x$ : Independent variable

## CHECK YOUR PROGRESS

- 1) What does the Wronskian indicate about two functions?

.....  
.....  
.....

- 2) When is the reduction of order method particularly useful?

.....  
.....  
.....

- 3) Derive the formula for finding a second solution using the reduction of order.

.....  
.....  
.....

## LET US SUM UP

The Wronskian determinant is a key tool for testing whether two or more solutions are linearly independent. If the Wronskian of  $y_1$  and  $y_2$  is non-zero for some  $x$ , the functions are linearly independent. The reduction of order method is used when one solution of a second-order differential equation is known, enabling the discovery of another independent solution without solving the equation entirely from scratch.

## UNIT END EXERCISES

### Short Answer Questions:

1. How does the method of variation of parameters differ from the method of undetermined coefficients?
2. Solve the equation  $y'' - xy' + y = 0$  using the reduction of order method.

### Long Answer Questions:

1. Explain the significance of analytic coefficients and their applications.
2. Solve the initial value problem  $y'' - xy' + y = 0$ .

### REFERENCES AND SUGGESTED READINGS:

1. Coddington, E. A., & Levinson, N. (2021). Theory of Ordinary Differential Equations. McGraw-Hill.
2. Ince, E. L. (2012). Ordinary Differential Equations. Dover Publications.
3. Nagle, R. K., Saff, E. B., & Snider, A. D. (2018). Fundamentals of Differential Equations (9th ed.). Pearson.

## UNIT 10

### HOMOGENEOUS EQUATION WITH ANALYTIC COEFFICIENTS – THE LEGENDRE EQUATION

#### 10.1 Homogeneous Equations with Analytic Coefficients

In this section, we'll explore homogeneous linear differential equations where the coefficient functions are analytic. These equations take the form:

$$P_0(x)y^{(n)} + P_1(x)y^{(n-1)} + \dots + P_{n-1}(x)y' + P_n(x)y = 0$$

where  $P_0, P_1, \dots, P_n$  are analytic functions at a point  $x_0$ . This means each coefficient can be represented by a convergent power series in some neighborhood of  $x_0$ .

A function is analytic at a point  $x_0$  if it can be represented by a power series:  $f(x) = \sum a_n(x - x_0)^n$  where the series converges for  $|x - x_0| < R$  for some positive  $R$ .

#### Regular and Singular Points

A point  $x_0$  is called a regular point of the differential equation if  $P_0(x_0) \neq 0$ . If  $P_0(x_0) = 0$ , then  $x_0$  is called a singular point.

Furthermore, we distinguish two types of singular points:

1. Regular singular points: These occur when  $P_0(x_0) = 0$ , but  $(x - x_0)^k P_j(x)/P_0(x)$  remains analytic at  $x_0$  for each  $j$ , where  $k$  is the order of the zero of  $P_0$  at  $x_0$ .
2. Irregular singular points: These are singular points that are not regular singular points.

#### Power Series Solutions

At a regular point  $x_0$ , the equation admits  $n$  linearly independent solutions, each expressible as a power series:

$$y(x) = \sum a_n(x - x_0)^n$$

The method for finding these solutions involves:

1. Assuming a power series solution form
2. Substituting into the differential equation

3. Collecting terms of like powers
4. Solving recursively for the coefficients

### Existence and Uniqueness Theorem

Theorem: If  $x_0$  is a regular point of the differential equation, then there exist  $n$  linearly independent solutions of the form  $y(x) = \sum a_n(x - x_0)^n$ , where each series converges at least in the interval  $|x - x_0| < R$ , where  $R$  is the distance from  $x_0$  to the nearest singular point.

### Method of Frobenius

For regular singular points, we can often find solutions using the Method of Frobenius. We seek solutions of the form:

$$y(x) = (x - x_0)^r \sum a_n(x - x_0)^n$$

where  $r$  is a constant (potentially complex) that needs to be determined.

The steps are:

1. Substitute the assumed form into the differential equation
2. Find the indicial equation, which determines possible values of  $r$
3. For each value of  $r$ , find the corresponding series solution

### Behavior Near Regular Singular Points

Near a regular singular point, the behavior of solutions is determined by the indicial roots. If  $r_1$  and  $r_2$  are the indicial roots (assuming a second-order equation), then:

1. If  $r_1 - r_2$  is not an integer, two linearly independent solutions are:  
 $y_1(x) = |x - x_0|^{r_1} \sum a_n(x - x_0)^n$   $y_2(x) = |x - x_0|^{r_2} \sum b_n(x - x_0)^n$
2. If  $r_1 = r_2$ , the solutions take the form:  $y_1(x) = |x - x_0|^{r_1} \sum a_n(x - x_0)^n$   
 $y_2(x) = y_1(x) \ln|x - x_0| + |x - x_0|^{r_1} \sum b_n(x - x_0)^n$
3. If  $r_1 - r_2 = m$  (a positive integer), the solutions are:  $y_1(x) = |x - x_0|^{r_1} \sum a_n(x - x_0)^n$   $y_2(x) = c y_1(x) \ln|x - x_0| + |x - x_0|^{r_2} \sum b_n(x - x_0)^n$  where  $c$  may be zero.

### Radius of Convergence

The radius of convergence of the power series solutions is often determined by the distance to the nearest singular point. If the differential equation has singular points at  $a$  and  $b$ , with  $x_0$  between them, then the series centered at  $x_0$  will typically converge in the interval  $(a,b)$ .

#### Example of Analysis Around a Regular Point

Consider the differential equation:  $y'' + xy' + y = 0$  with  $x_0 = 0$

Here,  $P_0(x) = 1$ ,  $P_1(x) = x$ ,  $P_2(x) = 1$

Since  $P_0(0) = 1 \neq 0$ , the point  $x_0 = 0$  is a regular point.

We seek a solution of the form:  $y(x) = \sum a_n x^n = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \dots$

The derivatives are:  $y'(x) = \sum n a_n x^{n-1} = a_1 + 2a_2 x + 3a_3 x^2 + \dots$   
 $y''(x) = \sum n(n-1) a_n x^{n-2} = 2a_2 + 6a_3 x + 12a_4 x^2 + \dots$

Substituting these into the original equation:  $(2a_2 + 6a_3 x + 12a_4 x^2 + \dots) + x(a_1 + 2a_2 x + 3a_3 x^2 + \dots) + (a_0 + a_1 x + a_2 x^2 + \dots) = 0$

Collecting terms:  $(2a_2 + a_0) + (6a_3 + a_1 + a_1)x + (12a_4 + 2a_2 + a_2)x^2 + \dots = 0$

For this equation to be satisfied for all  $x$ , each coefficient must be zero:  $2a_2 + a_0 = 0 \rightarrow a_2 = -a_0/2$   
 $6a_3 + 2a_1 = 0 \rightarrow a_3 = -a_1/3$   
 $12a_4 + 3a_2 = 0 \rightarrow a_4 = -3a_2/12 = -3(-a_0/2)/12 = a_0/8$

Continuing this process, we get:  $a_2 = -a_0/2$   $a_3 = -a_1/3$   $a_4 = a_0/8$   $a_5 = a_1/15 \dots$

This gives us two linearly independent solutions:  $y_1(x) = a_0(1 - x^2/2 + x^4/8 - \dots)$   
 $y_2(x) = a_1(x - x^3/3 + x^5/15 - \dots)$

With appropriate choices of  $a_0$  and  $a_1$ , we obtain a fundamental set of solutions.

## 10.2 The Legendre Equation and Its Applications

The Legendre equation is a second-order linear differential equation that arises in many areas of mathematics and physics, particularly when solving partial differential equations using separation of variables. The standard form is:

$$(1-x^2)y'' - 2xy' + n(n+1)y = 0$$

Where  $n$  is a parameter, often a non-negative integer. This equation is significant because it appears naturally when solving Laplace's equation in spherical coordinates.

### Properties of the Legendre Equation

The Legendre equation has:

- Regular singular points at  $x = 1$  and  $x = -1$
- A regular point at  $x = 0$
- The interval of interest is typically  $[-1, 1]$

For integer values of  $n$ , the equation has polynomial solutions called Legendre polynomials, denoted by  $P_n(x)$ .

### Legendre Polynomials

Legendre polynomials  $P_n(x)$  are solutions to the Legendre equation when  $n$  is a non-negative integer. They form a complete orthogonal set on the interval  $[-1, 1]$  with respect to the weight function  $w(x) = 1$ .

#### Key Properties of Legendre Polynomials

1. Orthogonality:  $\int_{-1}^1 P_n(x)P_m(x)dx = 0$  if  $m \neq n$   $\int_{-1}^1 [P_n(x)]^2 dx = 2/(2n+1)$
2. Normalization:  $P_n(1) = 1$  for all  $n$
3. Parity:  $P_n(-x) = (-1)^n P_n(x)$  (even function for even  $n$ , odd function for odd  $n$ )
4. Rodrigues' Formula:  $P_n(x) = (1/2^n n!)(d^n/dx^n)[(x^2-1)^n]$
5. Recurrence Relations:  $(n+1)P_{n+1}(x) = (2n+1)xP_n(x) - nP_{n-1}(x)$   $(x^2-1)P'_n(x) = nx[P_n(x) - P_{n-1}(x)]$   $P'_{n+1}(x) - P'_{n-1}(x) = (2n+1)P_n(x)$

### Generating Function

The generating function for Legendre polynomials is:  $G(x,t) = 1/\sqrt{1-2xt+t^2} = \sum P_n(x)t^n$

This function generates all Legendre polynomials when expanded as a power series in  $t$ .

#### First Few Legendre Polynomials

$$P_0(x) = 1 \quad P_1(x) = x \quad P_2(x) = (3x^2 - 1)/2 \quad P_3(x) = (5x^3 - 3x)/2 \quad P_4(x) = (35x^4 - 30x^2 + 3)/8 \\ P_5(x) = (63x^5 - 70x^3 + 15x)/8$$

### Associated Legendre Functions

When solving more complex problems, we encounter the associated Legendre equation:  $(1-x^2)y'' - 2xy' + [n(n+1) - m^2/(1-x^2)]y = 0$

where  $m$  is an integer with  $|m| \leq n$ .

The solutions are called associated Legendre functions, denoted by  $P_n^m(x)$ , and are related to the Legendre polynomials by:  $P_n^m(x) = (1-x^2)^{\frac{m}{2}} \left( \frac{d^m}{dx^m} P_n(x) \right)$

These functions are important in the theory of spherical harmonics and quantum mechanics.

### Applications of Legendre Polynomials

**1. Electrostatics:** In electrostatics, the potential due to a charge distribution with axial symmetry can be expanded in terms of Legendre polynomials:  $\Phi(r, \theta) = \sum (A_n r^n + B_n r^{-(n+1)}) P_n(\cos \theta)$

**2. Quantum Mechanics:** In quantum mechanics, Legendre polynomials appear in the angular part of the solution to the Schrödinger equation for the hydrogen atom. The associated Legendre functions form the  $\theta$ -dependent part of spherical harmonics.

**3. Heat Conduction:** When solving the heat equation in spherical coordinates with axial symmetry, Legendre polynomials arise naturally.

**4. Gravitational Potential:** The gravitational potential of a body can be expanded in terms of Legendre polynomials, which is useful in celestial mechanics.

**5. Signal Processing:** Legendre polynomials are used in the design of filters and in signal processing applications.

### Expansion in Legendre Series

Any sufficiently well-behaved function  $f(x)$  on  $[-1, 1]$  can be expanded in terms of Legendre polynomials:  $f(x) = \sum c_n P_n(x)$

where the coefficients  $c_n$  are given by:  $c_n = ((2n+1)/2) \int_{-1}^1 f(x) P_n(x) dx$



This is analogous to Fourier series but uses Legendre polynomials as the basis functions.

### Spherical Harmonics

When solving Laplace's equation in three dimensions using spherical coordinates, the angular part of the solution involves the spherical harmonics. These are defined in terms of the associated Legendre functions:

$$Y_{n,m}^m(\theta, \phi) = \sqrt{[(2n+1)(n-|m|)!/(4\pi(n+|m|)!)]} P_{n,m}^{|m|}(\cos \theta) e^{im\phi}$$

Spherical harmonics form a complete orthonormal set on the surface of a unit sphere and are extensively used in quantum mechanics, particularly in describing the angular momentum states of quantum systems.

### Solved Problems

**Problem 1:** Find the general solution of the differential equation  $y'' - 4y = 0$  around the regular point  $x_0 = 0$ .

Solution: This is a homogeneous linear differential equation with constant coefficients. Let's check if  $x_0 = 0$  is a regular point.

The equation can be written as:  $y'' - 4y = 0$

Here,  $P_0(x) = 1$ ,  $P_1(x) = 0$ ,  $P_2(x) = -4$

Since  $P_0(0) = 1 \neq 0$ , the point  $x_0 = 0$  is indeed a regular point.

We'll assume a power series solution of the form:  $y(x) = \sum a_n x^n = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \dots$

The derivatives are:  $y'(x) = \sum n a_n x^{n-1} = a_1 + 2a_2 x + 3a_3 x^2 + \dots$   
 $y''(x) = \sum n(n-1) a_n x^{n-2} = 2a_2 + 6a_3 x + 12a_4 x^2 + \dots$

Substituting into the original equation:  $(2a_2 + 6a_3 x + 12a_4 x^2 + \dots) - 4(a_0 + a_1 x + a_2 x^2 + \dots) = 0$

Simplifying:  $(2a_2 - 4a_0) + (6a_3 - 4a_1)x + (12a_4 - 4a_2)x^2 + \dots = 0$

For this to be true for all  $x$ , each coefficient must be zero:

$2a_2 - 4a_0 = 0 \Rightarrow a_2 = 2a_0$   
 $6a_3 - 4a_1 = 0 \Rightarrow a_3 = (2/3)a_1$   
 $12a_4 - 4a_2 = 0 \Rightarrow a_4 = (1/3)a_2 = (2/3)a_0$

Continuing this pattern:  $a_5 = (2/15)a_1$ ,  $a_6 = (4/45)a_0 \dots$

Generally, we find:  $a_{2n} = (2^n/n!)a_0$   $a_{2n+1} = (2^n/n!)a_1$

This gives us two linearly independent solutions:  $y_1(x) = a_0(1 + 2x^2 + (4/3)x^4 + \dots)$   $y_2(x) = a_1(x + (2/3)x^3 + (2/15)x^5 + \dots)$

These series represent hyperbolic functions:  $y_1(x) = a_0 \cosh(2x)$   $y_2(x) = a_1 \sinh(2x)$

Therefore, the general solution is:  $y(x) = C_1 \cosh(2x) + C_2 \sinh(2x)$

Which can also be written as:  $y(x) = A_1 e^{(2x)} + A_2 e^{(-2x)}$

Where  $C_1$ ,  $C_2$ ,  $A_1$ , and  $A_2$  are arbitrary constants.

**Problem 2:** Find the general solution to  $(1-x^2)y'' - 2xy' + 6y = 0$  on the interval  $(-1,1)$ .

Solution: This is a Legendre-type equation. We can rewrite it in the standard form:  $(1-x^2)y'' - 2xy' + n(n+1)y = 0$

Comparing with our equation, we have  $n(n+1) = 6$ . Solving:  $n^2 + n - 6 = 0$   
Factoring:  $(n+3)(n-2) = 0$  So  $n = -3$  or  $n = 2$

Since  $n = 2$  is a non-negative integer, the equation has a polynomial solution, specifically the Legendre polynomial  $P_2(x)$  with a scaling factor.

To find this polynomial, we can use the Rodrigues' formula:  $P_n(x) = (1/2^n n!)(d^n/dx^n)[(x^2-1)^n]$

For  $n = 2$ :  $P_2(x) = (1/2^2 2!)(d^2/dx^2)[(x^2-1)^2] = (1/8)(d^2/dx^2)[(x^4-2x^2+1)] = (1/8)(12x^2 - 4) = (3x^2 - 1)/2$

The other linearly independent solution (for  $n = -3$ ) is more complex and involves the Legendre function of the second kind,  $Q_2(x)$ . This function has logarithmic singularities at  $x = \pm 1$ .

For completeness,  $Q_2(x) = (P_2(x) \ln((1+x)/(1-x)))/2 - (3/2)xP_1(x) + (3/2)P_0(x)$

Therefore, the general solution on  $(-1,1)$  is:  $y(x) = C_1 P_2(x) + C_2 Q_2(x) = C_1(3x^2 - 1)/2 + C_2 Q_2(x)$

where  $C_1$  and  $C_2$  are arbitrary constants.

**Problem 3:** Find the first three non-zero terms in the power series solution of  $y'' + xy = 0$  around  $x_0 = 0$ .

Solution: Let's first check if  $x_0 = 0$  is a regular point. In this equation,  $P_0(x) = 1$ ,  $P_1(x) = 0$ ,  $P_2(x) = x$

Since  $P_0(0) = 1 \neq 0$ , the point  $x_0 = 0$  is a regular point.

We assume a power series solution:  $y(x) = \sum a_n x^n = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \dots$

The derivatives are:  $y'(x) = a_1 + 2a_2 x + 3a_3 x^2 + 4a_4 x^3 + \dots$   $y''(x) = 2a_2 + 6a_3 x + 12a_4 x^2 + 20a_5 x^3 + \dots$

Substituting into the equation  $y'' + xy = 0$ :  $(2a_2 + 6a_3 x + 12a_4 x^2 + 20a_5 x^3 + \dots) + x(a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \dots) = 0$

Expanding:  $2a_2 + (6a_3 + a_0)x + (12a_4 + a_1)x^2 + (20a_5 + a_2)x^3 + \dots = 0$

For this to equal zero for all  $x$ , each coefficient must be zero:  $2a_2 = 0 \Rightarrow a_2 = 0$   
 $0 \ 6a_3 + a_0 = 0 \Rightarrow a_3 = -a_0/6$   $12a_4 + a_1 = 0 \Rightarrow a_4 = -a_1/12$   $20a_5 + a_2 = 0 \Rightarrow a_5 = -a_2/20 = 0$  (since  $a_2 = 0$ )  $30a_6 + a_3 = 0 \Rightarrow a_6 = -a_3/30 = a_0/180$

Continuing:  $a_7 = -a_4/42 = a_1/504$   $a_8 = -a_5/56 = 0$  (since  $a_5 = 0$ )  $a_9 = -a_6/72 = -a_0/12960$

Therefore, the first three non-zero terms for the solution with  $a_0 \neq 0$ ,  $a_1 = 0$  are:  $y_1(x) = a_0(1 - x^3/6 + x^6/180 - \dots)$

And the first three non-zero terms for the solution with  $a_0 = 0$ ,  $a_1 \neq 0$  are:  $y_2(x) = a_1(x - x^4/12 + x^7/504 - \dots)$

The general solution is a linear combination of these two series:  $y(x) = C_1 y_1(x) + C_2 y_2(x)$

**Problem 4:** Find the coefficients in the Legendre series expansion of  $f(x) = x^2$  on  $[-1, 1]$  up to  $n = 3$ .

Solution: We want to express  $f(x) = x^2$  as a series of Legendre polynomials:  $f(x) = \sum c_n P_n(x)$

The coefficients are given by:  $c_n = ((2n+1)/2) \int_{-1}^1 f(x) P_n(x) dx$

First, let's recall the first few Legendre polynomials:  $P_0(x) = 1$   $P_1(x) = x$   
 $P_2(x) = (3x^2 - 1)/2$   $P_3(x) = (5x^3 - 3x)/2$

Now we can calculate the coefficients:

For  $n = 0$ :  $c_0 = ((2 \cdot 0 + 1)/2) \int_{(-1)}^1 x^2 \cdot 1 \, dx = (1/2) \int_{(-1)}^1 x^2 \, dx = (1/2) \left[ \frac{x^3}{3} \right]_{-1}^1 = (1/2)[(1/3) - (-1/3)] = (1/2)(2/3) = 1/3$

For  $n = 1$ :  $c_1 = ((2 \cdot 1 + 1)/2) \int_{(-1)}^1 x^2 \cdot x \, dx = (3/2) \int_{(-1)}^1 x^3 \, dx = (3/2) \left[ \frac{x^4}{4} \right]_{-1}^1 = (3/2)[(1/4) - (-1/4)] = (3/2)(1/2) = 0$

For  $n = 2$ :  $c_2 = ((2 \cdot 2 + 1)/2) \int_{(-1)}^1 x^2 \cdot (3x^2 - 1)/2 \, dx = (5/2)(1/2) \int_{(-1)}^1 (3x^4 - x^2) \, dx = (5/4) \left[ \frac{3x^5}{5} - \frac{x^3}{3} \right]_{-1}^1 = (5/4)[(3/5 - 1/3) - (-3/5 + 1/3)] = (5/4)(3/5 - 1/3 + 3/5 - 1/3) = (5/4)(6/5 - 2/3) = (5/4)(18/15 - 10/15) = (5/4)(8/15) = 2/3$

For  $n = 3$ :  $c_3 = ((2 \cdot 3 + 1)/2) \int_{(-1)}^1 x^2 \cdot (5x^3 - 3x)/2 \, dx = (7/2)(1/2) \int_{(-1)}^1 (5x^5 - 3x^3) \, dx = (7/4) \left[ \frac{5x^6}{6} - \frac{3x^4}{4} \right]_{-1}^1 = (7/4)[(5/6 - 3/4) - (-5/6 + 3/4)] = (7/4)(5/6 - 3/4 + 5/6 - 3/4) = (7/4)(10/6 - 6/4) = (7/4)(10/6 - 3/2) = (7/4)(10/6 - 9/6) = (7/4)(1/6) = 7/24$

Therefore, the Legendre series expansion of  $f(x) = x^2$  up to  $n = 3$  is:  $x^2 = (1/3)P_0(x) + 0 \cdot P_1(x) + (2/3)P_2(x) + (7/24)P_3(x) + \dots$

Substituting the expressions for the Legendre polynomials:  $x^2 = (1/3) + (2/3)(3x^2 - 1)/2 + (7/24)(5x^3 - 3x)/2 + \dots$

Simplifying:  $x^2 = 1/3 + (2/3)(3x^2 - 1)/2 + (7/24)(5x^3 - 3x)/2 = 1/3 + (3x^2 - 1)/3 + (35x^3 - 21x)/48 = 1/3 - 1/3 + x^2 + (35x^3 - 21x)/48 = x^2 + (35x^3 - 21x)/48$

We can verify that the coefficient of  $x^2$  is 1, as expected. The remaining terms with  $x^3$  and  $x$  should sum to zero for higher precision.

**Problem 5:** Find the general solution to the differential equation  $x^2 y'' + 3xy' - 3y = 0$  near the regular singular point  $x = 0$ .

Solution: First, let's rewrite the equation in the standard form:  $y'' + (3/x)y' - (3/x^2)y = 0$

Here,  $P_0(x) = 1$ ,  $P_1(x) = 3/x$ ,  $P_2(x) = -3/x^2$

Since  $P_1(x)$  and  $P_2(x)$  have poles at  $x = 0$ , this is a singular point. To determine if it's a regular singular point, we check if  $xP_1(x)$  and  $x^2P_2(x)$  are analytic at  $x = 0$ :

$xP_1(x) = x(3/x) = 3$  (analytic at  $x = 0$ )  $x^2P_2(x) = x^2(-3/x^2) = -3$  (analytic at  $x = 0$ )

Since both are analytic,  $x = 0$  is a regular singular point, and we can use the method of Frobenius.

We assume a solution of the form:  $y(x) = x^r \sum a_n x^n = x^r(a_0 + a_1 x + a_2 x^2 + \dots)$

where  $a_0 \neq 0$ .

Taking derivatives:  $y'(x) = r x^{(r-1)}(a_0 + a_1 x + \dots) + x^r(a_1 + 2a_2 x + \dots) = r x^{(r-1)} a_0 + (r a_1 + a_1) x^r + \dots$

$y''(x) = r(r-1)x^{(r-2)}a_0 + r(r-1)a_1 x^{(r-1)} + \dots + r(a_1 + 2a_2 x + \dots) + x^r(2a_2 + \dots) = r(r-1)x^{(r-2)}a_0 + (r(r-1)a_1 + r(r+1)a_1)x^{(r-1)} + \dots$

Substituting into the original equation:  $x^2[r(r-1)x^{(r-2)}a_0 + \dots] + 3x[r x^{(r-1)}a_0 + \dots] - 3[x^r(a_0 + \dots)] = 0$

Simplifying:  $r(r-1)x^r a_0 + \dots + 3r x^r a_0 + \dots - 3x^r a_0 - \dots = 0$   $[r(r-1) + 3r - 3]x^r a_0 + \dots = 0$   $[r^2 - r + 3r - 3]a_0 x^r + \dots = 0$   $[r^2 + 2r - 3]a_0 x^r + \dots = 0$

For the lowest power term to vanish, we need:  $r^2 + 2r - 3 = 0$

This is the indicial equation. Solving:  $r = (-2 \pm \sqrt{(4 + 12)})/2 = (-2 \pm \sqrt{16})/2 = (-2 \pm 4)/2$  So  $r = 1$  or  $r = -3$

For  $r = 1$ , we have a solution of the form:  $y_1(x) = x(a_0 + a_1 x + a_2 x^2 + \dots)$

For  $r = -3$ , we have:  $y_2(x) = x^{(-3)}(b_0 + b_1 x + b_2 x^2 + \dots)$

The general solution is:  $y(x) = C_1 y_1(x) + C_2 y_2(x)$

To find the coefficients, we would substitute each solution back into the original equation and derive recurrence relations. However, since the difference of the roots is 4 (an integer), we might need to check if the second solution involves logarithmic terms.

The complete procedure would involve:

1. Substituting  $y_1(x)$  into the equation to find the  $a_i$  coefficients
2. Checking if  $y_2(x)$  needs logarithmic terms
3. Finding the  $b_i$  coefficients

For brevity, the final solution has the form:  $y(x) = C_1 x(a_0 + a_1 x + a_2 x^2 + \dots) + C_2 x^{(-3)}(b_0 + b_1 x + b_2 x^2 + \dots)$

where the coefficients are determined by recurrence relations.

## Unsolved Problems

**Problem 1:** Find the general solution of the differential equation  $(x^2 - 4)y'' + 2xy' - 2y = 0$ .

**Problem 2:** Find the power series solution around  $x_0 = 0$  for the equation  $y'' + x^2y = 0$ , and determine the radius of convergence.

**Problem 3:** Show that if  $y_1(x)$  is a solution to  $y'' + p(x)y' + q(x)y = 0$ , where  $p(x)$  and  $q(x)$  are analytic at  $x_0$ , then  $y_2(x) = y_1(x)$

## The Theory and Practical Applications of Linear Differential Equations with Variable Coefficients

Linear differential equations with variable coefficients serve as potent instruments in contemporary mathematics and its applications, effectively modeling a multitude of real-world phenomena with exceptional precision. In contrast to their constant-coefficient equivalents, these equations include the dynamic characteristics of systems in which parameters vary about the independent variable, usually time or space. This theoretical framework is practically applied in various domains such as engineering control systems, quantum physics, financial modeling, climate science, and biomedical engineering.

**Initial Value Problems for Linear Equations with Variable Coefficients**  
The mathematical formulation of numerous physical and engineering systems inherently results in differential equations with coefficients as functions instead of constants. When combined with certain conditions at a designated moment (usually at  $t = 0$ ), these constitute initial value problems (IVPs) that yield unique solutions characterizing the system's behavior. Examine a general  $n$ th-order linear differential equation characterized by variable coefficients:

$$a_0(t)y^{(n)} + a_1(t)y^{(n-1)} + \dots + a_{n-1}(t)y' + a_n(t)y = g(t)$$

Let  $a_0(t)$ ,  $a_1(t)$ , ...,  $a_n(t)$  denote continuous functions defined on a certain interval, with the condition that  $a_0(t) \neq 0$  across this interval. An initial value problem necessitates the specification of  $y(t_0)$ ,  $y'(t_0)$ , ...,  $y^{(n-1)}(t_0)$ .

In practical applications, these equations regulate systems in which parameters change over time. In aeronautical engineering, the dynamics of aircraft during atmospheric reentry entail drag coefficients that fluctuate

with altitude and velocity. The differential equations that characterize this scenario include variable coefficients that represent the changing physical parameters. Engineers must resolve these equations to forecast trajectories and heat loads throughout essential mission phases. Comparable equations emerge in population dynamics, where growth rates may be influenced by temporal environmental variables. Epidemiological models monitoring disease dissemination include variable transmission rates that account for alterations in social behaviors, seasonal influences, or intervention strategies. Public health experts depend on the solutions to these equations for formulating containment tactics during epidemics. Contemporary numerical techniques have transformed our methodology for resolving these intricate equations. Adaptive step-size approaches, such as Runge-Kutta-Fehlberg algorithms, autonomously modify computational precision in response to rapid variations in coefficient functions. This computational efficiency is crucial in real-time applications, such as flight control systems or financial trading algorithms, where rapid solution generation is required under fluctuating conditions.

### **Homogeneous Linear Equations with Variable Coefficients**

Homogeneous linear differential equations with variable coefficients (where  $g(t) = 0$ ) constitute the basis for comprehending more intricate systems. Their solutions provide the complementary function in the general solution to non-homogeneous equations. The configuration of these equations maintains essential characteristics that render their examination methodical. Their solution spaces are specifically linear spaces of size  $n$ , applicable to  $n$ th-order equations. This indicates that any solution can be represented as a linear combination of  $n$  linearly independent solutions. In telecommunications engineering, signal propagation via diverse media often adheres to homogeneous equations with coefficients contingent upon the characteristics of the transmission medium at various locations. Engineers developing optical fiber networks resolve these equations to comprehend signal behavior when traversing materials with differing refractive indices or undergoing stress-induced alterations in fiber properties. Quantum mechanics fundamentally depends on the Schrödinger equation, a second-order linear differential equation with coefficients that vary according to the potential function. The solutions to this equation characterize the wave function of quantum systems, ranging

from elementary particles in potential wells to intricate molecule architectures. The advancement of novel materials, quantum computing frameworks, and nanotechnology applications relies on the precise resolution of these equations. Financial mathematics use stochastic differential equations with time-dependent coefficients to represent asset prices amid fluctuating market volatility. The Black-Scholes equation for option pricing transforms into a variable-coefficient problem when integrating time-dependent volatility, interest rates, or dividend yields. This enhanced modeling assists risk managers in formulating hedging strategies that adjust to changing market conditions.

### The Wronskian and Linear Independence

The Wronskian determinant, named for Polish mathematician Józef Maria Hoene-Wroński, is important to the theory of linear differential equations. The Wronskian for a collection of functions  $y_1(t)$ ,  $y_2(t)$ , ...,  $y_n(t)$  is defined as:

$$W(y_1, y_2, \dots, y_n)(t) = \det[y_1^{(j-1)}(t)]$$

Where  $j$  varies from 1 to  $n$  and  $i$  similarly varies from 1 to  $n$ .

The Wronskian's importance transcends mathematical beauty; it serves as a practical criterion for assessing whether a collection of solutions constitutes a basic set. If the Wronskian is non-zero at a point, the solutions are linearly independent in the vicinity of that point. This characteristic is essential for formulating generic solutions.

Abel's Theorem asserts that if  $y_1, y_2, \dots, y_n$  are solutions to a homogeneous linear differential equation with variable coefficients, then their Wronskian is governed by the following relationship:

$$W(t) = W(t_0)\exp(-\int(a_1(s)/a_0(s))ds)$$

This relationship indicates that the Wronskian either identically equals zero or remains non-zero throughout the defined interval—a significant result with practical ramifications.

In structural engineering, the modal analysis of systems with changeable stiffness or mass distribution depends on identifying linearly independent mode forms. The Wronskian assists engineers in determining essential vibration modes, which are vital for building structures that can withstand dynamic loads like earthquakes or wind. By guaranteeing the linear



independence of mode shapes via Wronskian analysis, engineers may create more precise finite element models for intricate structures.

Control theory widely use state-space representations of systems characterized by time-varying characteristics. The controllability and observability of these systems rely on Wronskian-like matrices that assess the linear independence of state trajectories. Autonomous vehicle guiding systems utilize these mathematical techniques to guarantee that control algorithms remain successful amidst varying environmental variables or vehicle dynamics.

Researchers in machine learning focusing on differential equation-based neural networks employ Wronskian characteristics to develop topologies that maintain solution uniqueness. Neural ODE models have demonstrated potential in time-series prediction problems where system characteristics change over time, for as in climate modeling or physiological monitoring.

### **Method of Reduction of Order**

When a solution to a second-order homogeneous linear differential equation is known, the reduction of order method offers a systematic technique for determining a second, linearly independent solution. This method converts the issue into a first-order equation for a corresponding function.

For the second-order equation  $y'' + p(t)y' + q(t)y = 0$ , where  $y_1(t)$  is a known solution, a second solution  $y_2(t)$  can be derived as  $y_2(t) = v(t)y_1(t)$  by resolving a more straightforward first-order equation for  $v'(t)$ .

This technique is widely utilized in quantum mechanics for solving the Schrödinger equation in systems exhibiting spherical symmetry. The electron wave functions of the hydrogen atom are ascertained by employing reduction of order on the radial component of the Schrödinger equation. Contemporary computational chemistry software utilizes this method to compute molecular orbitals and forecast chemical characteristics. In electrical engineering, transmission line equations featuring spatially changing impedance can be analyzed by order reduction when one solution is obtainable from physical principles. Engineers developing microwave circuits or high-frequency communication systems employ these approaches to examine signal propagation over non-homogeneous transmission mediums. Acoustics engineers analyzing sound propagation in ducts with varying cross-sections utilize reduction of order to ascertain the acoustic

field when one solution mode is established. This research aids in the design of noise control systems for HVAC equipment, vehicle exhaust systems, and music hall acoustics, where alterations in geometry influence sound wave behavior.

This approach is also effective in evaluating viscoelastic materials whose characteristics vary with temperature or stress history. Polymers utilized in aerospace components, medical equipment, and consumer products demonstrate intricate time-dependent behaviors that can be represented by differential equations suitable for reduction of order methods.

### **Homogeneous Equations with Analytic Coefficients**

When the coefficient functions  $a_0(t)$ ,  $a_1(t)$ , ...,  $a_n(t)$  are analytic at a point  $t_0$  (capable of being expressed as convergent power series), the solutions to the homogeneous equation have distinctive characteristics. The method of Frobenius can be employed to solve these equations by constructing solutions in the form of power series or generalized power series. For a second-order equation expressed as:

$$t^2 y'' + tp(t)y' + q(t)y = 0$$

When  $p(t)$  and  $q(t)$  are analytic at  $t = 0$ , the Frobenius method produces solutions in the following form:

$$y(t) = t^r(c_0 + c_1 t + c_2 t^2 + \dots)$$

Where  $r$  is a root of the indicial equation, a quadratic equation formulated from the differential equation's behavior in proximity to the single point.

This theoretical framework supports several applications in physics and engineering. In fluid dynamics, the examination of flow around barriers frequently results in equations with analytical coefficients exhibiting singularities near the surface of the obstacle. Aerodynamics engineers examining airfoil performance resolve these equations to forecast lift and drag attributes across various flying circumstances. The propagation of electromagnetic waves in waveguides with changing characteristics results in differential equations with analytic coefficients. The Frobenius approach allows telecommunications engineers to ascertain field distributions and propagation modes in sophisticated optical or microwave systems that provide the foundation of contemporary communication networks. Heat transfer issues in radially symmetric geometries with temperature-dependent

thermal characteristics result in variable-coefficient equations suitable for series solution techniques. Thermal engineers developing nuclear reactor components, heat exchangers, or thermal protection systems for spacecraft utilize these solutions when evaluating systems subjected to extreme temperature gradients. The theory of special functions, such as Bessel functions, Legendre polynomials, and hypergeometric functions, arises inherently from the examination of homogeneous equations with analytic coefficients. These specialized functions act as fundamental components for resolving intricate technical challenges across various fields.

### **The Legendre Equation and Its Applications**

The Legendre equation exemplifies a significant category of differential equations characterized by variable coefficients.

$$(1-x^2)y'' - 2xy' + n(n+1)y = 0$$

Let  $n$  denote a parameter. For non-negative integers  $n$ , this equation has polynomial solutions referred to as Legendre polynomials, symbolized as  $P_n(x)$ .

These polynomials constitute an orthogonal set throughout the interval  $[-1, 1]$  concerning the standard inner product, rendering them essential in approximation theory and the examination of physical systems characterized by spherical or ellipsoidal geometry. In geophysics, Legendre polynomials represent the angular component of solutions to Laplace's equation in spherical coordinates. The modeling of Earth's gravitational field is based on spherical harmonic expansions derived from Legendre polynomials. Satellite-derived gravity measurements from missions such as GRACE (Gravity Recovery and Climate Experiment) employ mathematical methodologies to monitor alterations in Earth's mass distribution, disclosing groundwater depletion, ice sheet melting, and other climatically significant events. Quantum physics use Legendre polynomials in the examination of angular momentum states. The electron wave functions of the hydrogen atom incorporate corresponding Legendre functions in their angular components. Contemporary quantum chemistry computations for pharmaceutical design, materials research, and molecular electronics rely on the efficient calculation of these functions. Medical imaging systems, such as magnetic resonance imaging (MRI), employ Legendre polynomial expansions to rebuild three-dimensional

images from measurement data. The mathematical characteristics of these polynomials provide effective algorithms for image processing and reconstruction, enhancing diagnostic capacities for neurological illnesses, cancer detection, and surgical planning. Antenna design for telecommunications networks often incorporates Legendre functions in the analysis of radiation patterns. Engineers designing phased array radars, satellite communication antennas, or 5G cellular network equipment enhance directivity and coverage using expansion techniques derived from their specialized roles. Weather prediction methods utilize Legendre polynomial expansions to represent atmospheric variables on spherical domains. Global circulation models that mimic climate trends and forecast extreme weather events utilize spectral approaches employing these functions to effectively resolve the governing equations of atmospheric dynamics.

### **Numerical Techniques for Equations with Variable Coefficients**

Although analytical solutions offer significant theoretical insights, numerous practical applications necessitate numerical methods. The intricacy of variable coefficient equations frequently requires computer techniques for solution generation. Finite difference methods estimate derivatives at discrete locations, converting the differential equation into a system of algebraic equations. These approaches must meticulously manage variable coefficients by assessing them at suitable grid points. Adaptive mesh refinement techniques are especially beneficial when coefficient functions exhibit fast variation in specific areas. Spectral approaches provide solutions as expansions in basis functions, sometimes utilizing orthogonal polynomials such as Legendre or Chebyshev polynomials. For variable coefficient equations, these approaches produce dense matrices while attaining excellent accuracy with a limited number of terms. The finite element technique partitions the domain into elements and estimates the solution with basis functions within each element. This method inherently supports varied coefficients and intricate geometries, rendering it common in engineering applications. Simulations of air flows using computational fluid dynamics utilize numerical techniques to resolve equations with diffusion coefficients that fluctuate with altitude and temperature. Weather forecasting systems and climate models depend on effective variable-coefficient solvers to anticipate atmospheric behavior

across various scales. Semiconductor device simulation entails the use of drift-diffusion equations characterized by spatially variable mobility and diffusion coefficients, which are contingent upon doping profiles and electric fields. Electronics makers employ specialized solvers for these equations for designing transistors, solar cells, and integrated circuits that drive contemporary technology. Biomedical applications encompass the simulation of drug diffusion across heterogeneous tissues characterized by spatially variable diffusion coefficients. Pharmaceutical researchers enhance drug delivery systems and forecast therapeutic success with numerical solutions to variable-coefficient challenges.

### **Asymptotic Techniques for Variable Coefficient Equations**

When parameter values pose difficulties for direct numerical or analytical solutions, asymptotic approaches offer useful approximations. These techniques examine the behavior of equations in limiting scenarios, specifically when a parameter approaches extreme values, either extremely large or very small. The matched asymptotic expansions approach links solutions applicable in distinct locations by aligning them in a shared intermediate overlap zone. This method is especially successful for equations characterized by quickly varying coefficients or boundary layers.

The WKB (Wentzel-Kramers-Brillouin) theory approximates solutions to equations of the following form:

$$\varepsilon^2 y'' + p(t)y = 0$$

Where  $\varepsilon$  represents a minor parameter. This approach produces oscillatory solutions characterized by slowly changing amplitude and rapidly fluctuating phase, suitable for wave propagation issues involving variable medium properties.

Multiscale analysis distinguishes dynamics at several temporal or spatial scales, providing consistently accurate approximations for issues with gradually changing coefficients. Applications of quantum mechanics encompass semiclassical approximations for the Schrödinger equation with slowly fluctuating potentials. These methods link quantum and classical representations of particle motion, crucial for comprehending atomic and molecular

spectroscopy. Optics researchers examining light propagation in gradient-index media utilize WKB algorithms to ascertain ray trajectories and wave characteristics. Optical waveguides, metamaterials, and photonic devices characterized by spatially variable refractive indices get advantages from these asymptotic methodologies. Structural mechanics issues concerning thin shells or beams with varying thickness employ asymptotic approaches to create reduced-order models. Aerospace engineers utilize these estimates to reconcile structural integrity with weight limitations when building lightweight structures for aircraft or spacecraft.

### **Stability Assessment for Systems with Variable Coefficients**

The stability of solutions to differential equations with variable coefficients poses distinct challenges in comparison to systems with fixed coefficients. Lyapunov theory offers methodologies for assessing stability without the necessity of explicitly solving the equations. In linear systems  $\dot{x} = A(t)x$ , where  $A(t)$  is a matrix with variable coefficients, stability is contingent upon the characteristics of the state transition matrix. When  $A(t)$  possesses specific structures, such as periodicity or near-periodicity, Floquet theory provides further insights. Control systems with time-varying parameters necessitate rigorous stability analysis to guarantee performance amid fluctuating situations. Adaptive control techniques that adjust control parameters based on system changes depend on stability criteria for variable coefficient systems. The evaluation of power grid stability entails differential equations with coefficients influenced by generation levels, load demands, and network topology. Engineers engaged in the development of smart grid technology and renewable energy integration methods scrutinize these equations to avert cascade failures and guarantee dependable electricity delivery. Biological systems frequently display time-dependent features as a result of environmental factors or developmental alterations. Population dynamics models, brain networks featuring plastic synapses, and metabolic pathways with regulated enzyme activity all provide variable-coefficient equations, the stability of which dictates system behavior.

### **Applications in Signal Processing and Telecommunications**

Contemporary signal processing heavily depends on linear systems exhibiting time-varying properties. Adaptive filters, which adjust their

coefficients according on the characteristics of the input signal, utilize variable-coefficient difference equations, the discrete counterpart of differential equations. These mathematical frameworks facilitate noise suppression in dynamic settings, channel equalization for wireless communication, and augmentation of biomedical signals for diagnostic applications. Echo cancellation algorithms in teleconferencing systems perpetually adjust filter coefficients to accommodate fluctuating acoustic surroundings. Radar systems that analyze signals from moving targets resolve differential equations in which Doppler effects introduce coefficients that vary with time. Military and civilian radar applications, such as air traffic control and meteorological observation, rely on these mathematical methods to derive target information from received signals. Speech recognition systems represent vocal tract features as time-varying filters, resulting in variable-coefficient equations that encapsulate the dynamics of speech generation. This theoretical framework supports voice assistants, transcription services, and speaker identification technology.

### **Monetary Applications**

Financial mathematics increasingly utilizes variable-coefficient differential equations to represent intricate market dynamics. The Black-Scholes-Merton model for option pricing can be adapted to include time-varying volatility, interest rates, and dividend yields, resulting in variable-coefficient partial differential equations. These advanced models encapsulate market characteristics such as volatility clustering, the term structure of interest rates, and seasonal dividend trends. Financial risk managers employ solutions to these equations for formulating hedging strategies for derivative portfolios in realistic market situations. Term structure models for interest rates frequently use stochastic differential equations with time-varying parameters that represent market expectations and central bank actions. These models facilitate bond valuation, mortgage rate prediction, and monetary policy evaluation. Credit risk assessment employs default intensity models featuring time-varying parameters that mirror fluctuating economic conditions. Banks and financial organizations employ these models for loan pricing, securitization structuring, and capital reserve management.

### **Applications of Biomedical Engineering**

Healthcare technologies increasingly utilize variable-coefficient differential equations to simulate physiological systems with parameters that fluctuate according to patient condition, pharmaceutical effects, or circadian rhythms. Pharmacokinetic-pharmacodynamic (PK-PD) models delineate drug absorption, distribution, metabolism, and excretion through factors contingent upon patient features and physiological conditions. These models inform individualized dosing strategies, pharmaceutical development, and therapeutic enhancement.

Modeling cardiac electrical activity entails reaction-diffusion equations using geographically and temporally variable conductivity tensors that represent the variety of heart tissue and pathological conditions. Cardiologists employ these models to comprehend arrhythmias, refine pacemaker configurations, and formulate therapies for cardiac disorders. Models of brain activity integrate neuronal field equations alongside connection patterns that change over learning, development, or disease advancement. Neuroscientists investigating epilepsy, Alzheimer's disease, or awareness utilize these mathematical frameworks to link observed phenomena with fundamental neuronal principles.

### **Climatology and Ecological Simulation**

Environmental systems inherently encompass characteristics that fluctuate spatially and temporally, rendering variable-coefficient differential equations vital in climate research and ecology. Global climate models resolve equations of atmospheric and oceanic dynamics utilizing coefficients that are contingent upon latitude, height, temperature, and more variables. These intricate models forecast future climatic scenarios, assess human impacts, and examine mitigation measures for climate change. Groundwater movement in heterogeneous aquifers adheres to Darcy's law, characterized by spatially variable hydraulic conductivity. Hydrologists apply answers to these variable-coefficient equations in the design of water delivery systems, the remediation of contaminated areas, and the management of aquifer recharge. Ecosystem models monitor population dynamics and resource flows using factors influenced by seasonal variations, regional variability, and interspecies interactions. Observation biologists and resource managers utilize these models to formulate sustainable harvesting practices, construct protected areas, or forecast the spread of invasive species.



## **Control Systems and Robotics**

Contemporary control theory extensively addresses systems with parameters that vary throughout operation. Gain scheduling approaches develop controllers that adjust to variations in operating points by resolving families of variable-coefficient differential equations. These technologies facilitate flight control systems that ensure stability across varying airspeeds and altitudes, process control systems that adapt to fluctuating feedstock characteristics, and robotic manipulators that manage items of diverse weights or forms. Model predictive control methods consistently resolve variable-coefficient optimization problems to ascertain appropriate control actions amidst fluctuating restrictions and objectives. These sophisticated controllers drive driverless vehicles, optimize industrial processes, and manage energy systems. Robotics applications encompass adaptive motion planning in dynamic situations, wherein robot dynamics and environmental interactions provide variable-coefficient equations. Collaborative robots operating alongside people in industrial, healthcare, or service sectors depend on solutions to these equations for planning safe and efficient movements.

## **Obstacles and Prospective Pathways**

Notwithstanding considerable progress, numerous obstacles persist in the theory and implementation of variable-coefficient differential equations. The pursuit of computational efficiency in high-dimensional systems characterized by rapidly fluctuating coefficients persists in driving algorithm development. Machine learning techniques are progressively combined with conventional numerical methods to address intricate, data-driven coefficient functions. Uncertainty quantification for systems with stochastically variable coefficients constitutes a dynamic field of research. Applications like climate forecasting, financial risk evaluation, and medical treatment strategizing necessitate not only solutions but also confidence metrics for those solutions. Multiscale phenomena with coefficients that vary across disparate scales require specific methods that connect microscopic and macroscopic descriptions. Hierarchical structured materials, biological systems ranging from molecular to organismal sizes, and socioeconomic systems linking individual behaviors to collective dynamics all offer prospects for theoretical advancements. The amalgamation of variable-coefficient differential

equations with data science techniques creates novel opportunities for hybrid modeling methodologies. These strategies integrate theoretical frameworks with empirical data to ascertain coefficient functions, evaluate models, and provide predictions in contexts where solely theoretical or purely data-driven methods would be inadequate.

Linear differential equations with variable coefficients constitute a robust mathematical framework characterized by significant theoretical sophistication and extensive practical applicability. This theory offers systematic methods for modeling intricate, dynamic systems, encompassing core notions of starting value issues and the Wronskian determinant, as well as specialized techniques such as reduction of order and series solutions. The applications encompass nearly all scientific and engineering fields, illustrating the ubiquitous nature of these mathematical constructs. As computer capabilities progress and interdisciplinary borders converge, the significance of these equations in tackling real-world situations increasingly escalates. The development of this discipline demonstrates the collaborative connection between abstract mathematical theory and practical problem-solving. Theoretical insights stimulate novel applications, whereas practical obstacles drive mathematical advancements. This reciprocal process propels advancement in both fields, illustrating the efficacy of mathematical modeling in comprehending and influencing our environment. In a time of unparalleled technological transformation and intricate global issues, proficiency in variable-coefficient differential equations equips researchers, engineers, and policymakers with vital instruments for analysis, forecasting, and design. The ongoing advancement of this mathematical framework is poised to unveil new potentials across various domains, including quantum computing, artificial intelligence, climate modeling, and personalized medicine, thereby reaffirming the enduring significance of mathematical theory in tackling humanity's most urgent challenges.

## **SELF ASSESSMENT QUESTIONS**

### **Multiple Choice Questions (MCQs)**

1. A second-order linear differential equation with variable coefficients has the general form:
  - a)  $y'' + p(x)y' + q(x)y = 0$
  - b)  $y'' + ay' + by = 0$

- c)  $y' + py = q$
  - d) None of the above
2. The Wronskian is used to determine:
    - a) The order of the equation
    - b) The linear dependence or independence of solutions
    - c) The presence of singular points
    - d) None of the above
  3. If the Wronskian of two solutions is nonzero, then the solutions are:
    - a) Linearly dependent
    - b) Linearly independent
    - c) Equal to each other
    - d) None of the above
  4. The reduction of order method is used when:
    - a) One solution is known
    - b) The equation has constant coefficients
    - c) The equation is non-homogeneous
    - d) None of the above
  5. A differential equation is said to have analytic coefficients if:
    - a) The coefficients are differentiable infinitely many times
    - b) The coefficients are constants
    - c) The equation has no singular points
    - d) None of the above
  6. The Legendre equation arises in:
    - a) Quantum mechanics
    - b) Classical mechanics
    - c) Both (a) and (b)
    - d) None of the above
  7. The general solution of a second-order linear differential equation requires:
    - a) Two linearly independent solutions
    - b) A single solution
    - c) Three solutions
    - d) None of the above

8. The variation of parameters method is used to:
- Solve non-homogeneous equations
  - Solve homogeneous equations
  - Compute the Wronskian
  - None of the above
9. A solution to the Legendre equation is given by:
- Legendre polynomials
  - Exponential functions
  - Logarithmic functions
  - None of the above
10. If  $y_1$  is a known solution of a second-order equation, the reduction of order method finds:
- A second linearly independent solution
  - The characteristic equation
  - The Wronskian
  - None of the above

Answer Key:

1	a	3	b	5	a	7	a	9	a
2	b	4	a	6	c	8	a	10	a

## CHECK YOUR PROGRESS

- 1) Write the Legendre differential equation.

.....  
.....  
.....

- 2) What are Legendre polynomials?

.....  
.....  
.....

- 3) Write the first three Legendre polynomials.

.....  
.....  
.....

## LET US SUM UP

When the coefficients of a differential equation are analytic functions, the solution can often be expressed as a power series around a given point. A significant example is the Legendre equation, given by  $(1-x^2)y''-2xy'+n(n+1)y=0$ , whose solutions are the Legendre polynomials  $P_n(x)$ . These equations frequently arise in physics, particularly in potential theory and spherical harmonics.

## **UNIT END EXERCISES**

### **Short Answer Questions:**

1. State the conditions for the existence and uniqueness of solutions.
2. What are singular points, and how do they affect differential equations?
3. Give an application of the Legendre equation in physics.

### **Long Answer Questions:**

1. Discuss the physical and mathematical significance of the Legendre equation.
2. What are singular points in differential equations? Explain their classification.
3. Compare and contrast the methods of variation of parameters and reduction of order.

### **REFERENCES AND SUGGESTED READINGS:**

1. Ince, E. L. (2012). Ordinary Differential Equations. Dover Publications.
2. Nagle, R. K., Saff, E. B., & Snider, A. D. (2018). Fundamentals of Differential Equations (9th ed.). Pearson.
3. Birkhoff, G., & Rota, G. C. (2010). Ordinary Differential Equations (4th ed.). Wiley.
4. Bender, C. M., & Orszag, S. A. (2013). Advanced Mathematical Methods for Scientists and Engineers. Springer.

## BLOCK 4

### UNIT 11

#### LINEAR EQUATION WITH REGULAR SINGULAR POINTS: EULER EQUATION

##### Objectives

- Understand Euler's equation and its role in solving differential equations.
- Learn about second-order equations with regular singular points.
- Study exceptional cases in singular point analysis.
- Explore the Bessel equation and its applications.

##### 11.1 Introduction to Regular Singular Points

When dealing with differential equations, we often encounter singularities - points where the equation behaves in unusual ways. A particularly important class of singularities in the study of differential equations is known as "regular singular points."

Consider a second-order linear differential equation in the standard form:

$$y'' + p(x)y' + q(x)y = 0$$

Where  $p(x)$  and  $q(x)$  are functions of  $x$ . A point  $x_0$  is called a singular point of this equation if either  $p(x)$  or  $q(x)$  is not analytic at  $x_0$  (meaning they have some kind of discontinuity or undefined behavior at that point).

Now, a singular point  $x_0$  is called a regular singular point if the functions  $(x-x_0)p(x)$  and  $(x-x_0)^2q(x)$  are both analytic at  $x_0$ . In other words, when we multiply  $p(x)$  by  $(x-x_0)$  and  $q(x)$  by  $(x-x_0)^2$ , the resulting functions should be well-behaved at  $x_0$ .

To understand this better, we can rewrite our differential equation in a slightly different form:

$$(x-x_0)^2y'' + (x-x_0)p(x)y' + q(x)y = 0$$

If we divide by  $(x-x_0)^2$ , we get:

$$y'' + [p(x)/(x-x_0)]y' + [q(x)/(x-x_0)^2]y = 0$$

For a regular singular point, the functions  $P(x) = (x-x_0)p(x)$  and  $Q(x) = (x-x_0)^2q(x)$  are analytic at  $x_0$ , which means they can be expressed as power series around  $x_0$ . So we can write:

$$P(x) = (x-x_0)p(x) = p_0 + p_1(x-x_0) + p_2(x-x_0)^2 + \dots \quad Q(x) = (x-x_0)^2q(x) = q_0 + q_1(x-x_0) + q_2(x-x_0)^2 + \dots$$

When we substitute these back, our differential equation becomes:

$$y'' + [P(x)/(x-x_0)]y' + [Q(x)/(x-x_0)^2]y = 0$$

or

$$y'' + [(p_0 + p_1(x-x_0) + \dots)/(x-x_0)]y' + [(q_0 + q_1(x-x_0) + \dots)/(x-x_0)^2]y = 0$$

This form is particularly useful for finding solutions around regular singular points.

### Why Regular Singular Points Matter

Regular singular points are important because:

1. They represent a class of singularities for which we can find series solutions using a modified power series approach.
2. Many physical problems lead to differential equations with regular singular points.
3. The behavior of solutions near regular singular points provides important information about the overall solution.

### Example of Identifying Regular Singular Points

Let's examine the equation:

$$x^2y'' + xy' + (x^2 - 1)y = 0$$

We can rewrite this in the standard form:

$$y'' + (1/x)y' + (1 - 1/x^2)y = 0$$

Here,  $p(x) = 1/x$  and  $q(x) = 1 - 1/x^2$

The point  $x = 0$  is a singular point because  $p(x)$  and  $q(x)$  are not analytic at  $x = 0$ .

To check if it's a regular singular point:



- $(x-0)p(x) = x \cdot (1/x) = 1$ , which is analytic at  $x = 0$
- $(x-0)^2q(x) = x^2 \cdot (1 - 1/x^2) = x^2 - 1$ , which is also analytic at  $x = 0$

Therefore,  $x = 0$  is a regular singular point of this differential equation.

In the next section, we'll see how to solve a special class of differential equations with regular singular points known as Euler's equations.

## 11.2 Euler's Equation and Its Solution

Euler's equation is a special type of differential equation with regular singular points. It has the form:

$$x^2y'' + axy' + by = 0$$

where  $a$  and  $b$  are constants. We can also write it in the standard form:

$$y'' + (a/x)y' + (b/x^2)y = 0$$

Notice that  $x = 0$  is a regular singular point because:

- $(x-0)(a/x) = a$ , which is analytic at  $x = 0$
- $(x-0)^2(b/x^2) = b$ , which is also analytic at  $x = 0$

Euler's equation is important because:

1. It represents the simplest type of equation with a regular singular point.
2. Solutions to more complex equations with regular singular points often involve techniques derived from solving Euler's equation.
3. Many physical phenomena are described by Euler-type equations.

### Method of Solution: Substitution Approach

One way to solve Euler's equation is to make the substitution  $x = e^t$ , which transforms the equation into one with constant coefficients.

Let's substitute  $x = e^t$ , which means:

- $y(x) = y(e^t) = Y(t)$
- $dy/dx = (dY/dt) \cdot (dt/dx) = (dY/dt) \cdot (1/x) = e^{-t} \cdot (dY/dt)$

- $$\begin{aligned} \frac{d^2y}{dx^2} &= \frac{d}{dx}\left(\frac{dy}{dx}\right) = \frac{d}{dx}\left(e^{(-t)} \cdot \left(\frac{dY}{dt}\right)\right) = e^{(-t)} \cdot \frac{d}{dx}\left(\frac{dY}{dt}\right) - e^{(-t)} \cdot \left(\frac{dY}{dt}\right) \cdot \left(\frac{1}{x}\right) \\ &= e^{(-t)} \cdot \left(\frac{d^2Y}{dt^2}\right) \cdot \left(\frac{1}{x}\right) - e^{(-2t)} \cdot \left(\frac{dY}{dt}\right) = e^{(-2t)} \cdot \left[\frac{d^2Y}{dt^2} - \frac{dY}{dt}\right] \end{aligned}$$

Substituting these into the Euler equation  $x^2y'' + axy' + by = 0$ :

$$x^2 \cdot e^{(-2t)} \cdot \left[\frac{d^2Y}{dt^2} - \frac{dY}{dt}\right] + ax \cdot e^{(-t)} \cdot \left(\frac{dY}{dt}\right) + b \cdot Y = 0$$

$$\text{Simplifying: } e^{(2t)} \cdot e^{(-2t)} \cdot \left[\frac{d^2Y}{dt^2} - \frac{dY}{dt}\right] + a \cdot e^t \cdot e^{(-t)} \cdot \left(\frac{dY}{dt}\right) + b \cdot Y = 0$$

$$\text{Which gives us: } \frac{d^2Y}{dt^2} - \frac{dY}{dt} + a \cdot \left(\frac{dY}{dt}\right) + b \cdot Y = 0$$

$$\text{Rearranging: } \frac{d^2Y}{dt^2} + (a-1) \cdot \left(\frac{dY}{dt}\right) + b \cdot Y = 0$$

This is a second-order linear differential equation with constant coefficients, which we know how to solve.

### Method of Solution: Power Series Approach

Another approach is to try a solution of the form  $y = x^r$ , where  $r$  is a constant to be determined.

If  $y = x^r$ , then:

- $$y' = rx^{(r-1)}$$
- $$y'' = r(r-1)x^{(r-2)}$$

$$\text{Substituting into the Euler equation: } x^2 \cdot r(r-1)x^{(r-2)} + ax \cdot rx^{(r-1)} + b \cdot x^r = 0$$

$$\text{Simplifying: } r(r-1)x^r + ar \cdot x^r + b \cdot x^r = 0$$

$$\text{Factoring out } x^r: x^r[r(r-1) + ar + b] = 0$$

$$\text{Since } x^r \text{ is not identically zero for } x \neq 0, \text{ we must have: } r(r-1) + ar + b = 0$$

This is called the indicial equation or characteristic equation for Euler's equation. Rearranging:  $r^2 + (a-1)r + b = 0$

This is a quadratic equation in  $r$  that we can solve to find the possible values of  $r$ .

### Cases for Solutions to Euler's Equation

The nature of the solutions depends on the roots of the indicial equation  $r^2 + (a-1)r + b = 0$ :

Case 1: Two Distinct Real Roots ( $r_1$  and  $r_2$ )

If the indicial equation has two distinct real roots  $r_1$  and  $r_2$ , then the general solution to the Euler equation is:

$$y(x) = c_1 x^{(r_1)} + c_2 x^{(r_2)}$$

where  $c_1$  and  $c_2$  are arbitrary constants.

Case 2: Repeated Real Root ( $r_1 = r_2 = r$ )

If the indicial equation has a repeated root  $r$ , then the general solution is:

$$y(x) = c_1 x^r + c_2 x^r \cdot \ln(x)$$

Case 3: Complex Conjugate Roots ( $r_1 = \alpha + i\beta$ ,  $r_2 = \alpha - i\beta$ )

If the indicial equation has complex conjugate roots  $\alpha \pm i\beta$ , the general solution can be written as:

$$y(x) = x^\alpha [c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x))]$$

Example: Solving an Euler Equation

Let's solve the equation:  $x^2 y'' - 3xy' + 4y = 0$

Step 1: Identify that this is an Euler equation with  $a = -3$  and  $b = 4$ .

Step 2: Form the indicial equation:  $r^2 + (a-1)r + b = 0$   $r^2 + (-3-1)r + 4 = 0$   $r^2 - 4r + 4 = 0$   $(r - 2)^2 = 0$

Step 3: Since we have a repeated root  $r = 2$ , the general solution is:  $y(x) = c_1 x^2 + c_2 x^2 \ln(x)$

This gives us the complete solution to the Euler equation.

## CHECK YOUR PROGRESS

- 1) Write the standard form of Euler's equation.

.....  
.....  
.....

- 2) What substitution converts Euler's equation into one with constant coefficients?

.....  
.....  
.....

- 3) Define the indicial equation.

.....  
.....  
.....

## LET US SUM UP

The Euler or Cauchy–Euler equation is a special type of differential equation where the coefficients are powers of  $x$ , such as  $x^2y''+axy'+by=0$ . It can be reduced to a constant coefficient equation by substituting  $y=x^m$ , which leads to an algebraic (indicial) equation in  $m$ . This type of equation frequently appears in problems involving scaling or power-law behavior.

## UNIT END EXERCISES

### Short Answer Questions:

1. What is the significance of the indicial equation in the Frobenius method?
2. How does the Bessel equation arise in physics?
3. Give an example of an equation with a regular singular point.

### Long Answer Questions:

1. Solve the Bessel equation  $x^2y'' + xy' + (x^2 - 1)y = 0$  using series expansion.
2. Discuss the physical applications of Bessel functions in engineering and physics.
3. Explain exceptional cases in the Frobenius method with examples.

### REFERENCES AND SUGGESTED READINGS:

1. Böcher, M. (2015). Regular Points of Linear Differential Equations of the Second Order, American Mathematical Society.
2. Watson, G. N. (2020). A Treatise on the Theory of Bessel Functions. Cambridge University Press.
3. Stakgold, I., & Holst, M. J. (2019), Green's Functions and Boundary Value Problems (3rd ed.). Wiley.

## UNIT 12

### SECOND ORDER EQUATIONS WITH REGULAR SINGULAR POINTS

#### 12.1 Second-Order Equations with Regular Singular Points

Now that we understand Euler's equation, we can tackle more general second-order differential equations with regular singular points.

#### Series Solutions around Regular Singular Points

Consider a general second-order differential equation with a regular singular point at  $x = 0$ :

$$x^2 y'' + x p(x) y' + q(x) y = 0$$

Where  $p(x)$  and  $q(x)$  are analytic at  $x = 0$  and can be expressed as power series:

$$p(x) = p_0 + p_1 x + p_2 x^2 + \dots \quad q(x) = q_0 + q_1 x + q_2 x^2 + \dots$$

To find a solution, we try a modified power series of the form:

$$y(x) = x^r \sum_{n=0}^{\infty} a_n x^n = x^r (a_0 + a_1 x + a_2 x^2 + \dots)$$

where  $r$  is a constant to be determined and  $a_0 \neq 0$ .

The method of finding solutions involves:

1. Substituting the series into the differential equation.
2. Finding the indicial equation to determine possible values of  $r$ .
3. Determining the recurrence relation for the coefficients  $a_n$ .
4. Constructing the solutions based on the nature of the roots of the indicial equation.

Let's work through this process:

#### Step 1: Derive the Indicial Equation

When we substitute the series solution into the differential equation and collect the lowest power terms (which will involve  $x^r$ ), we get what's called the indicial equation:

$$r(r-1) + p_0 r + q_0 = 0$$

This is a quadratic equation in  $r$ , and its roots determine the form of our solutions.

### Step 2: Analyze the Roots of the Indicial Equation

Let's denote the roots of the indicial equation as  $r_1$  and  $r_2$ , with  $r_1 \geq r_2$ .

There are three possible cases:

1. The roots differ by a non-integer:  $r_1 - r_2 \neq \text{integer}$
2. The roots are equal:  $r_1 = r_2$
3. The roots differ by a positive integer:  $r_1 - r_2 = \text{positive integer}$

### Step 3: Construct the Solutions Based on the Roots

Case 1: Roots Differ by a Non-Integer

If  $r_1 - r_2$  is not an integer, we obtain two linearly independent solutions:

$$y_1(x) = x^{(r_1)} \sum_{n=0}^{\infty} a_n x^n \quad y_2(x) = x^{(r_2)} \sum_{n=0}^{\infty} b_n x^n$$

where  $a_0 \neq 0$  and  $b_0 \neq 0$ .

Case 2: Equal Roots

If  $r_1 = r_2 = r$ , then we get:

$$y_1(x) = x^r \sum_{n=0}^{\infty} a_n x^n \quad y_2(x) = y_1(x) \ln(x) + x^r \sum_{n=1}^{\infty} b_n x^n$$

Case 3: Roots Differ by a Positive Integer

If  $r_1 - r_2 = m$  (a positive integer), then:

$$y_1(x) = x^{(r_1)} \sum_{n=0}^{\infty} a_n x^n \quad y_2(x) = C y_1(x) \ln(x) + x^{(r_2)} \sum_{n=0}^{\infty} b_n x^n$$

where  $C$  may be zero or non-zero depending on the specific equation.

### Method of Frobenius

The procedure we've outlined is known as the Method of Frobenius. It provides a systematic way to find series solutions around regular singular points.

Here's a step-by-step approach:

1. Identify a regular singular point  $x_0$ .

2. Shift the equation to make  $x_0 = 0$  (if necessary) by substituting  $x \rightarrow x + x_0$ .
3. Try a solution of the form  $y = x^r \sum_{n=0}^{\infty} a_n x^n$  with  $a_0 \neq 0$ .
4. Substitute into the differential equation and collect terms with the same power of  $x$ .
5. From the lowest power terms, derive the indicial equation.
6. Based on the roots of the indicial equation, determine the form of the solutions.
7. Find the recurrence relation for the coefficients  $a_n$  and solve for them.
8. Construct the general solution.

Example: Applying the Method of Frobenius

Let's solve the equation:  $x^2 y'' + x(1-x)y' - (1+x)y = 0$

Step 1: Verify that  $x = 0$  is a regular singular point.  $p(x) = (1-x)$ , so  $x p(x) = x(1-x)$  is analytic at  $x = 0$ .  $q(x) = -(1+x)$ , so  $x^2 q(x) = -x^2(1+x)$  is analytic at  $x = 0$ .

Step 2: Try a solution of the form  $y = x^r \sum_{n=0}^{\infty} a_n x^n$ .

Step 3: Derive the indicial equation. For the lowest power terms ( $x^r$ ), we get:  
 $r(r-1) + r - 1 = 0$ ,  $r^2 - r + r - 1 = 0$ ,  $r^2 - 1 = 0$ ,  $(r+1)(r-1) = 0$

So the roots are  $r_1 = 1$  and  $r_2 = -1$ .

Step 4: Since  $r_1 - r_2 = 2$  (a positive integer), we use Case 3. The first solution is:  $y_1(x) = x^1(a_0 + a_1 x + a_2 x^2 + \dots)$

Step 5: Substitute back and find the recurrence relation for  $a_n$  to complete the solution.

### Solved and Unsolved Problems

#### Solved Problem 1: Identify Regular Singular Points

Find all singular points of the differential equation and determine which ones are regular singular points:

$$x(x-2)y'' + (x+1)y' - 3y = 0$$



Solution: First, let's rewrite the equation in standard form:

$$y'' + [(x+1)/(x(x-2))]y' - [3/(x(x-2))]y = 0$$

Here  $p(x) = (x+1)/(x(x-2))$  and  $q(x) = -3/(x(x-2))$

The singular points occur when the coefficient of  $y''$  is zero, which happens when  $x = 0$  or  $x = 2$ .

For  $x = 0$ :

- $(x-0)p(x) = x \cdot (x+1)/(x(x-2)) = (x+1)/(x-2)$ , which has a finite limit as  $x \rightarrow 0$
- $(x-0)^2q(x) = x^2 \cdot (-3)/(x(x-2)) = -3x/(x-2)$ , which has a finite limit as  $x \rightarrow 0$

Therefore,  $x = 0$  is a regular singular point.

For  $x = 2$ :

- $(x-2)p(x) = (x-2) \cdot (x+1)/(x(x-2)) = (x+1)/x$ , which has a finite limit as  $x \rightarrow 2$
- $(x-2)^2q(x) = (x-2)^2 \cdot (-3)/(x(x-2)) = -3(x-2)/x$ , which has a finite limit as  $x \rightarrow 2$

Therefore,  $x = 2$  is also a regular singular point.

### Solved Problem 2: Solve an Euler Equation

Solve the Euler equation:  $x^2y'' + 5xy' + 4y = 0$

Solution: This is an Euler equation with  $a = 5$  and  $b = 4$ .

The indicial equation is:  $r^2 + (a-1)r + b = 0$   $r^2 + (5-1)r + 4 = 0$   $r^2 + 4r + 4 = 0$   
 $(r + 2)^2 = 0$

We have a repeated root  $r = -2$ .

For a repeated root, the general solution is:  $y(x) = c_1x^{(-2)} + c_2x^{(-2)}\ln(x)$

### Solved Problem 3: Find Recurrence Relation

For the differential equation  $x^2y'' + xy' + (x^2 - 1)y = 0$ , find the recurrence relation for the coefficients in the series solution around  $x = 0$ .

Solution: Let's try a solution of the form  $y = x^r \sum_{n=0}^{\infty} a_n x^n$ .

Step 1: Find the indicial equation. The equation can be written as:  $y'' + (1/x)y' + (1 - 1/x^2)y = 0$

The indicial equation is:  $r(r-1) + r + (-1) = 0 \Rightarrow r^2 = 1 \Rightarrow r = \pm 1$

So the roots are  $r_1 = 1$  and  $r_2 = -1$ .

Step 2: Let's find the recurrence relation for the first solution with  $r = 1$ . Substituting  $y = x^1 \sum_{n=0}^{\infty} a_n x^n$  into the original equation and collecting terms with the same power of  $x$ , we get:

$$\sum_{n=0}^{\infty} [(n+1)(n+2)a_{n+2} + a_n]x^{n+1} = 0$$

For this to be zero for all  $x$ , each coefficient must be zero:  $(n+1)(n+2)a_{n+2} + a_n = 0$

$$\text{Solving for } a_{n+2}: a_{n+2} = -a_n / [(n+1)(n+2)]$$

This is our recurrence relation.

#### Solved Problem 4: Find Series Solution

Find the first four terms of the series solution to the differential equation:

$$xy'' - y' + 4x^3y = 0$$

with the initial condition  $y(0) = 1, y'(0) = 2$ .

Solution: First, let's rewrite the equation in standard form:  $y'' - (1/x)y' + 4x^2y = 0$

This has a regular singular point at  $x = 0$ .

Let's try a solution of the form  $y = \sum_{n=0}^{\infty} a_n x^n$ . We need to find  $a_0, a_1, a_2$ , and  $a_3$ .

$$\text{Substituting into the equation: } \sum_{n=0}^{\infty} n(n-1)a_n x^{n-2} - \sum_{n=0}^{\infty} n a_n x^{n-1} + 4 \sum_{n=0}^{\infty} a_n x^{n+2} = 0$$

$$\text{Shifting indices to match powers of } x: \sum_{n=2}^{\infty} (n+2)(n+1)a_{n+2} x^n - \sum_{n=1}^{\infty} (n+1)a_{n+1} x^n + 4 \sum_{n=0}^{\infty} a_n x^{n+2} = 0$$

Collecting terms for each power of  $x$ : For  $n = 0$ :  $2 \cdot 1 \cdot a_2 - 1 \cdot a_1 = 0 \Rightarrow a_2 = a_1/2$

For  $n = 1$ :  $3 \cdot 2 \cdot a_3 - 2 \cdot a_2 = 0 \Rightarrow a_3 = 2a_2/6 = a_1/6$

From the initial conditions:  $y(0) = a_0 = 1, y'(0) = a_1 = 2$

Therefore:  $a_0 = 1, a_1 = 2, a_2 = a_1/2 = 2/2 = 1, a_3 = a_1/6 = 2/6 = 1/3$

The first four terms of the series solution are:  $y(x) = 1 + 2x + x^2 + (1/3)x^3 + \dots$

### Solved Problem 5: Find General Solution

Find the general solution to the differential equation:  $x^2y'' - x(x+2)y' + (x+2)y = 0$

Solution: Let's verify that  $x = 0$  is a regular singular point and find the solutions around this point.

Rewriting in standard form:  $y'' - [(x+2)/x]y' + [(x+2)/x^2]y = 0$

For  $x = 0$ :

- $x \cdot (-(x+2)/x) = -(x+2)$ , which is analytic at  $x = 0$
- $x^2 \cdot ((x+2)/x^2) = x+2$ , which is analytic at  $x = 0$

So  $x = 0$  is a regular singular point.

Let's try a solution of the form  $y = x^r \sum_{n=0}^{\infty} a_n x^n$ .

The indicial equation is:  $r(r-1) - 2r + 2 = 0$   $r^2 - r - 2r + 2 = 0$   $r^2 - 3r + 2 = 0$   $(r-1)(r-2) = 0$

So the roots are  $r_1 = 2$  and  $r_2 = 1$ .

Since  $r_1 - r_2 = 1$  (a positive integer), we have:  $y_1(x) = x^2(a_0 + a_1x + a_2x^2 + \dots)$   
 $y_2(x) = Cy_1(x)\ln(x) + x(b_0 + b_1x + b_2x^2 + \dots)$

For this particular equation, further calculation shows that  $C = 0$ , so the general solution is:  $y(x) = c_1x^2(a_0 + a_1x + a_2x^2 + \dots) + c_2x(b_0 + b_1x + b_2x^2 + \dots)$

### Unsolved Problem 1

Determine if  $x = 0$  is a regular singular point for the differential equation:  $x^3y'' + x^2y' - 2y = 0$

If it is, find the indicial equation and its roots.

### Unsolved Problem 2

Solve the Euler equation:  $x^2y'' - 3xy' - 3y = 0$

### Unsolved Problem 3

Find the general solution to the differential equation:  $x^2y'' + 3xy' + (x^2 - 1)y = 0$

#### Unsolved Problem 4

Derive the recurrence relation for the coefficients in the series solution to:  
 $x^2y'' + xy' + (x - 1)y = 0$   
around the regular singular point  $x = 0$ .

#### Unsolved Problem 5

For the differential equation:  $x^2y'' - x(2-x)y' + 2(1-x)y = 0$

Determine all singular points and classify them as regular or irregular. Then find the general solution around  $x = 0$ .

In this comprehensive exploration of differential equations with regular singular points, we have:

1. Defined and characterized regular singular points in second-order linear differential equations
2. Studied Euler's equations as the simplest type of equations with regular singular points
3. Learned multiple methods for solving Euler's equations
4. Developed the Method of Frobenius for finding series solutions around regular singular points
5. Analyzed different cases based on the roots of the indicial equation
6. Worked through several solved examples to illustrate the techniques
7. Provided challenging unsolved problems for practice

The theory of differential equations with regular singular points has numerous applications in physics, engineering, and other sciences. The methods we've developed, particularly the Method of Frobenius, provide powerful tools for solving these equations and understanding the behavior of their solutions near singular points.

Regular singular points represent a special case where, despite the presence of a singularity, we can still find well-behaved series solutions. This distinguishes them from irregular singular points, which require different

and often more complex approaches. By mastering the techniques presented here, you'll be equipped to handle a wide range of differential equations that arise in various applications.

## 12.2 Frobenius Method for Solving Singular Equations

The Frobenius method is a powerful technique for solving linear ordinary differential equations with regular singular points. Unlike the power series method which works for ordinary points, the Frobenius method allows us to find solutions near singular points where the standard approach fails.

### Introduction to Regular Singular Points

A second-order linear differential equation in standard form is written as:

$$y'' + P(x)y' + Q(x)y = 0$$

A point  $x = x_0$  is called a regular singular point if both  $(x - x_0)P(x)$  and  $(x - x_0)^2Q(x)$  are analytic at  $x = x_0$ . This means that while  $P(x)$  and  $Q(x)$  may have poles at  $x_0$ , these poles are of limited order (at most 1 for  $P$  and at most 2 for  $Q$ ).

When we encounter a regular singular point, the standard power series method fails. However, the Frobenius method allows us to find solutions by assuming a modified form of the solution.

### The Frobenius Method Approach

The key insight of the Frobenius method is to look for solutions of the form:

$$y(x) = (x - x_0)^r \sum a_n (x - x_0)^n$$

where  $r$  is a constant that we need to determine, and  $\{a_n\}$  are coefficients to be found. Without loss of generality, we can assume  $a_0 \neq 0$ .

For simplicity, we'll often take  $x_0 = 0$ , which means we're looking for solutions of the form:

$$y(x) = x^r \sum a_n x^n = x^r (a_0 + a_1 x + a_2 x^2 + \dots)$$

The Frobenius method consists of the following steps:

1. Verify that  $x = x_0$  is indeed a regular singular point
2. Express  $P(x)$  and  $Q(x)$  as Laurent series around  $x_0$

3. Substitute the assumed form of the solution into the differential equation
4. Find the indicial equation to determine possible values of  $r$
5. For each value of  $r$ , find the recurrence relation for the coefficients  $a_n$
6. Construct the solutions

### Finding the Indicial Equation

When we substitute our assumed solution form into the differential equation and collect terms with the smallest power of  $x$ , we get the indicial equation. This is typically a quadratic equation in  $r$  that determines the possible values for  $r$ .

If  $P(x) = p_1/(x - x_0) + p_0 + p_1(x - x_0) + \dots$  and  $Q(x) = q_2/(x - x_0)^2 + q_1/(x - x_0) + q_0 + \dots$

Then the indicial equation is:

$$r(r-1) + p_1r + q_2 = 0$$

This is also often written as:

$$r^2 + (p_1-1)r + q_2 = 0$$

The roots of this equation,  $r_1$  and  $r_2$ , are critical for determining the nature of the solutions.

### Cases Based on Indicial Equation Roots

1. Case 1:  $r_1$  and  $r_2$  are distinct and don't differ by an integer
  - Two linearly independent solutions exist:  $y_1(x) = x^{r_1}(a_0 + a_1x + a_2x^2 + \dots)$   $y_2(x) = x^{r_2}(b_0 + b_1x + b_2x^2 + \dots)$
2. Case 2:  $r_1$  and  $r_2$  are equal ( $r_1 = r_2 = r$ )
  - The first solution is:  $y_1(x) = x^r(a_0 + a_1x + a_2x^2 + \dots)$
  - The second solution involves a logarithmic term:  $y_2(x) = y_1(x)\ln(x) + x^r(c_1x + c_2x^2 + \dots)$
3. Case 3:  $r_1$  and  $r_2$  differ by an integer ( $r_1 - r_2 = N$ , where  $N$  is a positive integer)

- The solution corresponding to the larger root  $r_1$  is:  $y_1(x) = x^{r_1}(a_0 + a_1x + a_2x^2 + \dots)$
- The second solution may or may not involve a logarithmic term, depending on certain conditions

### The Recurrence Relation

After finding  $r$ , we substitute our assumed solution into the differential equation and collect coefficients of each power of  $x$ . This gives us a recurrence relation for the coefficients  $a_n$ .

The general form of the recurrence relation is complex and depends on the specific equation, but it allows us to compute  $a_1, a_2, a_3$ , etc. in terms of  $a_0$  (which we typically set to 1).

### Example: Bessel's Equation

Bessel's equation is a classic example where the Frobenius method is applied:

$$x^2y'' + xy' + (x^2 - n^2)y = 0$$

This equation has a regular singular point at  $x = 0$ . The indicial equation is  $r^2 - n^2 = 0$ , giving  $r = \pm n$ .

The resulting solutions are the Bessel functions of the first and second kind,  $J_n(x)$  and  $Y_n(x)$ .

### Worked Examples

Let's apply the Frobenius method to several examples:

#### Example 1: Euler's Equation

Consider the Euler equation:

$$x^2y'' + 3xy' - y = 0$$

Step 1: Verify that  $x = 0$  is a regular singular point.  $P(x) = 3/x$ , so  $xP(x) = 3$  is analytic at  $x = 0$ .  $Q(x) = -1/x^2$ , so  $x^2Q(x) = -1$  is analytic at  $x = 0$ . Therefore,  $x = 0$  is a regular singular point.

Step 2: Find the indicial equation. For Euler's equation, the indicial equation is:  $r(r-1) + 3r - 1 = 0$   $r^2 + 2r - 1 = 0$

Step 3: Solve the indicial equation. Using the quadratic formula:  $r = (-2 \pm \sqrt{(4+4)})/2 = -1 \pm \sqrt{2}$

So  $r_1 = -1 + \sqrt{2} \approx 0.414$  and  $r_2 = -1 - \sqrt{2} \approx -2.414$

Step 4: Since  $r_1$  and  $r_2$  differ by 2.828, which is not an integer, we can find two linearly independent solutions.

Step 5: For Euler's equation, the solutions can be written directly:  $y_1(x) = x^{(-1+\sqrt{2})}$   
 $y_2(x) = x^{(-1-\sqrt{2})}$

The general solution is:  $y(x) = C_1 x^{(-1+\sqrt{2})} + C_2 x^{(-1-\sqrt{2})}$

### Example 2: Legendre's Equation

Consider Legendre's equation:

$$(1-x^2)y'' - 2xy' + n(n+1)y = 0$$

To apply the Frobenius method, we need to transform this equation to have a singular point at  $x = 0$ . Let's focus instead on the singular points at  $x = \pm 1$ .

For  $x = 1$ , we make the substitution  $t = x-1$ :

$$\text{The equation becomes: } t(2-t)y'' + (2-2t)y' + n(n+1)y = 0$$

Step 1: Verify that  $t = 0$  is a regular singular point.  $P(t) = (2-2t)/(t(2-t)) = 2/(t(2-t))$ , so  $tP(t) = 2/(2-t)$  is analytic at  $t = 0$ .  $Q(t) = n(n+1)/(t(2-t))$ , so  $t^2Q(t) = tn(n+1)/(2-t)$  is analytic at  $t = 0$ . Therefore,  $t = 0$  is a regular singular point.

Step 2: Find the indicial equation. The indicial equation is:  $r(r-1) + r - 0 = 0$   
 $r^2 = 0$

Step 3: Solve the indicial equation. We have  $r_1 = r_2 = 0$  (repeated root).

Step 4: Since we have equal roots, one solution will involve a logarithmic term.

Step 5: The first solution is:  $y_1(t) = \sum a_n t^n = a_0 + a_1 t + a_2 t^2 + \dots$

We can find the recurrence relation by substituting this into the original equation and collect coefficients of each power of  $t$ .

The second solution, due to the repeated root, will have the form:  $y_2(t) = y_1(t)\ln(t) + t^0(c_1 t + c_2 t^2 + \dots)$



### Example 3: Bessel's Equation of Order 0

Consider Bessel's equation of order 0:

$$x^2 y'' + xy' + x^2 y = 0$$

Step 1: Verify that  $x = 0$  is a regular singular point.  $P(x) = 1/x$ , so  $xP(x) = 1$  is analytic at  $x = 0$ .  $Q(x) = x^2/x^2 = 1$ , so  $x^2Q(x) = x^2$  is analytic at  $x = 0$ . Therefore,  $x = 0$  is a regular singular point.

Step 2: Find the indicial equation. The indicial equation is:  $r(r-1) + r + 0 = 0$   
 $r^2 = 0$

Step 3: Solve the indicial equation. We have  $r_1 = r_2 = 0$  (repeated root).

Step 4: Since we have equal roots, one solution will involve a logarithmic term.

Step 5: Let's find the first solution:  $y(x) = x^0(a_0 + a_1x + a_2x^2 + \dots)$

Substituting into the original equation:  $x^2 y'' + xy' + x^2 y = 0$

After collecting terms and equating coefficients, we get: For  $n \geq 2$ :  $a_n = -a_{n-2} / (n^2)$

This gives:  $a_2 = -a_0/4$   $a_4 = -a_2/16 = a_0/64$   $a_6 = -a_4/36 = -a_0/2304 \dots$

And all odd coefficients  $a_1, a_3, a_5, \dots$  are 0.

Setting  $a_0 = 1$ , we get:  $y_1(x) = 1 - x^2/4 + x^4/64 - x^6/2304 + \dots$

This is the Bessel function of the first kind,  $J_0(x)$ .

The second solution involves a logarithmic term and gives the Bessel function of the second kind,  $Y_0(x)$ .

### Example 4: Airy's Equation

Consider Airy's equation:

$$y'' - xy = 0$$

This equation does not have a regular singular point at  $x = 0$ , but rather at infinity. However, we can apply a transformation to study it with the Frobenius method.

If we make the substitution  $t = x(3/2)$ , the equation transforms to have a regular singular point at  $t = 0$ .

The transformed equation is:  $y'' + (1/4t^2)y = 0$

Step 1: Verify that  $t = 0$  is a regular singular point.  $P(t) = 0$ , so  $tP(t) = 0$  is analytic at  $t = 0$ .  $Q(t) = 1/(4t^2)$ , so  $t^2Q(t) = 1/4$  is analytic at  $t = 0$ . Therefore,  $t = 0$  is a regular singular point.

Step 2: Find the indicial equation. The indicial equation is:  $r(r-1) + 0 + 1/4 = 0$   
 $r^2 - r + 1/4 = 0 \quad (r - 1/2)^2 = 0$

Step 3: Solve the indicial equation. We have  $r_1 = r_2 = 1/2$  (repeated root).

Step 4: Since we have equal roots, one solution will involve a logarithmic term.

Step 5: The solutions in terms of  $t$  are complex, but transforming back to  $x$ , we get the Airy functions  $Ai(x)$  and  $Bi(x)$  as the solutions to the original equation.

### Example 5: Hypergeometric Equation

Consider the hypergeometric equation:

$$x(1-x)y'' + [c - (a+b+1)x]y' - aby = 0$$

Step 1: Verify that  $x = 0$  is a regular singular point.  $P(x) = [c - (a+b+1)x]/(x(1-x))$ , so  $xP(x) = [c - (a+b+1)x]/(1-x)$  is analytic at  $x = 0$ .  $Q(x) = -ab/(x(1-x))$ , so  $x^2Q(x) = -abx/(1-x)$  is analytic at  $x = 0$ . Therefore,  $x = 0$  is a regular singular point.

Step 2: Find the indicial equation. The indicial equation is:  $r(r-1) + cr - 0 = 0$   
 $r^2 + (c-1)r = 0 \quad r(r + c - 1) = 0$

Step 3: Solve the indicial equation. We have  $r_1 = 0$  and  $r_2 = 1-c$ .

Step 4: The nature of the solutions depends on whether  $c$  is an integer.

Step 5: For  $r_1 = 0$ , the solution is:  $y_1(x) = 1 + (ab/c)x + [a(a+1)b(b+1)/(c(c+1)2!)]x^2 + \dots$

This is the hypergeometric function  ${}_2F_1(a, b; c; x)$ .

For  $r_2 = 1-c$ , if  $c$  is not an integer, the second solution is:  $y_2(x) = x^{(1-c)} [1 + \dots]$

If  $c$  is an integer, the second solution may involve a logarithmic term.

## Unsolved Problems

Here are five unsolved problems to practice applying the Frobenius method:

### Problem 1:

Solve the differential equation:  $2x^2y'' + 3xy' - y = 0$

### Problem 2:

Find the general solution to:  $x^2y'' + x(1-x)y' + y = 0$

### Problem 3:

Determine the nature of solutions to:  $x^2y'' + xy' + (x^2 - 1/4)y = 0$

### Problem 4:

Solve using the Frobenius method:  $x^2y'' - x(x+2)y' + (x+2)y = 0$

### Problem 5:

Find the first few terms of both solutions to:  $x^2y'' + xy' - (1+x)y = 0$

## 12.3 Exceptional Cases in Regular Singular Points

When applying the Frobenius method, there are certain exceptional cases that require special attention. These cases arise when the roots of the indicial equation satisfy specific conditions.

### Roots Differing by an Integer

If the roots of the indicial equation,  $r_1$  and  $r_2$ , differ by a positive integer  $N$  (where  $r_1 > r_2$  and  $r_1 - r_2 = N$ ), we have an exceptional case. In this scenario, the standard approach might fail to produce two linearly independent solutions.

For the larger root  $r_1$ , we can always find a solution of the form:

$$y_1(x) = x^{r_1} \sum a_n x^n = x^{r_1} (a_0 + a_1 x + a_2 x^2 + \dots)$$

However, for the smaller root  $r_2$ , the recurrence relation may break down when attempting to find the coefficient  $a_{-N}$ . This happens because the term corresponding to  $a_{-N}$  in the recurrence relation has a coefficient of zero.

The Logarithmic Case

When  $r_1 - r_2 = N$  (a positive integer), there are two possibilities for the second solution:

1. Case A: If a certain condition is met, the second solution has the form:  $y_2(x) = Cy_1(x)\ln(x) + x^{r_2} \sum b_n x^n$

where  $C$  is a constant that may be zero.

2. Case B: If the condition is not met, the second solution has the form:  $y_2(x) = x^{r_2} \sum b_n x^n$

The condition that determines whether a logarithmic term appears depends on the specific differential equation and involves the coefficient of  $a_N$  in the first solution.

### Equal Roots

When  $r_1 = r_2$  (the indicial equation has a repeated root), the logarithmic term always appears in the second solution:

$$y_2(x) = y_1(x)\ln(x) + x^r \sum b_n x^n$$

This is a special case of the scenario where the roots differ by an integer (with  $N = 0$ ).

### Detecting the Need for a Logarithmic Term

To determine whether a logarithmic term is needed, we follow these steps:

1. Find the first solution  $y_1(x)$  using the larger root  $r_1$
2. Try to find a second solution of the form  $y_2(x) = x^{r_2} \sum b_n x^n$
3. If we encounter a contradiction in the recurrence relation (typically at the  $N$ -th term), then a logarithmic term is necessary

The specific criterion can be expressed mathematically. If we have the recurrence relation for the coefficients in the form:

$$(n + r_2)(n + r_2 - 1 + p_1) a_n + \text{terms involving } a_0, a_1, \dots, a_{n-1} = 0$$

Then when  $n = N = r_1 - r_2$ , the first term becomes zero because  $(N + r_2) = r_1$ , and the indicial equation says that  $r_1(r_1 - 1 + p_1) + q_2 = 0$ .

At this point, we need to check whether the remaining terms add up to zero naturally. If they don't, we need to introduce a logarithmic term.

## Method of Frobenius for Logarithmic Solutions

When a logarithmic term is needed, we use the method of Frobenius to find the second solution:

1. Assume a solution of the form:  $y_2(x) = y_1(x)\ln(x) + \sum b_n x^{(n+r_2)}$
2. Substitute this into the differential equation and collect terms
3. Use the fact that  $y_1(x)$  is already a solution to simplify the resulting equation
4. Determine the coefficients  $b_n$  from the remaining terms

This approach ensures that we find two linearly independent solutions in all cases.

### Examples of Exceptional Cases

Let's examine some examples to illustrate these exceptional cases:

#### Example 1: Equal Roots

Consider the equation:  $x^2 y'' + xy' - x^2 y = 0$

The indicial equation is:  $r(r-1) + r - 0 = 0 \Rightarrow r^2 = 0$

This gives  $r_1 = r_2 = 0$  (equal roots).

The first solution has the form:  $y_1(x) = a_0 + a_1 x + a_2 x^2 + \dots$

Substituting into the original equation and collecting terms, we get: For  $n \geq 2$ :  $n^2 a_n - a_{n-2} = 0$  Thus,  $a_n = a_{n-2}/n^2$

With  $a_0 = 1$ , we get:  $a_2 = 1/4 \quad a_4 = a_2/16 = 1/64 \quad a_6 = a_4/36 = 1/2304 \dots$

And  $a_1 = a_3 = a_5 = \dots = 0$

So the first solution is:  $y_1(x) = 1 + x^2/4 + x^4/64 + x^6/2304 + \dots$

The second solution must include a logarithmic term:  $y_2(x) = y_1(x)\ln(x) + b_1 x + b_2 x^2 + \dots$

Substituting this into the differential equation and solving for the coefficients  $b_n$ , we would find the complete second solution.

#### Example 2: Roots Differing by an Integer

Consider the equation:  $x^2 y'' + x(1+x)y' + y = 0$

The indicial equation is:  $r(r-1) + r(1) + 0 = 0 \Rightarrow r^2 = 0$

This gives  $r_1 = r_2 = 0$  (equal roots).

The recurrence relation for the first solution gives:  $(n^2+n)a_n + a_{n-1} = 0$

With  $a_0 = 1$ , we get:  $a_1 = -a_0/(1^2+1) = -1/2$   $a_2 = -a_1/(2^2+2) = 1/12$   $a_3 = -a_2/(3^2+3) = -1/144 \dots$

So the first solution is:  $y_1(x) = 1 - x/2 + x^2/12 - x^3/144 + \dots$

Since the roots are equal, the second solution includes a logarithmic term:

$$y_2(x) = y_1(x)\ln(x) + x^0(b_1x + b_2x^2 + \dots)$$

### Example 3: Roots Differing by 2

Consider the equation:  $x^2y'' + x(3-x)y' - (1+x)y = 0$

The indicial equation is:  $r(r-1) + 3r - 1 = 0 \Rightarrow r^2 + 2r - 1 = 0$

Using the quadratic formula:  $r = (-2 \pm \sqrt{4+4})/2 = -1 \pm \sqrt{2}$

So  $r_1 = -1 + \sqrt{2} \approx 0.414$  and  $r_2 = -1 - \sqrt{2} \approx -2.414$

Since  $r_1 - r_2 = 2.828$ , which is not an integer, we have two linearly independent solutions of the form:  $y_1(x) = x^{r_1}(a_0 + a_1x + a_2x^2 + \dots)$   $y_2(x) = x^{r_2}(b_0 + b_1x + b_2x^2 + \dots)$

No logarithmic term is needed in this case.

### Frobenius Method with Three Regular Singular Points

Some differential equations have more than one regular singular point. A classic example is the hypergeometric equation, which has three regular singular points at  $x = 0$ ,  $x = 1$ , and  $x = \infty$ .

For such equations, we can apply the Frobenius method at each singular point to find local solutions, and then connect these solutions using analytic continuation.

### Power Series Versus Frobenius Method

It's important to understand when to use the power series method versus the Frobenius method:

1. Power Series Method: Used when expanding around an ordinary point

- Assumes solution of the form:  $y(x) = \sum a_n(x - x_0)^n$
  - Works when  $P(x)$  and  $Q(x)$  are analytic at  $x_0$
2. Frobenius Method: Used when expanding around a regular singular point
- Assumes solution of the form:  $y(x) = (x - x_0)^r \sum a_n(x - x_0)^n$
  - Works when  $(x - x_0)P(x)$  and  $(x - x_0)^2Q(x)$  are analytic at  $x_0$

Attempting to use a power series at a singular point will generally fail, as the radius of convergence would be zero.

### **Irregular Singular Points**

When a point  $x_0$  is singular but not regularly singular (i.e., either  $(x - x_0)P(x)$  or  $(x - x_0)^2Q(x)$  is not analytic at  $x_0$ ), we call it an irregular singular point. The Frobenius method does not work for irregular singular points. Other methods, such as the method of asymptotic expansions or the WKB approximation, are needed for such cases.

### **Special Functions and the Frobenius Method**

Many special functions in mathematics are defined as solutions to differential equations with regular singular points. The Frobenius method provides a systematic way to develop these functions as power series.

Examples include:

- Bessel functions (solutions to Bessel's equation)
- Legendre polynomials (solutions to Legendre's equation)
- Hypergeometric functions (solutions to the hypergeometric equation)
- Laguerre polynomials
- Chebyshev polynomials

Understanding the Frobenius method is crucial for working with these special functions and their applications in physics, engineering, and other fields.

The Frobenius method is a powerful technique for solving differential equations with regular singular points. The key steps are:

1. Identify regular singular points
2. Assume a solution of the form  $y(x) = x^r \sum a_n x^n$
3. Find the indicial equation and determine its roots
4. Based on the nature of the roots, construct one or two linearly independent solutions
5. Pay special attention to exceptional cases where the roots differ by an integer or are equal

The exceptional cases require careful analysis to determine whether a logarithmic term is needed in the second solution. The criterion is based on the recurrence relation for the coefficients and involves checking whether certain conditions are satisfied when the index reaches the value of the difference between the roots. By mastering the Frobenius method, including the handling of exceptional cases, you can solve a wide range of differential equations that arise in mathematical physics and other applications.



## CHECK YOUR PROGRESS

- 1) Define a regular singular point.

.....  
.....  
.....

- 2) What is the Frobenius series form of the solution?

.....  
.....  
.....

- 3) What does the indicial equation determine?

.....  
.....  
.....

## LET US SUM UP

A point  $x_0$  is called a regular singular point of the differential equation  $y'' + P(x)y' + Q(x)y = 0$  if both  $(x - x_0)P(x)$  and  $(x - x_0)^2Q(x)$  are analytic near  $x_0$ . The Frobenius method provides a systematic way of solving such equations by assuming a power series solution of the form  $y = \sum_{k=0}^{\infty} a_k (x - x_0)^{k+r}$ . The exponent  $r$  is determined by solving the indicial equation.

## UNIT END EXERCISES

### Short Answer Questions:

1. Define a regular singular point of a differential equation.
2. What is Euler's equation, and how is it solved?
3. Explain the Frobenius method for solving differential equations.

### Long Answer Questions:

1. Derive and solve Euler's equation  $x^2y'' + 3xy' + 2y = 0$
2. Explain the Frobenius method in detail and apply it to solve  $x^2y'' + xy' - y = 0$
3. Derive the indicial equation for a second-order equation with a regular singular point.

### REFERENCES AND SUGGESTED READINGS:

1. Arfken, G. B., Weber, H. J., & Harris, F. E. (2022). *Mathematical Methods for Physicists* (7th ed.). Academic Press.
2. Watson, G. N. (2020). *A Treatise on the Theory of Bessel Functions*. Cambridge University Press.
3. Weinberger, H. F. (2012). *A First Course in Partial Differential Equations with Complex Variables and Transform Methods*. Dover Publications.

## UNIT 13

### EXCEPTIONAL CASES – BESSEL EQUATION.

#### 13.1 The Bessel Equation and Its Properties

The Bessel equation is a second-order linear differential equation that appears frequently in problems involving cylindrical or spherical symmetry. It emerges naturally when solving partial differential equations like the wave equation, Laplace's equation, or the heat equation in cylindrical coordinates.

The standard form of the Bessel equation is:

$$x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + (x^2 - n^2)y = 0$$

where  $n$  is a parameter that may be any real or complex number, though it's most commonly a non-negative integer in physical applications. This equation is named after Friedrich Wilhelm Bessel, a German astronomer and mathematician who studied it extensively in the early 19th century.

#### Solutions to the Bessel Equation: Bessel Functions

The solutions to the Bessel equation are called Bessel functions. There are several types:

Bessel Functions of the First Kind:  $J_n(x)$

For any value of  $n$ , the Bessel function of the first kind, denoted  $J_n(x)$ , is defined by the series:

$$J_n(x) = \sum_{k=0}^{\infty} \frac{(-1)^k}{k! \Gamma(n+k+1)} \left(\frac{x}{2}\right)^{2k+n}$$

where  $\Gamma$  is the gamma function, which extends the factorial function to non-integer values.

When  $n$  is a non-negative integer, the series simplifies to:

$$J_n(x) = \sum_{k=0}^{\infty} \frac{(-1)^k}{k! (n+k)!} \left(\frac{x}{2}\right)^{2k+n}$$

For integer values of  $n$ ,  $J_n(x)$  is finite at  $x = 0$ , making it particularly useful for physical problems where a bounded solution at the origin is required.

Bessel Functions of the Second Kind:  $Y_n(x)$

The Bessel function of the second kind (also called the Neumann function or Weber function), denoted  $Y_n(x)$ , forms another linearly independent solution to the Bessel equation:

$$Y_n(x) = (J_n(x)\cos(n\pi) - J_{-n}(x)) / \sin(n\pi), \text{ for non-integer } n$$

$$\lim_{n \rightarrow m} (J_m(x)\cos(m\pi) - J_{-m}(x)) / \sin(m\pi), \text{ for integer } n$$

$Y_n(x)$  is singular at  $x = 0$ , so it's often excluded from physical problems requiring bounded solutions at the origin.

### **Modified Bessel Functions: $I_n(x)$ and $K_n(x)$**

If we replace  $x$  with  $ix$  in the Bessel equation, we get the modified Bessel equation:

$$x^2 d^2y/dx^2 + x dy/dx - (x^2 + n^2)y = 0$$

The solutions to this equation are the modified Bessel functions:

- The modified Bessel function of the first kind:  $I_n(x)$
- The modified Bessel function of the second kind (or MacDonald function):  $K_n(x)$

These functions are related to  $J_n(x)$  and  $Y_n(x)$  by:

$$I_n(x) = i(-n) J_n(ix) \quad K_n(x) = (\pi/2) i(n+1) [J_n(ix) + iY_n(ix)]$$

### **Important Properties of Bessel Functions**

#### **Recurrence Relations**

Bessel functions satisfy several important recurrence relations that make them easier to work with:

1.  $J_{(n-1)}(x) + J_{(n+1)}(x) = (2n/x) J_n(x)$
2.  $J_{(n-1)}(x) - J_{(n+1)}(x) = 2J'_n(x)$
3.  $J'_n(x) = (n/x) J_n(x) - J_{(n+1)}(x)$
4.  $J'_n(x) = J_{(n-1)}(x) - (n/x) J_n(x)$

Similar relations exist for  $Y_n(x)$ ,  $I_n(x)$ , and  $K_n(x)$ .

#### **Orthogonality**

The Bessel functions of the first kind satisfy an orthogonality relation:

$$\int_0^a x J_n(\alpha_m x) J_n(\alpha_k x) dx = 0, \text{ for } m \neq k$$

where  $\alpha_m$  and  $\alpha_k$  are the  $m$ th and  $k$ th positive roots of  $J_n(a x) = 0$ .

This orthogonality property makes Bessel functions useful in solving boundary-value problems and in Fourier-Bessel series.

### Asymptotic Behavior

For large values of  $x$ , the Bessel functions have the following asymptotic behavior:

$$J_n(x) \approx \sqrt{2/\pi x} \cos(x - n\pi/2 - \pi/4) \quad Y_n(x) \approx \sqrt{2/\pi x} \sin(x - n\pi/2 - \pi/4)$$

For small values of  $x$  when  $n > 0$ :

$$J_n(x) \approx (1/n!) * (x/2)^n \quad Y_{-n}(x) \approx -(n-1)!/\pi * (2/x)^n$$

### Zeros of Bessel Functions

The zeros of Bessel functions are important in many applications. Let's denote the  $k$ th positive zero of  $J_n(x)$  as  $j_{n,k}$ .

For large  $k$ , the zeros are approximately:

$$j_{n,k} \approx (k + n/2 - 1/4)\pi$$

The zeros of  $J_n(x)$  and  $J_{n+1}(x)$  interlace, meaning between any two consecutive zeros of  $J_n(x)$ , there's exactly one zero of  $J_{n+1}(x)$ .

### Differential Equations Related to the Bessel Equation

Several important equations in mathematical physics can be transformed into the Bessel equation or its variations:

#### The Airy Equation

The Airy equation is:

$$d^2y/dx^2 - xy = 0$$

Its solutions are the Airy functions, which can be expressed in terms of Bessel functions of order  $\pm 1/3$ .

#### The Spherical Bessel Equation

The spherical Bessel equation is:

$$x^2 d^2y/dx^2 + 2x dy/dx + [x^2 - n(n+1)]y = 0$$

Its solutions, the spherical Bessel functions  $j_n(x)$  and  $y_n(x)$ , are related to the regular Bessel functions by:

$$j_n(x) = \sqrt{\pi/2x} J_{n+1/2}(x) \quad y_n(x) = \sqrt{\pi/2x} Y_{n+1/2}(x)$$

### **The Associated Legendre Equation**

While not directly a Bessel equation, the associated Legendre equation is related and often appears alongside Bessel functions in physical problems, especially when separating variables in spherical coordinates.

### **Generating Functions and Integral Representations**

Generating Function for  $J_n(x)$

The generating function for Bessel functions of the first kind is:

$$\exp(x(t-1/t)/2) = \sum_{n=-\infty}^{\infty} t^n J_n(x)$$

This is useful for deriving properties of Bessel functions.

### **Integral Representations**

Bessel functions can also be represented by integrals:

$$J_n(x) = (1/\pi) \int_0^\pi \cos(n\theta - x \sin \theta) d\theta$$

This representation is useful in proving certain properties and in numerical computations.

### **Applications of Bessel Functions in Mathematics**

#### **Fourier-Bessel Series**

Functions defined on a disk can be expanded in terms of Bessel functions:

$$f(r) = \sum_{m=1}^{\infty} c_m J_0(j_{0,m} r/a)$$

where  $j_{0,m}$  are the zeros of  $J_0(x)$  and  $c_m$  are the coefficients determined by the orthogonality properties.

#### **Hankel Transform**

The Hankel transform uses Bessel functions as kernels:

$$F(k) = \int_0^\infty f(r) J_n(kr) r dr$$

This transform is particularly useful for problems with cylindrical symmetry.

## Computational Aspects of Bessel Functions

### Computing Bessel Functions

Bessel functions can be computed using:

1. Direct series evaluation (for small  $x$ )
2. Recurrence relations (for moderate  $x$ )
3. Asymptotic formulas (for large  $x$ )
4. Continued fractions
5. Numerical integration of the integral representations

### Special Values

Some special values of Bessel functions include:

- $J_0(0) = 1$ , while  $J_n(0) = 0$  for  $n > 0$
- $Y_n(0)$  is undefined (singular)
- $I_0(0) = 1$ , while  $I_n(0) = 0$  for  $n > 0$
- $K_n(0)$  is undefined (singular) for all  $n$

## 13.2 Applications of the Bessel Equation

Bessel functions appear in a wide range of physical and engineering applications. We'll explore some of the most important ones.

### Vibrating Membranes and Drums

The vibration of a circular membrane (like a drum) is governed by the wave equation in cylindrical coordinates:

$$\partial^2 u / \partial t^2 = c^2 (\partial^2 u / \partial r^2 + (1/r) \partial u / \partial r + (1/r^2) \partial^2 u / \partial \theta^2)$$

Using separation of variables  $u(r, \theta, t) = R(r)\Theta(\theta)T(t)$ , we get:

$$r^2 R'' + r R' + (\lambda^2 r^2 - n^2)R = 0$$

This is precisely the Bessel equation of order  $n$  with solution:

$$R(r) = A J_n(\lambda r) + B Y_n(\lambda r)$$

For a circular membrane with fixed edges (like a drum), we need  $R(a) = 0$ , where  $a$  is the radius. Since  $Y_n$  is singular at  $r = 0$ , we must set  $B = 0$ , and the boundary condition gives  $J_n(\lambda a) = 0$ , meaning  $\lambda = j_{n,k}/a$ , where  $j_{n,k}$  is the  $k$ th zero of  $J_n$ .

The natural frequencies of vibration are then:

$$\omega_{n,k} = (c/a) j_{n,k}$$

The general solution for the displacement of the membrane is a superposition of modes:

$$u(r, \theta, t) = \sum_{n=0}^{\infty} \sum_{k=1}^{\infty} [A_{n,k} \cos(\omega_{n,k} t) + B_{n,k} \sin(\omega_{n,k} t)] \times J_n(j_{n,k} r/a) \times [C_n \cos(n\theta) + D_n \sin(n\theta)]$$

### Heat Conduction in Cylindrical Bodies

The heat equation in cylindrical coordinates is:

$$\partial u / \partial t = \alpha (\partial^2 u / \partial r^2 + (1/r) \partial u / \partial r + (1/r^2) \partial^2 u / \partial \theta^2 + \partial^2 u / \partial z^2)$$

For problems with cylindrical symmetry ( $\partial u / \partial \theta = 0$ ,  $\partial u / \partial z = 0$ ), this simplifies to:

$$\partial u / \partial t = \alpha (\partial^2 u / \partial r^2 + (1/r) \partial u / \partial r)$$

Using separation of variables  $u(r, t) = R(r)T(t)$ , we get the Bessel equation for  $R(r)$ :

$$r^2 R'' + r R' + \lambda^2 r^2 R = 0$$

The solution involves Bessel functions, with the specific boundary conditions determining which Bessel functions to use.

### Electromagnetic Waves in Waveguides

In electromagnetic theory, cylindrical waveguides lead to Bessel equations. The propagation of electromagnetic waves in a circular waveguide is governed by Maxwell's equations, which, after separation of variables, lead to Bessel equations.

For TE modes (transverse electric), the boundary condition at the waveguide wall  $r = a$  gives:



$$J'_n(\kappa a) = 0$$

For TM modes (transverse magnetic), the boundary condition gives:

$$J_n(\kappa a) = 0$$

Where  $\kappa$  is related to the cutoff frequency of the waveguide.

### **Quantum Mechanics: Particle in a Cylindrical Box**

In quantum mechanics, the Schrödinger equation for a particle confined in a cylindrical box leads to Bessel equations. The wavefunctions involve Bessel functions, and the energy eigenvalues are related to the zeros of these functions.

### **Fluid Flow Through Pipes**

The velocity profile for laminar flow through a cylindrical pipe is related to Bessel functions. For pulsatile flow, the solution involves Bessel functions of the first kind.

### **Diffraction of Light**

In optics, the diffraction pattern of light passing through a circular aperture is described by Bessel functions. The intensity pattern is given by:

$$I(\theta) = I_0 [2J_1(ka \sin \theta)/(ka \sin \theta)]^2$$

where  $k$  is the wave number,  $a$  is the radius of the aperture, and  $\theta$  is the angle of diffraction.

### **Stress and Strain in Cylindrical Bodies**

In elasticity theory, the stress and strain in cylindrical bodies often involve Bessel functions. For example, the torsion of a circular shaft and the bending of cylindrical plates are problems where Bessel functions naturally appear.

### **Acoustics: Sound Propagation in Pipes**

The propagation of sound waves in cylindrical pipes is described by the wave equation in cylindrical coordinates, leading to Bessel functions. The resonant frequencies of organ pipes and wind instruments are related to the zeros of Bessel functions.

### **Electrical Conductors: Skin Effect**

The skin effect in electrical conductors, where alternating current tends to flow near the surface, is described by Bessel functions. The current density as a function of radius is given by:

$$J(r) = J_0 \times J_0(\sqrt{-i\omega\mu\sigma} r) / J_0(\sqrt{-i\omega\mu\sigma} a)$$

where  $J_0$  is the current density at the surface,  $\omega$  is the angular frequency,  $\mu$  is the permeability,  $\sigma$  is the conductivity, and  $a$  is the radius of the conductor.

### Earth's Magnetic Field

Models of the Earth's magnetic field use spherical harmonics, which are related to associated Legendre polynomials and spherical Bessel functions.

### Solved Problems

#### Solved Problem 1: Vibrating Circular Membrane

Problem: Find the normal modes of vibration for a circular membrane of radius  $a$  with fixed boundary.

Solution:

The displacement  $u(r, \theta, t)$  of a point on the membrane satisfies the wave equation:

$$\partial^2 u / \partial t^2 = c^2 (\partial^2 u / \partial r^2 + (1/r) \partial u / \partial r + (1/r^2) \partial^2 u / \partial \theta^2)$$

Using separation of variables,  $u(r, \theta, t) = R(r)\Theta(\theta)T(t)$ , we get:

$$T''(t) + \omega^2 T(t) = 0 \quad \Theta''(\theta) + n^2 \Theta(\theta) = 0 \quad r^2 R''(r) + r R'(r) + (\omega^2 r^2 / c^2 - n^2) R(r) = 0$$

The solutions are:  $T(t) = A \cos(\omega t) + B \sin(\omega t)$   $\Theta(\theta) = C \cos(n\theta) + D \sin(n\theta)$ , where  $n$  must be an integer for periodicity  $R(r) = E J_n(\omega r / c) + F Y_n(\omega r / c)$

Since  $Y_n$  is singular at  $r = 0$  and the solution must be bounded at the origin,  $F = 0$ .

The boundary condition  $u(a, \theta, t) = 0$  gives  $J_n(\omega a / c) = 0$ , which means  $\omega = (c/a) j_{(n,k)}$ , where  $j_{(n,k)}$  is the  $k$ th zero of  $J_n$ .

Therefore, the normal modes are:

$$u_{(n,k)}(r, \theta, t) = J_n(j_{(n,k)} r / a) [C_{(n,k)} \cos(n\theta) + D_{(n,k)} \sin(n\theta)] [A_{(n,k)} \cos(\omega_{(n,k)} t) + B_{(n,k)} \sin(\omega_{(n,k)} t)]$$

with frequencies  $\omega_{(n,k)} = (c/a) j_{(n,k)}$ .

The fundamental frequency (lowest) corresponds to  $j_{0,1} \approx 2.4048$ , giving  $\omega_{0,1} = 2.4048c/a$ .

### Solved Problem 2: Heat Conduction in a Solid Cylinder

Problem: A solid cylinder of radius  $a$  initially has temperature distribution  $T(r,0) = T_0(1-r^2/a^2)$ . The surface is kept at temperature 0. Find the temperature distribution  $T(r,t)$  for  $t > 0$ .

Solution:

The heat equation in cylindrical coordinates with radial symmetry is:

$$\partial T / \partial t = \alpha (\partial^2 T / \partial r^2 + (1/r) \partial T / \partial r)$$

with initial condition  $T(r,0) = T_0(1-r^2/a^2)$  and boundary condition  $T(a,t) = 0$ .

Let's define the dimensionless variables:

$$u = T/T_0, \rho = r/a, \tau = \alpha t/a^2$$

The heat equation becomes:

$$\partial u / \partial \tau = \partial^2 u / \partial \rho^2 + (1/\rho) \partial u / \partial \rho$$

with  $u(\rho,0) = 1-\rho^2$  and  $u(1,\tau) = 0$ .

Using separation of variables,  $u(\rho,\tau) = R(\rho)S(\tau)$ , we get:

$$S'(\tau) + \lambda^2 S(\tau) = 0 \quad \rho^2 R''(\rho) + \rho R'(\rho) + \lambda^2 \rho^2 R(\rho) = 0$$

The solutions are:  $S(\tau) = e^{(-\lambda^2 \tau)}$   $R(\rho) = AJ_0(\lambda \rho) + BY_0(\lambda \rho)$

Since  $Y_0$  is singular at  $\rho = 0$ ,  $B = 0$ . The boundary condition  $R(1) = 0$  gives  $J_0(\lambda) = 0$ , so  $\lambda = j_{0,k}$ , the  $k$ th zero of  $J_0$ .

The general solution is:

$$u(\rho,\tau) = \sum c_k J_0(j_{0,k} \rho) e^{-(j_{0,k})^2 \tau} \quad k=1$$

The coefficients  $c_k$  are determined from the initial condition:

$$1-\rho^2 = \sum c_k J_0(j_{0,k} \rho) \quad k=1$$

Using the orthogonality of Bessel functions:

$$c_k = (2/[j_{0,k}]^2 J_1(j_{0,k})) \int_0^1 \rho(1-\rho^2) J_0(j_{0,k}\rho) d\rho$$

This integral evaluates to:

$$c_k = 2/(j_{0,k} J_1(j_{0,k}))$$

Therefore, the temperature distribution is:

$$T(r,t) = 2T_0 \sum_{k=1}^{\infty} (1/(j_{0,k} J_1(j_{0,k}))) J_0(j_{0,k}r/a) e^{-(j_{0,k})^2 \alpha t/a^2}$$

### Solved Problem 3: Bessel Series Expansion

Problem: Expand the function  $f(x) = x$  for  $0 \leq x \leq 1$  in terms of Bessel functions of the first kind of order zero.

Solution:

We want to express  $f(x) = x$  as a series:

$$f(x) = \sum_{m=1}^{\infty} c_m J_0(j_{0,m}x)$$

where  $j_{0,m}$  is the  $m$ th positive zero of  $J_0$ .

Using the orthogonality property of Bessel functions:

$$\int_0^1 x J_0(j_{0,m}x) J_0(j_{0,n}x) dx = 0 \text{ for } m \neq n$$

and

$$\int_0^1 x [J_0(j_{0,m}x)]^2 dx = (1/2)[J_1(j_{0,m})]^2$$

we can find the coefficients:

$$c_m = (\int_0^1 x^2 J_0(j_{0,m}x) dx) / (\int_0^1 x [J_0(j_{0,m}x)]^2 dx)$$

Using integration by parts and the properties of Bessel functions:

$$\int_0^1 x^2 J_0(j_{0,m}x) dx = (2/j_{0,m}) J_1(j_{0,m})$$

Therefore:

$$c_m = (2/j_{0,m}) J_1(j_{0,m}) / ((1/2)[J_1(j_{0,m})]^2) = 4/(j_{0,m} J_1(j_{0,m}))$$

The Bessel series expansion is:

$$f(x) = x = \sum_{m=1}^{\infty} (4/(j_{0,m} J_1(j_{0,m}))) J_0(j_{0,m}x)$$

### Solved Problem 4: Wave Equation with Bessel Functions

Problem: Solve the wave equation  $\partial^2 u / \partial t^2 = c^2 \nabla^2 u$  in a circular region of radius  $a$  with boundary condition  $u(a, \theta, t) = 0$  and initial conditions  $u(r, \theta, 0) = f(r, \theta)$ ,  $\partial u / \partial t(r, \theta, 0) = g(r, \theta)$ .

Solution:

In cylindrical coordinates, the wave equation is:

$$\partial^2 u / \partial t^2 = c^2 (\partial^2 u / \partial r^2 + (1/r) \partial u / \partial r + (1/r^2) \partial^2 u / \partial \theta^2)$$

Using separation of variables,  $u(r, \theta, t) = R(r)\Theta(\theta)T(t)$ , we get:

$$T''(t) + \omega^2 T(t) = 0 \quad \Theta''(\theta) + n^2 \Theta(\theta) = 0 \quad r^2 R''(r) + r R'(r) + (\omega^2 r^2 / c^2 - n^2) R(r) = 0$$

The solutions are:  $T(t) = A \cos(\omega t) + B \sin(\omega t)$   $\Theta(\theta) = C \cos(n\theta) + D \sin(n\theta)$ , where  $n$  is an integer  $R(r) = E J_n(\omega r / c) + F Y_n(\omega r / c)$

Since  $Y_n$  is singular at  $r = 0$ ,  $F = 0$ . The boundary condition  $u(a, \theta, t) = 0$  gives  $J_n(\omega a / c) = 0$ , meaning  $\omega = (c/a) j_{(n,k)}$ .

The general solution is:

$$u(r, \theta, t) = \sum_n \sum_k [A_{(n,k)} \cos(\omega_{(n,k)} t) + B_{(n,k)} \sin(\omega_{(n,k)} t)] \times J_n(j_{(n,k)} r / a) \times [C_n \cos(n\theta) + D_n \sin(n\theta)] \quad n=0 \quad k=1$$

where  $\omega_{(n,k)} = (c/a) j_{(n,k)}$ .

The coefficients are determined from the initial conditions:

$$f(r, \theta) = \sum_n \sum_k A_{(n,k)} J_n(j_{(n,k)} r / a) [C_n \cos(n\theta) + D_n \sin(n\theta)] \quad n=0 \quad k=1$$

$$g(r, \theta) = \sum_n \sum_k B_{(n,k)} \omega_{(n,k)} J_n(j_{(n,k)} r / a) [C_n \cos(n\theta) + D_n \sin(n\theta)] \quad n=0 \quad k=1$$

Using the orthogonality properties of trigonometric functions and Bessel functions, we can find the coefficients.

For example, if  $f(r, \theta) = f(r)$  (independent of  $\theta$ ) and  $g(r, \theta) = 0$ , then:

$$A_{(0,k)} = (2 / (a^2 [J_1(j_{(0,k)})]^2)) \int_0^a r f(r) J_0(j_{(0,k)} r / a) dr \quad 0$$

$$B_{(n,k)} = 0 \text{ for all } n, k \quad A_{(n,k)} = 0 \text{ for } n > 0, \text{ all } k$$

And the solution simplifies to:

$$u(r, t) = \sum_k A_{(0,k)} \cos(\omega_{(0,k)} t) J_0(j_{(0,k)} r / a) \quad k=1$$

### Solved Problem 5: Quantum Particle in a Cylindrical Box

Problem: Find the energy eigenvalues and eigenfunctions for a quantum particle confined in a cylindrical box of radius  $a$  and height  $h$ .

Solution:

The time-independent Schrödinger equation in cylindrical coordinates is:

$$-\hbar^2/(2m) (\partial^2\psi/\partial r^2 + (1/r)\partial\psi/\partial r + (1/r^2)\partial^2\psi/\partial\theta^2 + \partial^2\psi/\partial z^2) = E\psi$$

With boundary conditions:  $\psi(a, \theta, z) = 0$  for all  $\theta$ ,  $0 \leq z \leq h$   $\psi(r, \theta, 0) = \psi(r, \theta, h) = 0$  for all  $r, \theta$

Using separation of variables,  $\psi(r, \theta, z) = R(r)\Theta(\theta)Z(z)$ , we get:

$$Z''(z) + k_z^2 Z(z) = 0 \quad \Theta''(\theta) + m^2 \Theta(\theta) = 0 \quad r^2 R''(r) + r R'(r) + (k_r^2 r^2 - m^2) R(r) = 0$$

Where  $k_r^2 + k_z^2 = 2mE/\hbar^2$ .

The solutions are:  $Z(z) = A \sin(k_z z)$ , with  $k_z = n\pi/h$ ,  $n = 1, 2, 3, \dots$   $\Theta(\theta) = B \cos(m\theta) + C \sin(m\theta)$ , where  $m$  is an integer  $R(r) = D J_m(k_r r)$ , with  $k_r = j_{(m,l)}/a$ , where  $j_{(m,l)}$  is the  $l$ th zero of  $J_m$

The energy eigenvalues are:

$$E_{(n,m,l)} = (\hbar^2/2m) [(j_{(m,l)}/a)^2 + (n\pi/h)^2]$$

And the normalized eigenfunctions are:

$$\psi_{(n,m,l)}(r, \theta, z) = N_{(n,m,l)} J_m(j_{(m,l)}r/a) [\cos(m\theta) \text{ or } \sin(m\theta)] \sin(n\pi z/h)$$

where  $N_{(n,m,l)}$  is a normalization constant:

$$N_{(n,m,l)} = (\sqrt{2}/h) / (a J_{(m+1)}(j_{(m,l)}) \sqrt{\pi}) \text{ for } m > 0 \quad N_{(n,0,l)} = (\sqrt{2}/h) / (a J_1(j_{(0,l)}) \sqrt{2\pi}) \text{ for } m = 0$$

The ground state corresponds to  $n = 1$ ,  $m = 0$ ,  $l = 1$ , with energy:

$$E_1 = (\hbar^2/2m) [(j_{(0,1)}/a)^2 + (\pi/h)^2]$$

## Unsolved Problems

### Unsolved Problem 1

A circular membrane of radius  $a$  is fixed at the boundary and has initial displacement  $u(r, 0) = u_0(1 - r^2/a^2)$  and zero initial velocity. Find the displacement  $u(r, t)$  for  $t > 0$ .

### Unsolved Problem 2

Solve the heat conduction problem in a hollow cylinder with inner radius  $a$  and outer radius  $b$ . The inner surface is insulated ( $\partial T / \partial r = 0$  at  $r = a$ ), and the outer surface is kept at temperature  $T = 0$ . The initial temperature distribution is  $T(r, 0) = T_0$ .

### Unsolved Problem 3

Find the first three terms of the asymptotic expansion of  $J_n(x)$  for large  $x$ .

### Unsolved Problem 4

A circular waveguide of radius  $a$  has perfectly conducting walls. Find the cutoff frequencies for the  $TE_{mn}$  and  $TM_{mn}$  modes, and determine which mode has the lowest cutoff frequency.

### Unsolved Problem 5

Prove the addition theorem for Bessel functions:

$$J_0(\sqrt{x^2 + y^2 - 2xy \cos \theta}) = J_0(x)J_0(y) + 2 \sum_{n=1}^{\infty} J_n(x)J_n(y) \cos(n\theta)$$

## SELF ASSESSMENT QUESTIONS

### Multiple Choice Questions (MCQs)

1. A regular singular point of a differential equation is a point where:
  - a) The equation is not defined
  - b) The coefficient functions have singularities that are not too severe
  - c) The solution does not exist
  - d) None of the above
2. Euler's equation has the form:
  - a)  $x^2 y'' + ax y' + by = 0$
  - b)  $y'' + p(x)y' + q(x)y = 0$
  - c)  $y' + py = 0$
  - d) None of the above
3. The Frobenius method is used to:
  - a) Solve equations with regular singular points
  - b) Solve equations with constant coefficients
  - c) Find the Wronskian
  - d) None of the above

4. A differential equation has a regular singular point if:
  - a) The coefficient functions satisfy a certain growth condition
  - b) The coefficient functions are discontinuous
  - c) The solution does not exist
  - d) None of the above
5. The characteristic equation in the Frobenius method is obtained from:
  - a) The lowest power of  $x$  in the series expansion
  - b) The highest power of  $x$  in the series expansion
  - c) The Wronskian determinant
  - d) None of the above
6. The Bessel equation arises in:
  - a) Vibrations of circular membranes
  - b) Heat conduction problems
  - c) Both (a) and (b)
  - d) None of the above
7. The solution of the Bessel equation involves:
  - a) Bessel functions of the first and second kind
  - b) Exponential functions
  - c) Polynomial solutions
  - d) None of the above
8. If two roots of the characteristic equation differ by an integer, the solutions are:
  - a) Linearly dependent
  - b) Linearly independent
  - c) Nonexistent
  - d) None of the above
9. The indicial equation is derived from:
  - a) The lowest exponent in the Frobenius method
  - b) The highest exponent in the Frobenius method
  - c) The Wronskian determinant
  - d) None of the above
10. The Bessel function  $J_n(x)J_{-n}(x)$  is defined as a series solution of:



a)  $x^2y''+xy'+(x^2-n^2)y=0$

b)  $y''+p(x)y'+q(x)y=0$

c)  $y'+py=0$

d) None of the above

Answer Key:

1	b	3	a	5	a	7	a	9	a
2	a	4	a	6	c	8	a	10	a

## CHECK YOUR PROGRESS

- 1) Write Bessel's differential equation.

.....  
.....  
.....

- 2) Define Bessel functions of the first and second kind.

.....  
.....  
.....

- 3) What are exceptional cases in the Frobenius method?

.....  
.....  
.....

- 4) Write one recurrence relation for Bessel functions.

.....  
.....  
.....

## LET US SUM UP

Bessel's equation, expressed as  $x^2y'' + xy' + (x^2 - n^2)y = 0$ , is a fundamental second-order differential equation with a regular singular point at  $x = 0$ . Its solutions are the Bessel functions of the first kind  $J_n(x)$  and of the second kind  $Y_n(x)$ . Exceptional cases occur when the difference between the roots of the indicial equation is an integer, requiring special forms of the second solution.

## UNIT END EXERCISES

### Short Answer Questions:

1. What are Bessel functions, and how are they defined?
2. Explain the importance of the characteristic equation in the Frobenius method.
3. What happens when the roots of the indicial equation differ by an integer?
4. How do singular points affect the solutions of differential equations?

### Long Answer Questions:

1. Solve the initial value problem for a differential equation with a singular point.
2. Discuss the connection between the Bessel equation and Fourier series.
3. Compare the Frobenius method with the method of undetermined coefficients.
4. Solve a second-order differential equation with a singular point using a power series method.

### REFERENCES AND SUGGESTED READINGS:

1. Arfken, G. B., Weber, H. J., & Harris, F. E. (2022). *Mathematical Methods for Physicists* (7th ed.). Academic Press.
2. Böcher, M. (2015). *Regular Points of Linear Differential Equations of the Second Order*, American Mathematical Society.
3. Watson, G. N. (2020). *A Treatise on the Theory of Bessel Functions*. Cambridge University Press.
4. Weinberger, H. F. (2012). *A First Course in Partial Differential Equations with Complex Variables and Transform Methods*. Dover Publications.
5. Stakgold, I., & Holst, M. J. (2019). *Green's Functions and Boundary Value Problems* (3rd ed.). Wiley.

## BLOCK 5

### UNIT 14

#### EXISTENCE AND UNIQUENESS OF SOLUTIONS TO FIRST-ORDER EQUATIONS

##### Objectives

- Understand conditions for the existence and uniqueness of solutions to first-order differential equations.
- Learn the method of solving separable differential equations.
- Study exact equations and integrating factors.
- Explore the method of successive approximations.
- Examine the Lipschitz condition and its role in uniqueness.
- Analyze the convergence of successive approximations.

##### 14.1 Introduction to Existence and Uniqueness Theorems

Differential equations are fundamental to describing natural phenomena and modeling real-world systems. When we formulate a differential equation to model a physical situation, two critical questions arise:

1. Does a solution to the differential equation actually exist?
2. If a solution exists, is it the only possible solution?

These questions lead us to the concepts of existence and uniqueness theorems, which provide conditions under which we can guarantee that a differential equation has a solution and that the solution is unique.

##### The Initial Value Problem

Before discussing existence and uniqueness, let's establish what we mean by a solution to a differential equation. Consider a first-order differential equation of the form:

$$dy/dx = f(x, y)$$

Along with an initial condition:

$$y(x_0) = y_0$$

This combination is called an Initial Value Problem (IVP). A solution to this IVP is a function  $y = \varphi(x)$  that:

- Satisfies the differential equation  $dy/dx = f(x, y)$  for all  $x$  in some interval containing  $x_0$
- Satisfies the initial condition  $\varphi(x_0) = y_0$

### **The Existence Theorem**

The existence theorem for first-order differential equations provides conditions under which we can guarantee that a solution to an IVP exists.

Existence Theorem (Informal Statement): If  $f(x, y)$  and  $\partial f/\partial y$  are continuous functions in some rectangle  $R$  containing the point  $(x_0, y_0)$ , then there exists at least one solution to the initial value problem:

- $dy/dx = f(x, y)$
- $y(x_0) = y_0$

This solution is valid in some interval containing  $x_0$ .

The existence theorem tells us that if our function  $f(x, y)$  is well-behaved (continuous) in a region containing our initial point, then a solution exists, at least for some interval around the initial point.

### **The Uniqueness Theorem**

The uniqueness theorem addresses the second question: whether the solution is unique.

Uniqueness Theorem (Informal Statement): If  $f(x, y)$  and  $\partial f/\partial y$  are continuous functions in some rectangle  $R$  containing the point  $(x_0, y_0)$ , then there exists exactly one solution to the initial value problem:

- $dy/dx = f(x, y)$
- $y(x_0) = y_0$

This unique solution is valid in some interval containing  $x_0$ .

Notice that the conditions for uniqueness are the same as those for existence in this statement. The key addition is that the partial derivative of  $f$  with respect to  $y$  must also be continuous.

## Practical Implications

These theorems have important practical implications:

1. **Predictability:** In physical systems, uniqueness guarantees that identical initial conditions always lead to the same outcome, ensuring predictability.
2. **Numerical Methods:** When implementing numerical methods to approximate solutions, we need to know that a solution exists and is unique to ensure our approximations converge to the correct solution.
3. **Interval of Existence:** The theorems guarantee solutions only on some interval containing the initial point, not necessarily for all values of  $x$ . Determining this interval can be crucial in applications.

## Geometric Interpretation

Geometrically, the differential equation  $dy/dx = f(x, y)$  defines a direction field (or slope field) in the  $xy$ -plane. At each point  $(x, y)$ , the value  $f(x, y)$  gives the slope of a small line segment.

- The existence theorem ensures that we can find a curve passing through  $(x_0, y_0)$  that follows the direction field.
- The uniqueness theorem ensures that only one such curve passes through  $(x_0, y_0)$ .

## Examples Where Uniqueness Fails

It's instructive to look at cases where the conditions for uniqueness fail:

**Example 1: Consider the differential equation:**

$$dy/dx = 3y^{(2/3)}$$

With the initial condition  $y(0) = 0$ .

The function  $f(x, y) = 3y^{(2/3)}$  is continuous, but its partial derivative with respect to  $y$ ,  $\partial f/\partial y = 2y^{(-1/3)}$ , is not continuous at  $y = 0$ . In this case, the IVP has multiple solutions:

$$y(x) = 0 \text{ for all } x \quad y(x) = x^3 \text{ for } x \geq 0, \quad y(x) = -x^3 \text{ for } x \leq 0$$

**Example 2: Consider:**

$$dy/dx = y/x$$

With the initial condition  $y(0) = 0$ .

Here,  $f(x, y) = y/x$  is not continuous at  $x = 0$ , violating the conditions of the existence theorem. Indeed, no solution can satisfy both the differential equation and the initial condition.

**Picard's Theorem**

A more detailed version of the existence and uniqueness theorem is given by Picard's theorem, which not only provides conditions for existence and uniqueness but also suggests a method for constructing the solution through successive approximations.

Picard's Theorem (Simplified): If  $f(x, y)$  satisfies a Lipschitz condition with respect to  $y$  in some region containing  $(x_0, y_0)$ , then the IVP has a unique solution in some interval containing  $x_0$ .

The Lipschitz condition essentially requires that the rate of change of  $f$  with respect to  $y$  is bounded, which is a slightly weaker condition than requiring  $\partial f/\partial y$  to be continuous.

**Global Existence**

The theorems discussed so far guarantee existence and uniqueness only locally, in some interval around the initial point. For some applications, we need to know whether the solution exists for all values of  $x$  in a given range.

Global Existence Theorem (Informal): If  $f(x, y)$  and  $\partial f/\partial y$  are continuous for all  $(x, y)$  in a strip  $a \leq x \leq b$ ,  $-\infty < y < \infty$ , and  $|f(x, y)| \leq M$  (a constant) in this strip, then any solution of  $dy/dx = f(x, y)$  with  $y(x_0) = y_0$  (where  $a \leq x_0 \leq b$ ) exists throughout the entire interval  $[a, b]$ .

This theorem is particularly useful when we can establish bounds on the growth of solutions.

**14.2 Equations with Separable Variables**

Separable differential equations represent one of the simplest classes of differential equations that can be solved analytically. A first-order differential equation is separable if it can be written in the form:

$$dy/dx = g(x)h(y)$$

where  $g$  is a function of  $x$  alone and  $h$  is a function of  $y$  alone.

The significance of separable equations lies in their direct method of solution and their frequent appearance in various applications, from physics to biology.

### The Method of Separation of Variables

The core idea behind solving separable equations is to rearrange the equation so that all terms containing  $y$  are on one side and all terms with  $x$  are on the other. Then, we integrate both sides.

For a differential equation in the form  $dy/dx = g(x)h(y)$ , we follow these steps:

1. Rearrange to separate variables:  $(1/h(y))dy = g(x)dx$
2. Integrate both sides:  $\int (1/h(y))dy = \int g(x)dx$
3. Solve for  $y$  if possible

Let's see this method in action with some examples.

### Solved Examples

#### Example 1: Basic Separation

Problem: Solve the differential equation  $dy/dx = xy$ .

**Solution:**

Step 1: Rearrange to separate variables.  $dy/y = x dx$

Step 2: Integrate both sides.  $\int (dy/y) = \int x dx \ln|y| = x^2/2 + C$  (where  $C$  is an arbitrary constant)

Step 3: Solve for  $y$ .  $|y| = e^{(x^2/2 + C)} = e^C \cdot e^{(x^2/2)}$   $y = \pm e^C \cdot e^{(x^2/2)}$

Since  $e^C$  is a positive constant, we can simplify by letting  $K = \pm e^C$ , which gives:  $y = K \cdot e^{(x^2/2)}$

Therefore, the general solution is  $y = K \cdot e^{(x^2/2)}$ , where  $K$  is an arbitrary non-zero constant.

If we have an initial condition, say  $y(0) = 2$ , we can determine  $K$ :  $2 = K \cdot e^{(0^2/2)}$   $2 = K \cdot 1$   $K = 2$



So the particular solution would be  $y = 2e^{(x^2/2)}$ .

### Example 2: Growth and Decay

Problem: Solve the differential equation  $dy/dx = ky$ , where  $k$  is a constant, with the initial condition  $y(0) = y_0$ .

#### Solution:

This is a classic equation describing exponential growth ( $k > 0$ ) or decay ( $k < 0$ ).

Step 1: Separate variables.  $dy/y = k dx$

Step 2: Integrate both sides.  $\int (dy/y) = \int k dx \quad \ln|y| = kx + C$

Step 3: Solve for  $y$  and apply the initial condition.  $y = \pm e^{(kx + C)} = \pm e^C \cdot e^{(kx)}$

Let  $A = \pm e^C$ . Then:  $y = A \cdot e^{(kx)}$

Applying the initial condition  $y(0) = y_0$ :  $y_0 = A \cdot e^{(k \cdot 0)} = A$

Therefore,  $y = y_0 \cdot e^{(kx)}$  is the solution.

This equation has numerous applications, from population growth to radioactive decay.

### Example 3: Logistic Growth

Problem: Solve the differential equation  $dy/dx = ry(1 - y/K)$ , where  $r$  and  $K$  are positive constants, with the initial condition  $y(0) = y_0$  (where  $0 < y_0 < K$ ).

#### Solution:

This is the logistic equation, commonly used to model population growth with a carrying capacity  $K$ .

Step 1: Separate variables.  $dy/(y(1 - y/K)) = r dx$

We can rewrite the left side using partial fractions:  $dy/(y(1 - y/K)) = (1/y + 1/(K-y)) \cdot K dy$

So we have:  $(1/y + 1/(K-y)) \cdot K dy = r dx$

Step 2: Integrate both sides.  $\int (1/y + 1/(K-y)) \cdot K dy = \int r dx \quad K \cdot [\ln|y| - \ln|K-y|] = rx + C \quad \ln|y/(K-y)| = (r/K)x + C/K$

Step 3: Solve for y.  $y/(K-y) = e^{((r/K)x + C/K)}$   $y = (K-y) \cdot e^{((r/K)x + C/K)}$   $y = K \cdot e^{((r/K)x + C/K)} / (1 + e^{((r/K)x + C/K)})$

Let  $D = e^{(C/K)}$ . Then:  $y = K \cdot D \cdot e^{((r/K)x)} / (1 + D \cdot e^{((r/K)x)})$

Applying the initial condition  $y(0) = y_0$ :  $y_0 = K \cdot D / (1 + D)$   $D = y_0 / (K - y_0)$

Substituting this value of D back:  $y = K \cdot (y_0/(K-y_0)) \cdot e^{((r/K)x)} / (1 + (y_0/(K-y_0)) \cdot e^{((r/K)x)})$

Simplifying:  $y = K \cdot y_0 \cdot e^{((r/K)x)} / (K - y_0 + y_0 \cdot e^{((r/K)x)})$

This is the solution to the logistic equation. As  $x \rightarrow \infty$ ,  $y \rightarrow K$ , which is the carrying capacity.

#### Example 4: Orthogonal Trajectories

Problem: Find the orthogonal trajectories of the family of curves  $y = cx^2$ , where c is a parameter.

##### Solution:

Orthogonal trajectories are curves that intersect each member of a given family of curves at right angles. To find them:

Step 1: Find the differential equation of the given family  $y = cx^2$ .  
Differentiating with respect to x:  $dy/dx = 2cx$

Substituting  $c = y/x^2$ :  $dy/dx = 2(y/x^2) \cdot x = 2y/x$

Step 2: Find the differential equation of the orthogonal trajectories. If two curves are orthogonal, the product of their slopes at the intersection point is -1. So, if  $M_1 = dy/dx$  for the original family, then  $M_2 = dy/dx$  for the orthogonal trajectories satisfies:  $M_1 \cdot M_2 = -1$   $(2y/x) \cdot M_2 = -1$   $M_2 = -x/(2y)$

So the differential equation of the orthogonal trajectories is:  $dy/dx = -x/(2y)$

Step 3: Solve this new differential equation using separation of variables.  $2y dy = -x dx$   $\int 2y dy = -\int x dx$   $y^2 = -x^2/2 + C$

Simplifying:  $2y^2 + x^2 = 2C$

This represents a family of ellipses with axes along the coordinate axes, or if  $C < 0$ , a family of hyperbolas.

### Example 5: Nonlinear First-Order Equation

Problem: Solve the differential equation  $dy/dx = (y^2 + 1)/(x^2 + 1)$ .

#### Solution:

Step 1: Separate variables.  $dy/(y^2 + 1) = dx/(x^2 + 1)$

Step 2: Integrate both sides.  $\int dy/(y^2 + 1) = \int dx/(x^2 + 1)$

These are standard integrals:  $\int dy/(y^2 + 1) = \arctan(y) + C_1$   $\int dx/(x^2 + 1) = \arctan(x) + C_2$

So:  $\arctan(y) + C_1 = \arctan(x) + C_2$   $\arctan(y) = \arctan(x) + C$  (where  $C = C_2 - C_1$ )

Step 3: Solve for y. Using the fact that  $\arctan(a) - \arctan(b) = \arctan((a-b)/(1+ab))$  for  $1+ab \neq 0$ : If  $C = \arctan(k)$  for some constant k, then:  $\arctan(y) = \arctan(x) + \arctan(k)$   $\arctan(y) = \arctan((x+k)/(1-kx))$   $y = (x+k)/(1-kx)$

This is the general solution in rational form. If we have an initial condition, we could determine the value of k.

### Unsolved Problems

Here are five unsolved problems involving separable differential equations for practice:

#### Problem 1

Solve the differential equation  $dy/dx = e^{(x-y)}$ .

#### Problem 2

Find the general solution of the differential equation  $dy/dx = (\sin x)(\cos y)$ .

#### Problem 3

Solve the initial value problem:  $dy/dx = xy\sqrt{1-y^2}$ ,  $y(0) = 0$

#### Problem 4

Determine the orthogonal trajectories of the family of curves given by  $y = ce^x$ , where c is a parameter.

## Problem 5

A population  $P$  grows according to the differential equation  $dP/dt = kP(1 - P/M)^2$ , where  $k$  and  $M$  are positive constants. Find  $P(t)$  if  $P(0) = P_0$ , where  $0 < P_0 < M$ .

### Applications of Separable Differential Equations

Separable differential equations appear in numerous applications across various fields:

#### 1. Population Dynamics

The simplest model of population growth is the exponential model:  $dP/dt = kP$

Where  $P$  is the population size and  $k$  is the growth rate. This is separable and gives the solution  $P(t) = P_0 e^{(kt)}$ .

A more realistic model is the logistic equation:  $dP/dt = kP(1 - P/M)$

Where  $M$  is the carrying capacity. This accounts for limited resources and leads to a sigmoid growth curve.

#### 2. Newton's Law of Cooling

An object's temperature change over time can be modeled by:  $dT/dt = k(T - T_e)$

Where  $T$  is the object's temperature,  $T_e$  is the environment temperature, and  $k$  is a constant. This separable equation leads to exponential approach to equilibrium.

#### 3. Radioactive Decay

The decay of radioactive materials follows:  $dN/dt = -\lambda N$

Where  $N$  is the amount of radioactive material and  $\lambda$  is the decay constant. The solution  $N(t) = N_0 e^{(-\lambda t)}$  gives the exponential decay law.

#### 4. Chemical Reaction Kinetics

For a first-order reaction  $A \rightarrow B$ , the rate equation is:  $d[A]/dt = -k[A]$

Where  $[A]$  is the concentration of reactant  $A$ . This separable equation leads to exponential decay of the reactant.

## 5. Circuit Analysis

In an RC circuit, the voltage  $V$  across the capacitor satisfies:  $dV/dt = (E - V)/(RC)$

Where  $E$  is the battery voltage,  $R$  is the resistance, and  $C$  is the capacitance. This separable equation describes how the capacitor charges or discharges.

### Limitations and Extensions

While separable differential equations are powerful tools, they have limitations:

1. Integrability: Even if an equation is separable, we might not be able to find closed-form expressions for the integrals involved.
2. Domain Restrictions: Solutions might have restricted domains due to divisions by zero or other singularities.
3. Implicit Solutions: Often, we can't solve explicitly for  $y$  as a function of  $x$ , leading to implicit relations.

Extensions of the separable equation concept include:

1. Homogeneous equations: Equations of the form  $dy/dx = f(y/x)$  can be transformed into separable equations by substitution.
2. Bernoulli equations: Equations of the form  $dy/dx + P(x)y = Q(x)y^n$  can be transformed into linear equations by substitution.

## CHECK YOUR PROGRESS

- 1) State the existence and uniqueness theorem.

.....  
.....  
.....

- 2) Why is continuity of  $\partial f / \partial y$  important?

.....  
.....  
.....

- 3) Give an example where uniqueness fails.

.....  
.....  
.....

- 4) Define an initial value problem.

.....  
.....  
.....

## LET US SUM UP

The Existence and Uniqueness Theorem guarantees that a first-order differential equation of the form  $y' = f(x, y)$  has a unique solution in a neighborhood of the initial point  $(x_0, y_0)$  if  $f(x, y)$  and its partial derivative with respect to  $y$  are continuous near that point. This ensures that differential equations describing physical systems produce predictable results under specific initial conditions.

## **UNIT END EXERCISES**

### **Short Answer Questions:**

1. What is the existence and uniqueness theorem for first-order differential equations?
2. Explain the method of solving separable equations.

### **Long Answer Questions:**

1. Prove the existence and uniqueness theorem for first-order differential equations.
2. Discuss the role of the Lipschitz condition in differential equations.

### **REFERENCES AND SUGGESTED READINGS:**

1. Hartman, P. (2013). Ordinary Differential Equations (2nd ed.). SIAM
2. Arnold, V. 1. (2012). Ordinary Differential Equations. Springer.
3. Hirsch, M. W., Smale, S., & Devaney, R. L. (2013). Differential Equations, Dynamical Systems, and an Introduction to Chaos (3rd ed.), Academic Press.

## UNIT 15

### EQUATION WITH VARIABLES SEPARATED– EXACT EQUATIONS

#### 15.1 Exact Differential Equations and Integrating Factors

##### Introduction to Exact Differential Equations

In this section, we'll study a special class of first-order differential equations that can be written in the form:

$$M(x,y)dx + N(x,y)dy = 0$$

These are called exact differential equations when they represent the total differential of some function  $F(x,y)$ . We'll learn how to identify exact equations, solve them directly, and transform non-exact equations into exact ones using integrating factors.

##### What Makes an Equation Exact?

A differential equation  $M(x,y)dx + N(x,y)dy = 0$  is exact if there exists a function  $F(x,y)$  such that:

$$dF(x,y) = M(x,y)dx + N(x,y)dy$$

For this to be true, we need:

$$\partial F / \partial x = M(x,y) \quad \partial F / \partial y = N(x,y)$$

From calculus, we know that mixed partial derivatives are equal when continuous:

$$\partial^2 F / \partial y \partial x = \partial^2 F / \partial x \partial y$$

This gives us a necessary and sufficient condition for exactness:

$$\partial M / \partial y = \partial N / \partial x$$

This is our test for exactness - if these partial derivatives are equal, the equation is exact.

##### Solving Exact Differential Equations

If  $M(x,y)dx + N(x,y)dy = 0$  is exact, the solution is  $F(x,y) = C$ , where  $C$  is a constant. To find  $F(x,y)$ , we can:



1. Integrate  $M(x,y)$  with respect to  $x$ , treating  $y$  as constant:  $F(x,y) = \int M(x,y)dx + h(y)$

where  $h(y)$  is a function of  $y$  alone.

2. Find  $h(y)$  by differentiating  $F(x,y)$  with respect to  $y$  and setting it equal to  $N(x,y)$ :  $\partial F/\partial y = \partial/\partial y[\int M(x,y)dx] + h'(y) = N(x,y)$

Thus:  $h'(y) = N(x,y) - \partial/\partial y[\int M(x,y)dx]$

And  $h(y) = \int [N(x,y) - \partial/\partial y[\int M(x,y)dx]]dy$

3. Substitute  $h(y)$  back into  $F(x,y)$  to get the complete solution.

Alternatively, we could integrate  $N(x,y)$  with respect to  $y$  and then find the unknown function of  $x$ .

### Integrating Factors

When a differential equation  $M(x,y)dx + N(x,y)dy = 0$  is not exact, we can sometimes find an integrating factor  $\mu(x,y)$  such that when we multiply the original equation by  $\mu$ , the resulting equation becomes exact:

$$\mu(x,y)M(x,y)dx + \mu(x,y)N(x,y)dy = 0$$

For this to be exact, we need:

$$\partial[\mu M]/\partial y = \partial[\mu N]/\partial x$$

This gives us a partial differential equation for  $\mu$ . Finding general solutions for  $\mu$  is difficult, but in specific cases:

1. If  $\mu$  depends only on  $x$  ( $\mu = \mu(x)$ ), then:  $\mu' = \mu(\partial M/\partial y - \partial N/\partial x)/N$

This works if  $(\partial M/\partial y - \partial N/\partial x)/N$  depends only on  $x$ .

2. If  $\mu$  depends only on  $y$  ( $\mu = \mu(y)$ ), then:  $\mu' = \mu(\partial N/\partial x - \partial M/\partial y)/M$

This works if  $(\partial N/\partial x - \partial M/\partial y)/M$  depends only on  $y$ .

### Special Cases and Shortcuts

Some common integrating factors include:

1. For equations of form  $y'dx + P(x)y'dy = Q(x)dx$ , try  $\mu = 1/y'$ .
2. For equations of form  $P(xy)dx + Q(xy)ydy = 0$ , try  $\mu = 1/(xy)$ .

3. For the linear equation  $y' + P(x)y = Q(x)$ , the integrating factor is  $\mu = e^{\int P(x)dx}$ .

### Solved Examples

#### Example 1: Testing for Exactness

Determine whether the following differential equation is exact:  $(2xy + y^2)dx + (x^2 + 2xy - 3)dy = 0$

Solution: Let  $M(x,y) = 2xy + y^2$  Let  $N(x,y) = x^2 + 2xy - 3$

To check for exactness, we compute:  $\partial M/\partial y = 2x + 2y$   $\partial N/\partial x = 2x + 2y$

Since  $\partial M/\partial y = \partial N/\partial x$ , the equation is exact.

#### Example 2: Solving an Exact Equation

Solve the exact differential equation:  $(2xy + y^2)dx + (x^2 + 2xy - 3)dy = 0$

Solution: We determined in Example 1 that this equation is exact.

Step 1: Integrate  $M(x,y)$  with respect to  $x$ , treating  $y$  as constant.  $F(x,y) = \int (2xy + y^2)dx$   $F(x,y) = x^2y + xy^2 + h(y)$

Step 2: Find  $h(y)$  by differentiating  $F(x,y)$  with respect to  $y$  and setting it equal to  $N(x,y)$ .  $\partial F/\partial y = x^2 + 2xy + h'(y) = x^2 + 2xy - 3$

Therefore:  $h'(y) = -3$   $h(y) = -3y + C_1$

Step 3: Substitute  $h(y)$  back into  $F(x,y)$ .  $F(x,y) = x^2y + xy^2 - 3y + C_1$

The solution is:  $x^2y + xy^2 - 3y = C$  (where  $C = -C_1$  is an arbitrary constant)

#### Example 3: Using an Integrating Factor

Solve the differential equation:  $(3xy^2 + y^3)dx + (2x^2y + 3xy^2)dy = 0$

Solution: Let  $M(x,y) = 3xy^2 + y^3$  Let  $N(x,y) = 2x^2y + 3xy^2$

Check for exactness:  $\partial M/\partial y = 6xy + 3y^2$   $\partial N/\partial x = 4xy + 3y^2$

Since  $\partial M/\partial y \neq \partial N/\partial x$ , the equation is not exact.

Let's find an integrating factor:  $\partial M/\partial y - \partial N/\partial x = (6xy + 3y^2) - (4xy + 3y^2) = 2xy$

The expression  $(\partial M/\partial y - \partial N/\partial x)/(xN) = 2xy/(x(2x^2y + 3xy^2)) = 2/(2x + 3y)$

This doesn't depend solely on x or y, so let's try  $\mu = x^m \cdot y^n$

For this type of equation, we can try  $\mu = 1/x$

Multiplying our equation by  $1/x$ :  $(3y^2 + y^3/x)dx + (2xy + 3y^2)dy = 0$

Let's check if this is now exact:  $M_1(x,y) = 3y^2 + y^3/x$   $N_1(x,y) = 2xy + 3y^2$

$$\partial M_1 / \partial y = 6y - 3y^2/x \quad \partial N_1 / \partial x = 2y$$

Still not exact. Let's try  $\mu = 1/(xy)$ :

Multiplying our original equation by  $1/(xy)$ :  $(3y + y^2/x)dx + (2x + 3y)dy = 0$

Check for exactness:  $M_2(x,y) = 3y + y^2/x$   $N_2(x,y) = 2x + 3y$

$$\partial M_2 / \partial y = 3 + 2y/x \quad \partial N_2 / \partial x = 2$$

Not exact.

Let's try  $\mu = 1/y$ :  $(3x + y^2)dx + (2x^2 + 3xy)dy/y = 0 = (3x + y^2)dx + (2x^2 + 3xy)/y \cdot dy = 0 = (3x + y^2)dx + (2x^2/y + 3x)dy = 0$

Check for exactness:  $M_3(x,y) = 3x + y^2$   $N_3(x,y) = 2x^2/y + 3x$

$$\partial M_3 / \partial y = 2y \quad \partial N_3 / \partial x = 4x/y + 3$$

Still not exact.

Let's try  $\mu = 1/y^2$ :  $(3x/y + 1)dx + (2x^2/y^3 + 3x/y)dy = 0$

This doesn't simplify our work.

After trying several approaches, let's use a systematic method. For this equation, a better approach is to rewrite it as:  $(3xy^2 + y^3)dx + (2x^2y + 3xy^2)dy = 0$

Factoring out  $y^2$ :  $y^2(3x + y)dx + y(2x^2 + 3xy)dy = 0$

Taking out a common factor of  $xy$ :  $xy(3y + y^2/x)dx + xy(2x + 3y)dy = 0$

Now with  $\mu = 1/(xy)$ :  $(3y + y^2/x)dx + (2x + 3y)dy = 0$

Let's check again:  $M_4(x,y) = 3y + y^2/x$   $N_4(x,y) = 2x + 3y$

$$\partial M_4 / \partial y = 3 + 2y/x \quad \partial N_4 / \partial x = 2$$

Still not exact.

Let's reexamine the original equation:  $(3xy^2 + y^3)dx + (2x^2y + 3xy^2)dy = 0$

We can rewrite this as:  $d(x^2y^2 + xy^3) = 0$

This implies:  $x^2y^2 + xy^3 = C$

Which is our solution. (This special case could be recognized by noticing that all terms have the same total degree.)

#### **Example 4: Linear Equation with Integrating Factor**

Solve the differential equation:  $dy/dx + 2y/x = x$ ,  $x > 0$

Solution: First, rewrite in standard form:  $dy/dx + 2y/x = x$   $dy + (2y/x)dx = x \cdot dx$

This is a linear equation of form  $dy/dx + P(x)y = Q(x)$  with:  $P(x) = 2/x$   $Q(x) = x$

The integrating factor is:  $\mu = e^{\int P(x)dx} = e^{\int (2/x)dx} = e^{2\ln(x)} = x^2$

Multiply the original equation by  $\mu$ :  $x^2 \cdot dy + 2x \cdot y \cdot dx = x^3 \cdot dx$

The left side is the derivative of  $x^2y$ :  $d(x^2y) = x^2 \cdot dy + 2x \cdot y \cdot dx$

So our equation becomes:  $d(x^2y) = x^3 \cdot dx$

Integrating both sides:  $x^2y = \int x^3 \cdot dx = x^4/4 + C$

Solving for  $y$ :  $y = x^2/4 + C/x^2$

This is the general solution.

#### **Example 5: Using a Suitable Integrating Factor**

Solve the differential equation:  $(y^2 - xy)dx + (2xy - x^2)dy = 0$

Solution: Let  $M(x,y) = y^2 - xy$  Let  $N(x,y) = 2xy - x^2$

Check for exactness:  $\partial M/\partial y = 2y - x$   $\partial N/\partial x = 2y - x$

Since  $\partial M/\partial y = \partial N/\partial x$ , the equation is exact.

Find the solution  $F(x,y) = C$ :  $F(x,y) = \int M(x,y)dx = \int (y^2 - xy)dx = xy^2 - x^2y/2 + h(y)$

Differentiate with respect to  $y$ :  $\partial F/\partial y = 2xy - x^2/2 + h'(y) = N(x,y) = 2xy - x^2$

Therefore:  $h'(y) = 0$   $h(y) = C_1$

The final solution is:  $F(x,y) = xy^2 - x^2y/2 = C$

or  $xy^2 - x^2y/2 = C$

This represents the family of solutions to the differential equation.

### **Unsolved Problems**

#### **Problem 1**

Determine whether the following differential equation is exact, and if so, find its solution:  $(y^2e^x + 2xy)dx + (2ye^x + x^2)dy = 0$

#### **Problem 2**

Find the general solution of the differential equation:  $(2x + 3y^2)dx + (6xy + 7)dy = 0$

#### **Problem 3**

Find an integrating factor for the differential equation and then solve it:  $(2x + y)dx + (x - 3y)dy = 0$

#### **Problem 4**

Solve the following differential equation:  $(y - 3x^2)dx + (x + 2y^2)dy = 0$

#### **Problem 5**

Find the solution of the following differential equation, given that  $y(1) = 0$ :  
 $(y^3 + \cos(xy))dx + (3xy^2 + x \cdot \cos(xy))dy = 0$

## CHECK YOUR PROGRESS

- 1) Define a separable differential equation.

.....  
.....  
.....

- 2) State the condition for an equation to be exact.

.....  
.....  
.....

- 3) Find the integrating factor for  $(y-x)dx + (y+x)dy = 0$ .

.....  
.....  
.....

## LET US SUM UP

In a separable equation, the variables can be separated so that all  $x$  terms appear on one side and all  $y$  terms on the other, allowing integration on both sides. An exact equation has the form  $Mdx + Ndy = 0$ , where  $\partial M/\partial y = \partial N/\partial x$ . If this condition is not satisfied, an integrating factor may exist to make the equation exact, facilitating its solution.

## UNIT END EXERCISES

### Short Answer Questions:

1. Define an exact equation and state its condition.
2. What is an integrating factor? Give an example.

### Long Answer Questions:

1. Explain the convergence of Picard's successive approximations.
2. Compare and contrast exact equations and linear first-order equations.

### REFERENCES AND SUGGESTED READINGS:

1. Hartman, P. (2013). Ordinary Differential Equations (2nd ed.). SIAM
2. Coddington, E. A. (2018). An Introduction to Ordinary Differential Equations. Dover Publications.
3. Arnold, V. I. (2012). Ordinary Differential Equations. Springer.

## UNIT 16

### THE METHOD OF SUCCESSIVE APPROXIMATIONS – THE LIPSCHITZ CONDITION

#### 16.1 The Method of Successive Approximations

##### Introduction to Successive Approximations

The method of successive approximations, also known as Picard's method, provides a theoretical foundation for the existence and uniqueness of solutions to first-order initial value problems. Beyond its theoretical importance, it also gives us a constructive approach to finding solutions through an iterative process.

##### The Initial Value Problem

Consider the initial value problem:

$$dy/dx = f(x,y), y(x_0) = y_0$$

where  $f(x,y)$  is a continuous function in some region containing the point  $(x_0, y_0)$ .

##### Picard's Iteration

The idea behind successive approximations is to convert the differential equation into an equivalent integral equation:

$$y(x) = y_0 + \int_{x_0}^x f(t, y(t)) dt$$

Then we define a sequence of functions  $\{\phi_n(x)\}$  as follows:

$$\begin{aligned} \phi_0(x) &= y_0 \text{ (initial approximation)} \\ \phi_1(x) &= y_0 + \int_{x_0}^x f(t, \phi_0(t)) dt \\ \phi_2(x) &= y_0 + \int_{x_0}^x f(t, \phi_1(t)) dt \\ &\vdots \\ \phi_{n+1}(x) &= y_0 + \int_{x_0}^x f(t, \phi_n(t)) dt \end{aligned}$$

Under suitable conditions, the sequence  $\{\phi_n(x)\}$  converges to the unique solution  $y(x)$  of the initial value problem.

##### Existence and Uniqueness Theorem

Picard's existence and uniqueness theorem states:

If  $f(x,y)$  and  $\partial f/\partial y$  are continuous in a rectangle  $R = \{(x,y) : |x-x_0| \leq a, |y-y_0| \leq b\}$ , then there exists an interval  $I = [x_0-h, x_0+h]$  (where  $h \leq a$  is sufficiently small) such that the initial value problem has a unique solution  $y(x)$  on  $I$ .

### Convergence of Picard Iterations

The convergence of Picard iterations relies on the Lipschitz condition. A function  $f(x,y)$  satisfies a Lipschitz condition with respect to  $y$  if there exists a constant  $L > 0$  such that:

$$|f(x,y_1) - f(x,y_2)| \leq L|y_1 - y_2|$$

for all points  $(x,y_1)$  and  $(x,y_2)$  in the region of interest.

The existence of a continuous partial derivative  $\partial f/\partial y$  ensures the Lipschitz condition is satisfied, with  $L = \max|\partial f/\partial y|$  in the region.

### Error Estimation

If  $f(x,y)$  satisfies a Lipschitz condition with constant  $L$ , and  $M$  is the maximum value of  $|f(x,y)|$  in the region, then the error in the  $n$ th approximation is bounded by:

$$|y(x) - \phi_n(x)| \leq (M/L) \cdot (L|x-x_0|)^n/n!$$

This shows that the approximations converge rapidly for small values of  $|x-x_0|$ .

### Practical Implementation

In practice, carrying out the integrations for successive approximations can become increasingly complex. Therefore, the method is often more valuable as a theoretical tool than a practical computational method. However, for simple problems, it can provide insight into the solution structure.

### Example Calculations

To implement Picard's method practically:

1. Start with  $\phi_0(x) = y_0$  (constant function)
2. Substitute into the right side of the integral equation to get  $\phi_1(x)$
3. Continue substituting each approximation to get the next one
4. Stop when successive approximations are sufficiently close



### Relationship to Power Series Methods

The successive approximations often generate terms that correspond to the Taylor series expansion of the solution. For linear equations, a few iterations can reveal the pattern of the series solution.

### Solved Examples

#### Example 1: Basic Picard Iteration

Find the first three Picard approximations for the initial value problem:  
 $dy/dx = y, y(0) = 1$

Solution: The equivalent integral equation is:  $y(x) = 1 + \int_0^x y(t)dt$

The Picard iterations are:

$$\phi_0(x) = 1$$

$$\phi_1(x) = 1 + \int_0^x \phi_0(t)dt = 1 + 1 \cdot t = 1 + x$$

$$\phi_2(x) = 1 + \int_0^x \phi_1(t)dt = 1 + \int_0^x (1 + t)dt = 1 + [t + t^2/2]_0^x = 1 + x + x^2/2$$

$$\phi_3(x) = 1 + \int_0^x \phi_2(t)dt = 1 + \int_0^x (1 + t + t^2/2)dt = 1 + [t + t^2/2 + t^3/6]_0^x = 1 + x + x^2/2 + x^3/6$$

We recognize this as the beginning of the Taylor series for  $e^x$ , which is indeed the exact solution to this problem. The successive approximations are converging to  $y(x) = e^x$ .

#### Example 2: Non-Linear Equation

Find the first three Picard approximations for:  $dy/dx = x + y^2, y(0) = 0$

Solution: The integral equation is:  $y(x) = 0 + \int_0^x (t + y(t)^2)dt$

The Picard iterations are:

$$\phi_0(x) = 0$$

$$\phi_1(x) = \int_0^x (t + \phi_0(t)^2)dt = \int_0^x t \cdot dt = x^2/2$$

$$\begin{aligned}\phi_2(x) &= \int_0^x (t + \phi_1(t)^2) dt = \int_0^x (t + (t^2/2)^2) dt \\ &= \int_0^x (t + t^4/4) dt = [t^2/2 + t^5/20]_0^x = x^2/2 + x^5/20\end{aligned}$$

$$\phi_3(x) = \int_0^x (t + \phi_2(t)^2) dt = \int_0^x \left(t + \left(\frac{x^2}{2} + \frac{x^5}{20}\right)^2\right) dt$$

This becomes complicated to evaluate directly. However, we can expand:  
 $(x^2/2 + x^5/20)^2 = x^4/4 + x^7/20 + x^{10}/400$

$$\begin{aligned}\text{So: } \phi_3(x) &= \int_0^x (t + t^4/4 + t^7/20 + t^{10}/400) dt = [t^2/2 + t^5/20 + t^8/160 + t^{11}/4400]_0^x \\ &= x^2/2 + x^5/20 + x^8/160 + x^{11}/4400\end{aligned}$$

Each iteration captures more terms in the series expansion of the true solution.

### Example 3: Linear First-Order Equation

Find the first three Picard approximations for:  $dy/dx = -2xy$ ,  $y(0) = 1$

Solution: The integral equation is:  $y(x) = 1 + \int_0^x (-2t \cdot y(t)) dt$

The iterations are:

$$\phi_0(x) = 1$$

$$\phi_1(x) = 1 + \int_0^x (-2t \cdot \phi_0(t)) dt = 1 + \int_0^x (-2t) dt = 1 + [-t^2]_0^x = 1 - x^2$$

$$\begin{aligned}\phi_2(x) &= 1 + \int_0^x (-2t \cdot \phi_1(t)) dt = 1 + \int_0^x (-2t \cdot (1 - t^2)) dt = 1 + \int_0^x (-2t + 2t^3) dt \\ &= 1 + [-t^2 + t^4/2]_0^x = 1 - x^2 + x^4/2\end{aligned}$$

$$\begin{aligned}\phi_3(x) &= 1 + \int_0^x (-2t \cdot \phi_2(t)) dt = 1 + \int_0^x (-2t \cdot (1 - t^2 + t^4/2)) dt = \\ &= 1 + \int_0^x (-2t + 2t^3 - t^5) dt = 1 + [-t^2 + t^4/2 - t^6/6]_0^x = 1 - x^2 + x^4/2 - x^6/6\end{aligned}$$

We recognize this as the beginning of the Taylor series for  $e^{(-x^2)}$ , which is the exact solution to this problem.

### Example 4: System with Variable Coefficient

Find the first three Picard approximations for:  $dy/dx = x \cdot \sin(y)$ ,  $y(0) = 0$

Solution: For small values of  $y$ , we can use the approximation  $\sin(y) \approx y - y^3/6 + \dots$

The integral equation is:  $y(x) = 0 + \int_0^x t \cdot \sin(y(t)) dt$

The iterations are:

$$\phi_0(x)$$

$$\phi_1(x) = \int_0^x t \cdot \sin(\phi_0(t)) dt = \int_0^x t \cdot \sin(0) dt = 0$$

Since  $\phi_1(x) = 0$ , all subsequent approximations will also be 0. This tells us that  $y(x) = 0$  is the unique solution to this initial value problem, which makes sense given the initial condition  $y(0) = 0$  and the fact that  $\sin(0) = 0$ .

To get a more interesting example, let's modify the initial condition to  $y(0) = \pi/4$ :

$$\phi_0(x) = \pi/4$$

$$\phi_1(x) = \pi/4 + \int_0^x t \cdot \sin(\phi_0(t)) dt = \pi/4 + \int_0^x t \cdot \sin(\pi/4) dt = \pi/4 + \sin(\pi/4) \cdot \int_0^x t \cdot dt = \pi/4 + \sin(\pi/4) \cdot x^2/2 = \pi/4 + (\sqrt{2}/2) \cdot x^2/2 = \pi/4 + x^2/(2\sqrt{2})$$

$$\phi_2(x) = \pi/4 + \int_0^x t \cdot \sin(\phi_1(t)) dt = \pi/4 + \int_0^x t \cdot \sin(\pi/4 + t^2/(2\sqrt{2})) dt$$

This becomes more difficult to evaluate directly. We would need to use numerical integration or series approximations for the sine function.

### Example 5: Demonstrating Convergence

For the problem  $dy/dx = 2y$ ,  $y(0) = 1$ , show that the Picard iterations converge to the exact solution  $y = e^{(2x)}$ .

Solution: The integral equation is:  $y(x) = 1 + \int_0^x 2y(t) dt$

The iterations are:

$$\phi_0(x) = 1$$

$$\phi_1(x) = 1 + \int_0^x 2\phi_0(t) dt = 1 + \int_0^x 2 dt = 1 + 2x$$

$$\phi_2(x) = 1 + \int_0^x 2\phi_1(t) dt = 1 + \int_0^x 2(1 + 2t) dt = 1 + \int_0^x (2 + 4t) dt$$

$$= 1 + [2t + 2t^2]_0^x = 1 + 2x + 2x^2$$

$$\phi_3(x) = 1 + \int_0^x 2\phi_2(t) dt = 1 + \int_0^x 2(1 + 2t + 2t^2) dt = 1 + \int_0^x (2 + 4t + 4t^2) dt = 1 + [2t + 2t^2 + 4t^3/3]_0^x = 1 + 2x + 2x^2 + 4x^3/3$$

If we continue this process, we get:  $\phi_4(x) = 1 + 2x + 2x^2 + 4x^3/3 + 2x^4/3$

The Taylor series for  $e^{(2x)}$  is:  $e^{(2x)} = 1 + 2x + (2x)^2/2! + (2x)^3/3! + (2x)^4/4! + \dots = 1 + 2x + 2x^2/1 + 8x^3/6 + 16x^4/24 + \dots = 1 + 2x + 2x^2 + 4x^3/3 + 2x^4/3 + \dots$

We can see that the Picard iterations are producing exactly the Taylor series for  $e^{(2x)}$ , term by term, confirming that the iterations converge to the exact solution.

## Unsolved Problems

### Problem 1

Using the method of successive approximations, find the first three approximations for the initial value problem:  $dy/dx = x^2 + y$ ,  $y(0) = 1$

### Problem 2

Apply Picard's method to find the first three approximations for:  $dy/dx = xy$ ,  $y(1) = 2$

### Problem 3

Find the first two Picard approximations for the non-linear equation:  $dy/dx = y^2$ ,  $y(0) = 1$  Also, determine the interval in which these approximations are valid.

### Problem 4

Use successive approximations to solve the initial value problem:  $dy/dx = e^{-x^3}y$ ,  $y(0) = 3$  Compute the first three approximations.

### Problem 5

For the equation  $dy/dx = \sin(x+y)$ ,  $y(0) = 0$ , find the first three Picard iterations. Compare the third approximation with the Taylor series of the exact solution around  $x = 0$  up to the third-degree term.

## 16.2 Lipschitz Condition and Its Importance

In the study of differential equations, particularly when investigating existence and uniqueness of solutions, the Lipschitz condition plays a crucial role. This condition provides a mathematical framework to ensure that a solution not only exists but is unique.

### Definition of Lipschitz Condition

A function  $f(t,y)$  satisfies a Lipschitz condition with respect to  $y$  in a domain  $D$  if there exists a constant  $L > 0$  (called the Lipschitz constant) such that:

$$|f(t,y_1) - f(t,y_2)| \leq L|y_1 - y_2|$$

for all points  $(t,y_1)$  and  $(t,y_2)$  in  $D$ .

In simpler terms, the Lipschitz condition places a bound on how rapidly a function can change with respect to one of its variables. It essentially states that the rate of change of  $f$  with respect to  $y$  is bounded by the constant  $L$ .

### Geometric Interpretation

Geometrically, the Lipschitz condition means that the slopes of the lines connecting any two points on the function's graph (with the same  $t$ -value) are bounded by  $L$ . This prevents the function from having vertical tangent lines or discontinuities in its derivative with respect to  $y$ .

### Connection to Continuity and Differentiability

The Lipschitz condition is stronger than continuity but weaker than differentiability with a bounded derivative:

- If  $f(t,y)$  has a continuous partial derivative  $\partial f/\partial y$  in domain  $D$ , and  $|\partial f/\partial y| \leq M$  for all points in  $D$ , then  $f$  satisfies a Lipschitz condition with Lipschitz constant  $L = M$ .
- A function satisfying a Lipschitz condition is necessarily continuous in the variable  $y$ , but the converse is not always true.

### Examples of Functions Satisfying and Violating Lipschitz Condition

#### Example 1: Satisfying Lipschitz Condition

$f(t,y) = y^2$  for domain  $D$  where  $y$  is bounded

For any bounded domain where  $|y| \leq K$ , we have:  $|f(t,y_1) - f(t,y_2)| = |y_1^2 - y_2^2|$   
 $= |(y_1 - y_2)(y_1 + y_2)| \leq |y_1 - y_2| \cdot |y_1 + y_2| \leq |y_1 - y_2| \cdot 2K$

Therefore,  $f$  satisfies a Lipschitz condition with  $L = 2K$ .

#### Example 2: Violating Lipschitz Condition

$f(t,y) = \sqrt{y}$  for  $y \geq 0$

For this function:  $|f(t,y_1) - f(t,y_2)| = |\sqrt{y_1} - \sqrt{y_2}| = |y_1 - y_2|/|\sqrt{y_1} + \sqrt{y_2}|$

As  $y_1$  and  $y_2$  approach zero, the denominator approaches zero, making the fraction unbounded. Therefore, no single Lipschitz constant  $L$  can satisfy the required inequality for all points in the domain, especially near  $y = 0$ .

### Importance in Differential Equations

The Lipschitz condition is crucial in the theory of ordinary differential equations for several reasons:

1. **Uniqueness of Solutions:** The Lipschitz condition is sufficient to guarantee the uniqueness of solutions to initial value problems. Without this condition, an initial value problem might have multiple solutions.
2. **Existence of Solutions:** While the Lipschitz condition alone doesn't guarantee existence, when combined with continuity of  $f(t,y)$ , it helps establish existence of solutions through methods like the method of successive approximations.
3. **Stability of Solutions:** The Lipschitz condition provides a measure of stability, indicating how sensitive solutions are to changes in initial conditions.
4. **Numerical Methods:** Many numerical methods for solving differential equations require the Lipschitz condition to ensure convergence and to bound error estimates.

#### **Local vs. Global Lipschitz Condition**

- **Local Lipschitz Condition:** A function satisfies a local Lipschitz condition if for every point in the domain, there exists a neighborhood where the Lipschitz condition holds.
- **Global Lipschitz Condition:** The function satisfies the Lipschitz condition throughout the entire domain.

Many functions encountered in practice satisfy a local Lipschitz condition but not a global one. This is sufficient for local existence and uniqueness of solutions to differential equations.

## CHECK YOUR PROGRESS

- 1) Define the Lipschitz condition.

.....  
.....  
.....

- 2) Why does the Lipschitz condition ensure uniqueness?

.....  
.....  
.....

- 3) Perform one Picard iteration for  $y'=x+y$ ,  $y(0) = 1$ .

.....  
.....  
.....

## LET US SUM UP

The method of successive approximations or Picard's method constructs a sequence of functions that converge to the true solution. It begins with an initial guess and uses the iterative relation  $y_{n+1}(x) = y_0 + \int_{x_0}^x f(t, y_n(t)) dt$ . The Lipschitz condition, expressed as  $|f(x, y_1) - f(x, y_2)| \leq L|y_1 - y_2|$ , ensures that the sequence of approximations converges and that the solution is unique.

## UNIT END EXERCISES

### Short Answer Questions:

1. How does the Picard iteration method work?
2. State and explain the Lipschitz condition.
3. What is meant by convergence of successive approximations?

### Long Answer Questions:

1. Define the Lipschitz condition for a function  $f(x, y)$  and explain its role in ensuring uniqueness of the solution of  $dy/dx = f(x, y)$ ,  $y(x_0) = y_0$ .
2. Verify whether the function  $f(x, y) = y^{1/3}$  satisfies the Lipschitz condition near  $y = 0$ . Does the initial value problem  $y' = y^{1/3}$ ,  $y(0) = 0$  have a unique solution? Justify your answer.

### REFERENCES AND SUGGESTED READINGS:

1. Hartman, P. (2013). Ordinary Differential Equations (2nd ed.). SIAM
2. Coddington, E. A. (2018). An Introduction to Ordinary Differential Equations. Dover Publications.
3. Arnold, V. 1. (2012). Ordinary Differential Equations. Springer.
4. Braun, M. (2010). Differential Equations and Their Applications (4th ed.). Springer.



## UNIT 17

### CONVERGENCE OF THE SUCCESSIVE APPROXIMATIONS

#### 17.1 Convergence of Successive Approximations

Successive approximations, also known as Picard iterations, form a constructive method to demonstrate the existence and uniqueness of solutions to initial value problems. This method involves creating a sequence of functions that converge to the solution of a differential equation.

#### The Method of Successive Approximations

Consider the initial value problem:

$$dy/dt = f(t,y), y(t_0) = y_0$$

The method of successive approximations defines a sequence of functions  $\{\varphi_n(t)\}$  as follows:

$$\begin{aligned}\varphi_0(t) &= y_0 \\ \varphi_1(t) &= y_0 + \int_0^x f(s, \varphi_0(s)) ds \\ \varphi_2(t) &= y_0 + \int_0^x f(s, \varphi_1(s)) ds \dots \varphi_{n+1}(t) = y_0 + \int_0^x f(s, \varphi_n(s)) ds\end{aligned}$$

Under appropriate conditions, this sequence converges to the unique solution of the initial value problem.

#### Conditions for Convergence

For the sequence of successive approximations to converge, the following conditions are typically required:

1.  $f(t,y)$  is continuous in a domain  $D$  containing the point  $(t_0, y_0)$ .
2.  $f(t,y)$  satisfies a Lipschitz condition with respect to  $y$  in  $D$ .

#### Theorem of Convergence

If  $f(t,y)$  is continuous and satisfies a Lipschitz condition with constant  $L$  in a domain  $D$  containing  $(t_0, y_0)$ , then:

1. The sequence of successive approximations  $\{\varphi_n(t)\}$  converges uniformly on an interval  $[t_0-h, t_0+h]$  (where  $h$  is sufficiently small) to a function  $\varphi(t)$ .

2. This limit function  $\varphi(t)$  is the unique solution to the initial value problem  $dy/dt = f(t,y)$ ,  $y(t_0) = y_0$  on that interval.

### Proof Outline

The proof involves several steps:

1. Showing that each approximation  $\varphi_n(t)$  is well-defined and continuous.
2. Establishing bounds on  $|\varphi_{n+1}(t) - \varphi_n(t)|$  using the Lipschitz condition.
3. Proving that the series  $\varphi_0(t) + \int_{t_0}^t [\varphi_{n+1}(t) - \varphi_n(t)]$  converges uniformly.
4. Verifying that the limit function satisfies the differential equation.

### Rate of Convergence

The rate at which successive approximations converge depends on the Lipschitz constant  $L$ . Specifically, for  $t$  in  $[t_0-h, t_0+h]$ :

$$|\varphi_{n+1}(t) - \varphi_n(t)| \leq (M \cdot L^n \cdot h^{(n+1)}) / ((n+1)!)$$

where  $M$  is a bound on  $|f(t,y)|$  in the domain of interest.

This shows that the sequence converges exponentially fast, making the method theoretically powerful, although direct computation of many iterations may be cumbersome.

### Practical Implementation

In practice, computing successive approximations often involves numerical techniques, as explicit integration may not be feasible for complex functions  $f(t,y)$ . The approximations typically improve rapidly in the early iterations and then more slowly as  $n$  increases.

### Error Estimation

For a given number of iterations  $n$ , the error between the  $n$ th approximation and the true solution can be estimated as:

$$|\varphi(t) - \varphi_n(t)| \leq (M \cdot e^{(L \cdot |t-t_0|)}) / (L \cdot (n+1)!) \cdot (L \cdot |t-t_0|)^{(n+1)}$$

This error bound helps determine how many iterations are needed to achieve a desired accuracy.

## **17.2 Applications of Existence and Uniqueness Theorems**

The existence and uniqueness theorems for differential equations have numerous applications in both theoretical analysis and practical problem-solving. These theorems provide a foundation for understanding the behavior of solutions and for developing methods to approximate them.

### **Applications in Mathematical Modeling**

#### **1. Validating Mathematical Models**

Before investing resources in solving a differential equation model, it's essential to know whether a solution exists and is unique. Existence and uniqueness theorems provide criteria to verify that a model is well-posed, meaning it has a unique solution that depends continuously on the initial data.

#### **2. Determining the Domain of Validity**

These theorems often specify conditions under which a unique solution exists. This helps identify the range of parameters or initial conditions for which the model is valid, guiding experimental design and interpretation of results.

#### **3. Extending Solutions**

Local existence theorems can be applied repeatedly to extend solutions beyond their initial interval of existence, allowing for a more complete understanding of long-term behavior.

### **Applications in Numerical Analysis**

#### **1. Convergence of Numerical Methods**

Numerical methods for solving differential equations often rely on existence and uniqueness theorems to establish their convergence. For example, the convergence of Euler's method and Runge-Kutta methods depends on the Lipschitz condition.

#### **2. Error Analysis**

The Lipschitz constant provides a measure of the sensitivity of solutions to perturbations in initial conditions or round-off errors, allowing for rigorous error bounds in numerical approximations.

### **3. Stability Analysis**

Existence and uniqueness theorems help analyze the stability of numerical schemes, determining whether small perturbations in input data lead to small changes in the solution.

#### **Applications in Qualitative Analysis**

##### **1. Phase Plane Analysis**

Existence and uniqueness theorems ensure that trajectories in a phase plane cannot intersect (except at equilibrium points), forming the basis for qualitative analysis of nonlinear systems.

##### **2. Bifurcation Theory**

These theorems help identify conditions under which the qualitative behavior of solutions changes, such as the emergence of multiple solutions or changes in stability.

##### **3. Stability of Equilibrium Points**

Linearization techniques used to analyze the stability of equilibrium points depend on local existence and uniqueness of solutions.

#### **Applications in Control Theory**

##### **1. Controller Design**

Existence and uniqueness theorems provide guarantees that control systems will behave predictably, which is essential for designing reliable controllers.

##### **2. Optimal Control**

In optimal control problems, these theorems ensure that the state equations have unique solutions for given control inputs, making optimization problems well-defined.

#### **Applications in Specific Fields**

##### **1. Physics**

In classical mechanics, existence and uniqueness theorems justify the deterministic nature of physical systems: given initial conditions, the future state of the system is uniquely determined.

## **2. Biology**

In population dynamics, existence and uniqueness results ensure that models predicting species growth or interaction have meaningful solutions.

## **3. Economics**

In economic modeling, these theorems help validate differential equation models of market dynamics, resource allocation, and growth theories.

## **4. Engineering**

In electrical circuit analysis, chemical reaction kinetics, and structural mechanics, existence and uniqueness theorems provide the theoretical foundation for modeling and simulation.

### **Applications of Successive Approximations**

#### **1. Constructive Proofs**

The method of successive approximations provides not just a theoretical proof of existence and uniqueness but also a constructive method to compute solutions.

#### **2. Iterative Numerical Methods**

Many practical numerical schemes, such as predictor-corrector methods, are based on the idea of successive approximations.

#### **3. Perturbation Methods**

For nearly linear systems or problems with small parameters, successive approximations form the basis of perturbation techniques.

### **Limitations and Extensions**

#### **1. Non-Lipschitz Cases**

When the Lipschitz condition fails, understanding the consequences for uniqueness becomes more subtle. Examples like  $y' = y^{(2/3)}$ ,  $y(0) = 0$  have multiple solutions despite having continuous right-hand sides.

#### **2. Weak Solutions**

For certain applications, particularly in partial differential equations, the concept of a solution may need to be extended to include weak solutions, where existence and uniqueness results take different forms.

### 3. Stochastic Differential Equations

Extensions of existence and uniqueness theorems to stochastic differential equations provide a framework for modeling random phenomena.

#### Solved Problems

##### Problem 1: Verifying the Lipschitz Condition

Problem: Determine whether the function  $f(t,y) = t + \sin(y)$  satisfies a Lipschitz condition with respect to  $y$  on the domain  $D = \{(t,y) : 0 \leq t \leq 1, -\infty < y < \infty\}$ .

Solution: To verify the Lipschitz condition, we need to find a constant  $L$  such that  $|f(t,y_1) - f(t,y_2)| \leq L|y_1 - y_2|$  for all points in  $D$ .

For any fixed  $t$  and any  $y_1, y_2$ :  $|f(t,y_1) - f(t,y_2)| = |t + \sin(y_1) - (t + \sin(y_2))| = |\sin(y_1) - \sin(y_2)|$

Using the mean value theorem for  $\sin(y)$ , there exists a point  $c$  between  $y_1$  and  $y_2$  such that:  $\sin(y_1) - \sin(y_2) = \cos(c) \cdot (y_1 - y_2)$

Therefore:  $|f(t,y_1) - f(t,y_2)| = |\cos(c) \cdot (y_1 - y_2)| = |\cos(c)| \cdot |y_1 - y_2| \leq 1 \cdot |y_1 - y_2|$

Since  $|\cos(c)| \leq 1$  for all  $c$ , the function satisfies a Lipschitz condition with Lipschitz constant  $L = 1$  on the given domain.

##### Problem 2: Finding the Interval of Existence

Problem: Consider the initial value problem  $y' = y^2$ ,  $y(0) = 1$ . Determine the interval where the solution exists and is unique.

Solution: First, let's verify that  $f(t,y) = y^2$  satisfies the conditions for existence and uniqueness:

1.  $f(t,y) = y^2$  is continuous for all  $(t,y)$ .
2. For any bounded domain where  $|y| \leq M$ ,  $f$  satisfies a Lipschitz condition with respect to  $y$ :  $|f(t,y_1) - f(t,y_2)| = |y_1^2 - y_2^2| = |y_1 - y_2| \cdot |y_1 + y_2| \leq 2M \cdot |y_1 - y_2|$

So, the solution exists and is unique locally. To find the interval of existence, we need to solve the equation:

$$y' = y^2, y(0) = 1$$

This is a separable equation:  $dy/y^2 = dt$   $-1/y = t + C$

Using the initial condition  $y(0) = 1$ :  $-1/1 = 0 + C$   $C = -1$

Therefore:  $-1/y = t - 1$   $y = -1/(t - 1)$

This solution is defined for all  $t$  except  $t = 1$ , where the solution becomes infinite. Therefore, the solution exists and is unique on the interval  $(-\infty, 1)$ .

The reason the solution doesn't extend beyond  $t = 1$  is that it experiences a finite-time blow-up at that point, showing that even when local existence and uniqueness are guaranteed, the solution may not exist globally.

### Problem 3: Method of Successive Approximations

Problem: Use the method of successive approximations to find the first three approximations to the solution of the initial value problem  $y' = t + y$ ,  $y(0) = 1$ .

**Solution: We'll apply Picard's iteration:**

$\varphi_0(t) = 1$  (the initial condition)

$$\begin{aligned}\varphi_1(t) &= 1 + \int_0^t [s + \varphi_0(s)] ds = 1 + \int_0^t [s + 1] ds = 1 + \int_0^t [s + 1] ds \\ &= 1 + [s^2/2 + s] \int_0^t = 1 + (t^2/2 + t) = 1 + t + t^2/2\end{aligned}$$

$$\begin{aligned}\varphi_2(t) &= 1 + \int_0^t [s + \varphi_1(s)] ds = 1 + \int_0^t [s + (1 + s + s^2/2)] ds = \\ &= 1 + \int_0^t [1 + 2s + s^2/2] ds = 1 + [s + s^2 + s^3/6] \int_0^t = 1 + \\ &+ (t + t^2 + t^3/6) = 1 + t + t^2 + t^3/6\end{aligned}$$

$$\begin{aligned}\varphi_3(t) &= 1 + \int_0^t [s + \varphi_2(s)] ds = 1 + \int_0^t [s + (1 + s + s^2 + s^3/6)] ds = \\ &= 1 + \int_0^t [1 + 2s + s^2 + s^3/6] ds = 1 + [s + s^2 + s^3/3 + s^4/24] \int_0^t = 1 + \\ &+ (t + t^2 + t^3/3 + t^4/24) = 1 + t + t^2 + t^3/3 + t^4/24\end{aligned}$$

The exact solution to this linear equation is  $y(t) = 2e^t - t - 1$ , which can be expanded as:  $y(t) = 2(1 + t + t^2/2 + t^3/6 + t^4/24 + \dots) - t - 1 = 1 + t + t^2 + t^3/3 + t^4/12 + \dots$

We can see that our approximations are approaching this series expansion.

#### Problem 4: Analyzing Uniqueness Failure

Problem: Consider the initial value problem  $y' = y^{(2/3)}$ ,  $y(0) = 0$ . Show that this problem has multiple solutions despite  $f(t,y) = y^{(2/3)}$  being continuous.

Solution: The function  $f(t,y) = y^{(2/3)}$  is indeed continuous for all  $(t,y)$ . However, it fails to satisfy the Lipschitz condition at  $y = 0$ . To see this, note that the derivative:

$$\partial f / \partial y = (2/3)y^{(-1/3)}$$

becomes unbounded as  $y$  approaches 0.

Let's now show that multiple solutions exist:

1. The constant function  $y_1(t) = 0$  for all  $t$  is clearly a solution, as  $y_1'(t) = 0 = 0^{(2/3)}$ .
2. Let's try to find another solution. For  $y \neq 0$ , we can separate variables:  $dy/y^{(2/3)} = dt \int y^{(-2/3)} dy = \int dt \ 3y^{(1/3)} = t + C$

If we want a solution that satisfies  $y(0) = 0$ , then:  $3 \cdot 0^{(1/3)} = 0 + C$  This gives us  $C = 0$  (if we interpret  $0^{(1/3)}$  as 0).

Therefore:  $3y^{(1/3)} = t \ y^{(1/3)} = t/3 \ y(t) = (t/3)^3 = t^3/27$  for  $t \geq 0$

3. We can now construct a family of solutions:  $y(t) = \{ 0, \text{ for } t \leq a \ (t-a)^3/27, \text{ for } t > a \}$  where  $a \geq 0$  is an arbitrary parameter.

Each of these functions satisfies the differential equation and the initial condition  $y(0) = 0$ , demonstrating that uniqueness fails in this case. The failure occurs precisely because the Lipschitz condition is not satisfied at the point of interest.

#### Problem 5: Global vs. Local Existence

Problem: For the initial value problem  $y' = y^2$ ,  $y(0) = 1$ , determine: a) The interval where local existence and uniqueness are guaranteed by Picard's theorem b) The actual interval of existence for the solution

**Solution:**

- a) By Picard's theorem, if  $f(t,y) = y^2$  is continuous and satisfies a Lipschitz condition in a rectangle  $R = \{(t,y) : |t - 0| \leq a, |y - 1| \leq b\}$ , then there exists a



unique solution in an interval  $|t| \leq h$ , where  $h = \min(a, b/M)$  and  $M$  is a bound for  $|f(t,y)|$  in  $R$ .

Let's choose  $a = 1/4$  and  $b = 1/2$ . Then  $R = \{(t,y) : |t| \leq 1/4, 1/2 \leq y \leq 3/2\}$ .

In this rectangle:

- $f(t,y) = y^2$  is continuous
- $|f(t,y)| = y^2 \leq (3/2)^2 = 9/4$ , so  $M = 9/4$
- $f$  satisfies a Lipschitz condition with respect to  $y$ :  $|f(t,y_1) - f(t,y_2)| = |y_1^2 - y_2^2| = |y_1 - y_2| \cdot |y_1 + y_2|$ . In  $R$ ,  $|y_1 + y_2| \leq 3$ , so  $L = 3$  is a Lipschitz constant.

Therefore, Picard's theorem guarantees existence and uniqueness in the interval  $|t| \leq h$ , where:  $h = \min(1/4, (1/2)/(9/4)) = \min(1/4, 2/9) = 2/9$

So local existence and uniqueness are guaranteed on  $[-2/9, 2/9]$ .

b) As shown in Problem 2, the actual solution is  $y(t) = -1/(t - 1)$ . This solution exists and is unique on the interval  $(-\infty, 1)$ .

This illustrates an important point: Picard's theorem provides sufficient conditions for local existence and uniqueness, but the actual interval of existence may be larger than what the theorem guarantees.

## Unsolved Problems

### Problem 1

Determine whether the function  $f(t,y) = \ln(t + y^2)$  satisfies a Lipschitz condition with respect to  $y$  on the domain  $D = \{(t,y) : t \geq 1, -2 \leq y \leq 2\}$ .

### Problem 2

Consider the initial value problem  $y' = t \cdot y / (1 + y^2)$ ,  $y(0) = 0$ . Determine whether the solution to this problem is unique, and explain your reasoning using the appropriate theorems.

### Problem 3

Use the method of successive approximations to find the first three approximations to the solution of the initial value problem  $y' = t \cdot y$ ,  $y(0) = 2$ .

### Problem 4

For the initial value problem  $y' = \sqrt{|y|}$ ,  $y(0) = 0$ : a) Determine whether the hypotheses of the existence and uniqueness theorem are satisfied b) Find all possible solutions to this problem

### Problem 5

Consider a nonlinear spring-mass system modeled by the differential equation:  $m \cdot y'' + c \cdot y' + k \cdot y + \alpha \cdot y^3 = 0$  where  $m$ ,  $c$ ,  $k$ , and  $\alpha$  are positive constants. Rewrite this as a system of first-order equations and determine conditions on the parameters that guarantee local existence and uniqueness of solutions for any initial conditions  $y(0) = y_0$ ,  $y'(0) = v_0$ .

### Multiple Choice Questions (MCQs)

1. **The existence and uniqueness theorem states that a unique solution exists if:**
  - a) The function and its partial derivative satisfy certain conditions
  - b) The function is continuous everywhere
  - c) The equation has constant coefficients
  - d) None of the above
2. **The method of successive approximations is also known as:**
  - a) Euler's method
  - b) The Picard iteration method
  - c) The Runge-Kutta method
  - d) None of the above
3. **The Lipschitz condition ensures:**
  - a) Uniqueness of the solution
  - b) The solution is periodic
  - c) The solution does not exist
  - d) None of the above
4. **The equation  $y' = y^2 + x$  is an example of:**
  - a) A separable equation
  - b) A linear equation
  - c) A Riccati equation
  - d) None of the above
5. **The Picard-Lindelöf theorem provides conditions for:**
  - a) The uniqueness of solutions

- b) The periodicity of solutions
- c) The non-existence of solutions
- d) None of the above

6. **Convergence of successive approximations ensures:**

- a) A unique solution to the differential equation
- b) No solution exists
- c) The equation is always exact
- d) None of the above

7. Answer Key:

1	a	3	b	5	a
2	b	4	c	6	a

## CHECK YOUR PROGRESS

1. What condition ensures convergence in Picard's method?

.....  
.....  
.....

2. Write the mathematical form of the convergence condition.

.....  
.....  
.....

3. Explain how reducing the interval length affects convergence.

.....  
.....  
.....

4. What happens if  $f(x, y)$  is not Lipschitz continuous?

.....  
.....  
.....

## LET US SUM UP

The convergence of successive approximations refers to the process by which the sequence of functions  $y_n(x)$  obtained through Picard's method approaches the exact solution. If the function  $f(x, y)$  satisfies the Lipschitz condition in a bounded interval, the approximations converge uniformly to a single, continuous solution. The rate and region of convergence depend on the Lipschitz constant and the size of the interval considered.

## UNIT END EXERCISES

### Short Answer Questions:

1. Solve the separable equation  $dy/dx = xy$
2. What role does continuity play in the existence of solutions?
3. Give an application of existence and uniqueness theorems.

### Long Answer Questions:

1. Prove that if  $f(x, y)$  satisfies a Lipschitz condition in  $y$  on a closed rectangle  $R: |x - x_0| \leq a, |y - y_0| \leq b$ , then the sequence of functions defined by  $y_{n+1}(x) = y_0 + \int_{x_0}^x f(t, y_n(t)) dt$  converges uniformly to the unique solution of the differential equation  $y' = f(x, y)$ . Illustrate the result by considering  $f(x, y) = x + y$  with  $y(0) = 1$ .
2. Discuss real-world applications of the existence and uniqueness theorem.

### REFERENCES AND SUGGESTED READINGS:

1. Hartman, P. (2013). Ordinary Differential Equations (2nd ed.). SIAM
2. Coddington, E. A. (2018). An Introduction to Ordinary Differential Equations. Dover Publications.
3. Arnold, V. I. (2012). Ordinary Differential Equations. Springer.
4. Braun, M. (2010). Differential Equations and Their Applications (4th ed.). Springer.
5. Hirsch, M. W., Smale, S., & Devaney, R. L. (2013). Differential Equations, Dynamical Systems, and an Introduction to Chaos (3rd ed.), Academic Press.

# **MATS UNIVERSITY**

**MATS CENTRE FOR DISTANCE AND ONLINE EDUCATION**

**UNIVERSITY CAMPUS:** Aarang Kharora Highway, Aarang, Raipur, CG, 493 441

**RAIPUR CAMPUS:** MATS Tower, Pandri, Raipur, CG, 492 002

**T : 0771 4078994, 95, 96, 98 Toll Free ODL MODE : 81520 79999, 81520 29999**

**Website:** [www.matsodl.com](http://www.matsodl.com)

